

# Discussion Papers in Economics

## Cost Monotonicity, Consistency and Minimum Cost spanning Tree Games.

Bhaskar Dutta and Anirban Kar

**Discussion Paper 02-04**

**July 2002**



Indian Statistical Institute, Delhi  
Planning Unit  
7 S.J.S. Sansanwal Marg, New Delhi 110 016, India

# Cost Monotonicity, Consistency and Minimum Cost Spanning Tree Games

Bhaskar Dutta and Anirban Kar\*

July 2001

This Version : July 2002

---

\*Dutta is at the University of Warwick, Coventry CV4 7AL, England, (B.Dutta@warwick.ac.uk). Kar is at the Indian Statistical Institute, New Delhi - 110 016, India, (anirban@isid.ac.in). We are most grateful to two anonymous referees and an Associate Editor for remarkably detailed comments on an earlier version of the paper. We also acknowledge helpful suggestions from A.van den Nouweland and H. Moulin.

## **Abstract**

We propose a new cost allocation rule for minimum cost spanning tree games. The new rule is a core selection and also satisfies cost monotonicity. We also give characterization theorems for the new rule as well as the much-studied Bird allocation. We show that the principal difference between these two rules is in terms of their consistency properties. JEL Classification Numbers: D7

*Keywords:* spanning tree, cost allocation, core selection, cost monotonicity, consistency.

# 1 Introduction

There is a wide range of economic contexts in which “aggregate costs” have to be allocated amongst individual agents or components who derive the benefits from a common project. A firm has to allocate overhead costs amongst its different divisions. Regulatory authorities have to set taxes or fees on individual users for a variety of services. Partners in a joint venture must share costs (and benefits) of the joint venture. In many of these examples, there is no external force such as the market, which determines the allocation of costs. Thus, the final allocation of costs is decided either by mutual agreement or by an “arbitrator” on the basis of some notion of *fairness*.

A central problem of cooperative game theory is how to divide the benefits of cooperation amongst individual players or agents. Since there is an obvious analogy between the division of costs and that of benefits, the tools of cooperative game theory have proved very useful in the analysis of cost allocation problems.<sup>1</sup> Much of this literature has focused on “general” cost allocation problems, so that the ensuing *cost game* is identical to that of a typical game in characteristic function form. This has facilitated the search for “appropriate” cost allocation rules considerably given the corresponding results in cooperative game theory.

The purpose of this paper is the analysis of allocation rules in a special class of cost allocation problems. The common feature of these problems is that a group of users have to be connected to a single supplier of some service. For instance, several towns may draw power from a common power plant, and hence have to share the cost of the distribution network. There is a non-negative cost of connecting each pair of users (towns) as well as a cost of connecting each user (town) to the common supplier (power plant). A cost game arises because cooperation reduces aggregate costs - it may be cheaper for town A to construct a link to town B which is “nearer” to the power plant, rather than build a separate link to the plant. Clearly, an efficient network must be a *tree*, which connects all users to the common supplier. That is

---

<sup>1</sup>Moulin[11] and Young [16] are excellent surveys of this literature.

why these games have been labelled *minimum cost spanning tree games*.

Notice that in the example mentioned above, it makes sense for town B to demand some compensation from A in order to let A use its own link to the power plant. But, how much should A agree to pay? This is where both strategic issues as well as considerations of fairness come into play. Of course, these issues are present in *any* surplus-sharing or cost allocation problem. What is special in our context is that the structure of the problem implies that the *domain* of the allocation rule will be smaller than that in a more general cost problem. This smaller domain raises the possibility of constructing allocation rules satisfying “nice” properties which cannot always be done in general problems. For instance, it is known that the *core* of a minimum cost spanning tree game is always non-empty.<sup>2</sup>

Much of the literature on minimum cost spanning tree games has focused on *algorithmic* issues.<sup>3</sup> In contrast, the derivation of attractive cost allocation rules or the analysis of axiomatic properties of different rules has received correspondingly little attention.<sup>4</sup> This provides the main motivation for this paper. We show that the allocation rule proposed by Bird [1], which always selects an allocation in the core of the game, does not satisfy *cost monotonicity*. Cost monotonicity is an extremely attractive property, and requires that the cost allocated to agent  $i$  does not increase if the cost of a link involving  $i$  goes down, nothing else changing. Notice that if a rule does not satisfy cost monotonicity, then it may not provide agents with the appropriate incentives to reduce the costs of constructing links.

The cost allocation rule, which coincides with the Shapley value of the cost game, satisfies cost monotonicity. However, the Shapley value is unlikely to be used in these contexts because it may not lie in the core. This implies that some group of agents may well find it beneficial to construct their own network if the Shapley value is used to allocate costs. We show that cost monotonicity and the core are not mutually exclusive<sup>5</sup> by constructing a new rule, which satisfies cost monotonicity and also

---

<sup>2</sup>See, for instance, Bird[1], Granot and Huberman [7].

<sup>3</sup>See for instance Granot and Granot [5], Granot and Huberman [6], Graham and Hell [4].

<sup>4</sup>Exceptions are Feltkampf [3], Kar [10]. See Sharkey [14] for a survey of this literature.

<sup>5</sup>This is where the small domain comes in useful. Young [16] shows that in the context of

selects an allocation in the core of the game.

We then go on to provide axiomatic characterizations of the Bird rule as well as the new rule constructed by us. An important type of axiom used by us is closely linked to the *reduced game properties* which have been extensively used in the axiomatic characterization of solutions in cooperative game theory.<sup>6</sup> These are consistency conditions, which place restrictions on how solutions of different but related games defined on different player sets behave. We show that the Bird rule and the new allocation rule satisfy different consistency conditions.

The plan of this paper is the following. In section 2, we define the basic structure of minimum cost spanning tree games. The main purpose of Section 3 is the construction of the new rule as well as the proof that it satisfies cost monotonicity and core selection. Section 4 contains the characterization results. An appendix contains the proofs of some lemmas.

## 2 The Framework

Let  $\mathcal{N} = \{1, 2, \dots\}$  be the set of all possible agents. We are interested in *graphs* or *networks* where the nodes are elements of a set  $N \cup \{0\}$ , where  $N \subset \mathcal{N}$ , and 0 is a distinguished node which we will refer to as the *source* or *root*.

Henceforth, for any set  $N \subset \mathcal{N}$ , we will use  $N^+$  to denote the set  $N \cup \{0\}$ .

A typical graph over  $N^+$  will be represented by  $g_N = \{(ij) | i, j \in N^+\}$ . Two nodes  $i$  and  $j \in N^+$  are said to be *connected* in  $g_N$  if  $\exists (i_1 i_2), (i_2 i_3), \dots, (i_{n-1} i_n)$  such that  $(i_k i_{k+1}) \in g_N$ ,  $1 \leq k \leq n-1$ , and  $i_1 = i, i_n = j$ . A graph  $g_N$  is called *connected* over  $N^+$  if  $i, j$  are connected in  $g_N$  for all  $i, j \in N^+$ . The set of connected graphs over  $N^+$  is denoted by  $\Gamma_N$ .

Consider any  $N \subset \mathcal{N}$ , where  $\#N = n$ . A *cost matrix*  $C = (c_{ij})$  represents the cost of *direct* connection between any pair of nodes. That is,  $c_{ij}$  is the cost of directly transferable utility games, there is no solution concept which picks an allocation in the core of the game when the latter is nonempty and also satisfies a property which is analogous to cost monotonicity.

<sup>6</sup>See Peleg[12], Thomson [15].

connecting any pair  $i, j \in N^+$ . We assume that each  $c_{ij} > 0$  whenever  $i \neq j$ . We also adopt the convention that for each  $i \in N^+$ ,  $c_{ii} = 0$ . So, each cost matrix over  $N^+$  is nonnegative, symmetric and of order  $n + 1$ . In this paper we will often use the term matrix instead of cost matrix. The set of all matrices for  $N$  is denoted by  $\mathcal{C}_N$ . However, we will typically drop the subscript  $N$  whenever there is no cause for confusion about the set of nodes.

Consider any  $C \in \mathcal{C}_N$ . A *minimum cost spanning tree* (m.c.s.t.) over  $N^+$  satisfies  $g_N = \operatorname{argmin}_{g \in \Gamma_N} \sum_{(ij) \in g} c_{ij}$ . Note that an m.c.s.t. need not be unique. Clearly a minimum cost spanning network must be a tree. Otherwise, we can delete an extra edge and still obtain a connected graph at a lower cost.

An m.c.s.t. corresponding to  $C \in \mathcal{C}_N$  will typically be denoted by  $g_N(C)$ .

**Example 1:** Consider a set of three rural communities  $\{A, B, C\}$ , which have to decide whether to build a system of irrigation channels to an existing dam, which is the source. Each community has to be *connected* to the dam in order to draw water from the dam. However, some connection(s) could be indirect. For instance, community  $A$  could be connected directly to the dam, while  $B$  and  $C$  are connected to  $A$ , and hence indirectly to the source.

There is a cost of building a channel connecting each pair of communities, as well as a channel connecting each community directly to the dam. Suppose, these costs are represented by the matrix  $C$  given below.

$$C = \begin{pmatrix} 0 & 2 & 4 & 1 \\ 2 & 0 & 1 & 3 \\ 4 & 1 & 0 & 2 \\ 1 & 3 & 2 & 0 \end{pmatrix}$$

The minimum cost of building the system of irrigation channels will be 4 units. Our object of interest in this paper is to see how the total cost of 4 units is to be distributed amongst  $A, B$  and  $C$ .

This provides the motivation for the next definition.

**Definition 1:** A *cost allocation rule* (or simply a *rule*) is a family of functions

$\{\psi^N\}_{N \subset \mathcal{N}}$  with  $N \subset \mathcal{N}$ ,  $\psi^N : \mathcal{C}_N \rightarrow \mathfrak{R}_+^N$  satisfying  $\sum_{i \in N} \psi_i^N(C) \geq \sum_{(ij) \in g_N(C)} c_{ij}$  for all  $C \in \mathcal{C}_N$ .

We will drop the superscript  $N$  for the rest of the paper.

So, given any set of nodes  $N$  and any matrix  $C$  of order  $(|N| + 1)$ , a rule specifies the costs attributed to agents in  $N$ . Note that the source 0 is not an *active* player, and hence does not bear any part of the cost.

A rule can be generated by any *single-valued* game-theoretic solution of a transferable utility game. Thus, consider the transferable utility game generated by considering the aggregate cost of a minimum cost spanning tree for each coalition  $S \subseteq N$ . Given  $C$  and  $S \subseteq N$ , let  $C_S$  be the matrix restricted to  $S^+$ . Then, consider a m.c.s.t.  $g_S(C_S)$  over  $S^+$ , and the corresponding minimum cost of connecting  $S$  to the source. Let this cost be denoted by  $c_S$ . For each  $N \subset \mathcal{N}$ , this defines a *cost game*  $(N, c)$  where for each  $S \subseteq N$ ,  $c(S) = c_S$ . That is,  $c$  is the cost function, and is analogous to a TU game. Then, if  $\Phi$  is a single-valued solution,  $\Phi(N, c)$  can be viewed as the rule corresponding to the matrix which generates the cost function  $c$ .<sup>7</sup>

One particularly important game-theoretic property, which will be used subsequently is that of the *core*. If a rule does not always pick an element in the core of the game, then some subset of  $N$  will find it profitable to break up  $N$  and construct its own minimum cost tree. This motivates the following definition.

**Definition 2:** A rule  $\phi$  is a *core selection* if for all  $N \subseteq \mathcal{N}$  and for all  $C \in \mathcal{C}_N$ ,  $\sum_{i \in S} \phi_i(C) \leq c(S)$ , where  $c(S)$  is the cost of the m.c.s.t. for  $S$ ,  $\forall S \subseteq N$ .

However, cost allocation rules can also be defined without appealing to the underlying cost game. For instance, this was the procedure followed by Bird [1]. In order to describe his procedure, we need some more notation.

The (unique) *path* from  $i$  to  $j$  in tree  $g$ , is a set  $U(i, j, g) = \{i_1, i_2, \dots, i_K\}$ , where each pair  $(i_{k-1}, i_k) \in g$ , and  $i_1, i_2, \dots, i_K$  are all distinct agents with  $i_1 = i, i_K = j$ . The *predecessor set* of an agent  $i$  in  $g$  is defined as  $P(i, g) = \{k | k \neq i, k \in U(0, i, g)\}$ . The *immediate predecessor* of agent  $i$ , denoted by  $\alpha(i)$ , is the agent who

<sup>7</sup>See Kar [10] for an axiomatic characterization of the Shapley value in m.c.s.t. games.

comes immediately before  $i$ , that is,  $\alpha(i) \in P(i, g)$  and  $k \in P(i, g)$  implies either  $k = \alpha(i)$  or  $k \in P(\alpha(i), g)$ .<sup>8</sup> The *followers* of agent  $i$ , are those agents who come immediately after  $i$ ;  $F(i) = \{j | \alpha(j) = i\}$ .

Bird's method is defined with respect to a *specific* tree. Let  $g_N$  be some m.c.s.t. corresponding to the matrix  $C$ . Then,

$$B_i(C) = c_{i\alpha(i)} \forall i \in N.$$

So, in the Bird allocation, each node pays the cost of connecting to its immediate predecessor in the appropriate m.c.s.t.

Notice that this does not define a rule if  $C$  gives rise to more than one m.c.s.t. However, when  $C$  does not induce a unique m.c.s.t., one can still use Bird's method on each m.c.s.t. derived from  $C$  and then take some convex combination of the allocations corresponding to each m.c.s.t. as the rule. In general, the properties of the resulting rule will not be identical to those of the rule given by Bird's method on matrices which induce unique m.c.s.t. s.

In section 4, we will use two domain restrictions on the set of permissible matrices. These are defined below.

**Definition 3:**  $\mathcal{C}^1 = \{C \in \mathcal{C} | C \text{ induces a } \textit{unique} \text{ m.c.s.t. } \forall N \subset \mathcal{N}\}$ .

**Definition 4:**  $\mathcal{C}^2 = \{C \in \mathcal{C}^1 | \text{no two edges of the unique m.c.s.t. have the same cost}\}$ .

Notice if  $C$  is not in  $\mathcal{C}^2$ , then even a "small" perturbation of  $C$  produces a matrix with the property that no two edges have the same cost. Of course, such a matrix must be in  $\mathcal{C}^2$ . So, even the stronger domain restriction is relatively mild, and the permissible sets of matrices are large.

### 3 Cost Monotonicity

The Bird allocation is an attractive rule because it is a core selection. In addition, it is easy to compute. However, it fails to satisfy *cost monotonicity*.

---

<sup>8</sup>Note that since  $g$  is a tree, the immediate predecessor must be unique.

**Definition 5:** Fix  $N \subset \mathcal{N}$ . Let  $i, j \in N^+$ , and  $C, C' \in \mathcal{C}_N$  be such that  $c_{kl} = c'_{kl}$  for all  $(kl) \neq (ij)$  and  $c_{ij} > c'_{ij}$ . Then, the rule  $\psi$  satisfies *Cost Monotonicity* if for all  $m \in N \cap \{i, j\}$ ,  $\psi_m(C) \geq \psi_m(C')$ .

Cost monotonicity is an extremely appealing property. The property applies to two matrices which differ only in the cost of connecting the pair  $(ij)$ ,  $c'_{ij}$  being lower than  $c_{ij}$ . Then, cost monotonicity requires that no agent in the pair  $\{i, j\}$  be charged more when the matrix changes from  $C$  to  $C'$ .

Despite its intuitive appeal, cost monotonicity has a lot of bite.<sup>9</sup> The following example shows that the Bird rule does not satisfy cost monotonicity.

**Example 2:** Let  $N = \{1, 2\}$ . The two matrices are specified below.

- (i)  $c_{01} = 4, c_{02} = 4.5, c_{12} = 3$ .
- (ii)  $c'_{01} = 4, c'_{02} = 3.5, c'_{12} = 3$ .

Then,  $B_1(C) = 4, B_2(C) = 3$ , while  $B_1(C') = 3, B_2(C') = 3.5$ . So, 2 is charged more when the matrix is  $C'$  although  $c'_{02} < c_{02}$  and the costs of edges involving 1 remain the same.

The rule corresponding to the Shapley value of the cost game does satisfy cost monotonicity. However, it does not always select an outcome which is in the core of the cost game. Our main purpose in this section is to define a new rule which will be a core selection and satisfy cost monotonicity. We are able to do this despite the impossibility result due to Young because of the special structure of minimum cost spanning tree games - these are a *strict* subset of the class of *balanced* games. Hence, monotonicity in the context of m.c.s.t. games is a weaker restriction.

We describe an algorithm whose outcome will be the cost allocation prescribed by the new rule. Our rule is defined for all matrices in  $\mathcal{C}$ . However, in order to economise on notation, we describe the algorithm for a matrix in  $\mathcal{C}^2$ . We then indicate how to construct the rule for all matrices.

---

<sup>9</sup>In fact, Young [16] shows that an analogous property in the context of TU games cannot be satisfied by any solution which selects a core outcome in balanced games.

Fix some  $N \subset \mathcal{N}$ , and choose some matrix  $C \in \mathcal{C}_N^2$ . Also, for any  $A \subset N$ , define  $A_c$  as the complement of  $A$  in  $N^+$ . That is  $A_c = N^+ \setminus A$ .

The algorithm proceeds as follows.

Let  $A^0 = \{0\}$ ,  $g^0 = \emptyset$ ,  $t^0 = 0$ .

*Step 1:* Choose the ordered pair  $(a^1 b^1)$  such that  $(a^1 b^1) = \operatorname{argmin}_{(i,j) \in A^0 \times A_c^0} c_{ij}$ .

Define  $t^1 = \max(t^0, c_{a^1 b^1})$ ,  $A^1 = A^0 \cup \{b^1\}$ ,  $g^1 = g^0 \cup \{(a^1 b^1)\}$ .

*Step k:* Define the ordered pair  $(a^k b^k) = \operatorname{argmin}_{(i,j) \in A^{k-1} \times A_c^{k-1}} c_{ij}$ ,  $A^k = A^{k-1} \cup \{b^k\}$ ,  $g^k = g^{k-1} \cup \{(a^k b^k)\}$ ,  $t^k = \max(t^{k-1}, c_{a^k b^k})$ . Also,

$$\psi_{b^{k-1}}^*(C) = \min(t^{k-1}, c_{a^k b^k}). \quad (1)$$

The algorithm terminates at step  $\#N = n$ . Then,

$$\psi_{b^n}^*(C) = t^n \quad (2)$$

The new rule  $\psi^*$  is described by equations (1), (2).

At any step  $k$ ,  $A^{k-1}$  is the set of nodes which have already been connected to the source 0. Then, a new edge is constructed at this step by choosing the *lowest-cost* edge between a node in  $A^{k-1}$  and nodes in  $A_c^{k-1}$ . The cost allocation of  $b^{k-1}$  is decided at step  $k$ . Equation (1) shows that  $b^{k-1}$  pays the minimum of  $t^{k-1}$ , which is the *maximum* cost amongst all edges which have been constructed in previous steps, and  $c_{a^k b^k}$ , the edge being constructed in step  $k$ . Finally, equation (2) shows that  $b^n$ , the last node to be connected, pays the maximum cost.<sup>10</sup>

**Remark 1:** The algorithm has been described for matrices in  $\mathcal{C}^2$ . Suppose that  $C \notin \mathcal{C}^2$ . Then, the algorithm is not well-defined because at some step  $k$ , two distinct edges  $(a^k b^k)$  and  $(\bar{a}^k \bar{b}^k)$  may minimise the cost of connecting nodes in  $A^{k-1}$  and  $A_c^{k-1}$ . But, there is an easy way to extend the algorithm to deal with matrices not in  $\mathcal{C}^2$ . Let  $\sigma$  be a strict ordering over  $N$ . Then,  $\sigma$  can be used as a tie-breaking rule - for instance, choose  $(a^k b^k)$  if  $b^k$  is ranked over  $\bar{b}^k$  according to  $\sigma$ . Any such tie-breaking rule makes the algorithm well-defined. Now, let  $\Sigma$  be the set of all strict orderings

<sup>10</sup>From Prim[13], it follows that  $g^n$  is also the m.c.s.t. corresponding to  $C$ .

over  $N$ . Then, the eventual cost allocation is obtained by taking the simple average of the “component” cost allocations obtained for each ordering  $\sigma \in \Sigma$ . That is, for any  $\sigma \in \Sigma$ , let  $\psi_\sigma^*(C)$  denote the cost allocation obtained from the algorithm when  $\sigma$  is used as the tie-breaking rule. Then,

$$\psi^*(C) = \frac{1}{\#\Sigma} \sum_{\sigma \in \Sigma} \psi_\sigma^*(C). \quad (3)$$

We illustrate this procedure in Example 5 below.

**Remark 2:** Notice that  $\psi^*$  only depends on the m.c.s.t.s corresponding to any matrix. This property of *Tree Invariance* adds to the computational simplicity of the rule, and distinguishes it from rules such as the Shapley Value and nucleolus.

We now construct a few examples to illustrate the algorithm.

**Example 3:** Suppose  $C^1$  is such that the m.c.s.t. is unique and is a *line*. That is, each node has at most one follower. Then the nodes can be labelled  $a_0, a_1, a_2, \dots, a_n$ , where  $a_0 = 0$ ,  $\#N = n$ , with the predecessor set of  $a_k$ ,  $P(a_k, g) = \{0, a_1, \dots, a_{k-1}\}$ . Then,

$$\forall k < n, \psi_{a_k}^*(C^1) = \min(\max_{0 \leq t < k} c_{a_t a_{t+1}}, c_{a_k a_{k+1}}) \quad (4)$$

and

$$\psi_{a_n}^*(C^1) = \max_{0 \leq t < n} c_{a_t a_{t+1}} \quad (5)$$

**Example 4:** Let  $N = \{1, 2, 3, 4\}$ , and

$$C^2 = \begin{pmatrix} 0 & 4 & 5 & 5 & 5 \\ 4 & 0 & 2 & 1 & 5 \\ 5 & 2 & 0 & 5 & 5 \\ 5 & 1 & 5 & 0 & 3 \\ 5 & 5 & 5 & 3 & 0 \end{pmatrix}$$

There is only one m.c.s.t. of  $C^2$ .

*Step 1:* We have  $(a^1 b^1) = (01)$ ,  $t^1 = c_{01} = 4$ ,  $A^1 = \{0, 1\}$ .

*Step 2:* Next,  $(a^2 b^2) = (13)$ ,  $\psi_1^*(C^2) = \min(t^1, c_{13}) = 1$ ,  $t^2 = \max(t^1, c_{13}) = 4$ ,  $A^2 = \{0, 1, 3\}$ .

*Step 3:* We now have  $(a^3b^3) = (12)$ ,  $\psi_3^*(C^2) = \min(t^2, c_{12}) = 2$ ,  $t^3 = \max(t^2, c_{12}) = 4$ ,  $A^3 = \{0, 1, 2, 3\}$ .

*Step 4:* Next,  $(a^4b^4) = (34)$ ,  $\psi_2^*(C^2) = \min(t^3, c_{34}) = 3$ ,  $t^4 = \max(t^3, c_{34}) = 4$ ,  $A^4 = \{0, 1, 2, 3, 4\}$ .

Since  $A^4 = N^+$ ,  $\psi_4^*(C^2) = t^4 = 4$ , and the algorithm is terminated.

So,  $\psi^*(C^2) = (1, 3, 2, 4)$ . This example shows that it is not necessary for a node to be assigned the cost of its preceding or following edge. Here 2 pays the cost of the edge (34), while 3 pays the cost of the edge (12).<sup>11</sup>

The next example involves a matrix which has more than one m.c.s.t. with edges which cost the same.

**Example 5:** Let  $N = \{1, 2, 3\}$ , and

$$C^3 = \begin{pmatrix} 0 & 4 & 4 & 5 \\ 4 & 0 & 2 & 2 \\ 4 & 2 & 0 & 5 \\ 5 & 2 & 5 & 0 \end{pmatrix}$$

$C^3$  has two m.c.s.t.s -  $g_N = \{(01), (12), (13)\}$  and  $g_N^1 = \{(02), (12), (13)\}$ . The edges (12) and (13) have the same cost.<sup>12</sup>

Suppose the algorithm is first applied to  $g_N$ . Then, we have  $b^1 = 1$ . In step 2,  $a^2 = 1$ , but  $b^2$  can be either 2 or 3. Taking each in turn, we get the vectors  $x^1 = (2, 2, 4)$  and  $x^2 = (2, 4, 2)$ .

Now, consider  $g_N^1$ , which is a line. So, as we have described in Example 3, the resulting cost allocation is  $\hat{x} = (2, 2, 4)$ .

The algorithm will “generate”  $g_N^1$  instead of  $g_N$  for all  $\sigma \in \Sigma$  which ranks 2 over 1. Hence, the “weight” attached to  $g_N^1$  is half. Similarly, the weight attached to  $x^1$  and  $x^2$  must be one-sixth and one-third respectively.

Hence,  $\psi^*(C^3) = (2, \frac{8}{3}, \frac{10}{3})$ .

We now show that  $\psi^*$  is a core selection and also satisfies Cost Monotonicity.

<sup>11</sup>See Figure 1 for the m.c.s.t. corresponding to  $C^2$ .

<sup>12</sup>Figure 2 displays minimum cost spanning trees of  $C^3$ .

**Theorem 1:** The rule  $\psi^*$  satisfies Cost Monotonicity and is a core selection.<sup>13</sup>

## 4 Characterization Theorems

In this section, we present characterizations of the allocation rules  $\psi^*$  and  $B$ .<sup>14</sup> These characterization theorems will be proved for the restricted domains  $\mathcal{C}^1$  for  $B$  and  $\mathcal{C}^2$  for  $\psi^*$ . Examples 8 and 9 explain why we choose these domain restrictions.

We first describe the axioms used in the characterization.

$$\text{Efficiency (EF): } \sum_{i \in N} \psi_i(C) = \sum_{(ij) \in g_N(C)} c_{ij}.$$

This axiom ensures that the agents together pay exactly the cost of the efficient network.

Before moving on to our next axiom, we introduce the concept of an *extreme point*. Let  $C \in \mathcal{C}_N$  be such that the m.c.s.t.  $g_N(C)$  is unique. Then,  $i \in N$  is called an *extreme point* of  $g_N(C)$  (or equivalently of  $C$ ), if  $i$  has no follower in  $g_N(C)$ .

*Extreme Point Monotonicity (EPM):* Let  $C \in \mathcal{C}_N$ , and  $i$  be an extreme point of  $C$ . Let  $\bar{C}$  be the restriction of  $C$  over the set  $N^+ \setminus \{i\}$ . A rule satisfies *Extreme Point Monotonicity* if  $\psi_k(\bar{C}) \geq \psi_k(C) \forall k \in N \setminus \{i\}$ .

Suppose  $i$  is an extreme point of  $g_N(C)$ . Note that  $i$  is of no use to the rest of the network since no node is connected to the source through  $i$ . Extreme Point Monotonicity essentially states that no “existing” node  $k$  will agree to pay a higher cost in order to include  $i$  in the network.

The next two axioms are *consistency* properties, analogous to reduced game properties introduced by Davis and Maschler [2] and Hart and Mas-Collel [9].<sup>15</sup>

<sup>13</sup>All proofs are contained in the appendix.

<sup>14</sup>See Feltkamp [3] for an alternative characterization of  $B$ .

<sup>15</sup>Thomson[15] contains an excellent discussion of consistency properties in various contexts. Granot and Maschler[8] also use a reduced game property for a class of related (but different) cost allocation problems.

Consider any  $C$  with a unique m.c.s.t.  $g_N(C)$ , and suppose that  $(i0) \in g_N(C)$ . Let  $x_i$  be the cost allocation ‘assigned’ to  $i$ . Suppose  $i$  ‘leaves’ the scene (or stops bargaining for a different cost allocation), but other nodes are allowed to connect through it. Then, the effective *reduced matrix* changes for the remaining nodes. We can think of two alternative ways in which the others can use node  $i$ .

- (i) The others can use node  $i$  only to connect to the source.
- (ii) Node  $i$  can be used more widely. That is, node  $j$  can connect to node  $k$  through  $i$ .

In case (i), the connection costs on  $N^+ \setminus \{i\}$  are described by the following equations:

$$\text{For all } j \neq i, \bar{c}_{j0} = \min(c_{j0}, c_{ji} + c_{i0} - x_i) \quad (6)$$

$$\text{If } \{j, k\} \cap \{i, 0\} = \emptyset, \text{ then } \bar{c}_{jk} = c_{jk} \quad (7)$$

Equation 6 captures the notion that node  $j$ ’s cost of connecting to the source is the cheaper of two options - the first option being the original one of connecting directly to the source, while the second is the indirect one of connecting through node  $i$ . In the latter case, the cost borne by  $j$  is adjusted for the fact that  $i$  pays  $x_i$ . Equation 7 captures the notion that node  $i$  can only be used to connect to the source.

Let  $C_{x_i}^{sr}$  represent the reduced matrix derived through equations 6, 7.

Consider now case (ii).

$$\text{For all } j, k \in N^+ \setminus \{i\}, \bar{c}_{jk} = \min(c_{jk}, c_{ji} + c_{ki} - x_i). \quad (8)$$

Equation 8 captures the notion that  $j$  can use  $i$  to connect to any other node  $k$ , where  $k$  is not necessarily the source.

Let  $C_{x_i}^{tr}$  represent the reduced matrix derived through equation 8.

We can now define the two consistency conditions.

*Source Consistency (SR)*: Let  $C \in \mathcal{C}_N^1$ , and  $(0i) \in g_N(C)$ . Then, the rule  $\psi$  satisfies *Source Consistency* if  $\psi_k(C_{\psi_i(C)}^{sr}) = \psi_k(C)$  for all  $k \in N \setminus \{i\}$  whenever  $C_{\psi_i(C)}^{sr} \in \mathcal{C}_{N \setminus i}^1$ .

*Tree Consistency (TR)*: Let  $C \in \mathcal{C}_N^2$ , and  $(0i) \in g_N(C)$ . Then, the rule  $\psi$  satisfies *Tree Consistency* if  $\psi_k(C_{\psi_i^{tr}}^{tr}(C)) = \psi_k(C)$  for all  $k \in N \setminus \{i\}$  whenever  $C_{\psi_i^{tr}}^{tr} \in \mathcal{C}_{N \setminus i}^2$ .

The two consistency conditions require that the cost allocated to any agent be the same on the original and reduced matrix. This ensures that once an agent connected to the source agrees to a particular cost allocation and then subsequently allows other agents to use its location for possible connections, the remaining agents do not have any incentive to reopen the debate about what is an appropriate allocation of costs.

We now present a characterization of  $\psi^*$  in terms of Tree Consistency, Efficiency and Extreme Point Monotonicity.

**Theorem 2** : Over the domain  $\mathcal{C}^2$ , a rule  $\psi$  satisfies TR, EF and EPM if and only if  $\psi = \psi^*$ .

We now show that the three axioms used in the theorem are independent.

**Example 6**: We construct a rule  $\phi$  which satisfies EPM and TR but violates EF.

$$\text{Let } \phi_k(C) = \psi_k^*(C) + \epsilon \forall k, \text{ where } \epsilon > \sum_{(ij) \in g_N(C)} c_{ij}.$$

Since  $\psi^*$  satisfies EPM,  $\phi$  also satisfies EPM. Moreover, the restriction on the value of  $\epsilon$  ensures that the reduced matrices always lie outside  $\mathcal{C}$ . So, TR is vacuously satisfied by  $\phi$ . Also, since  $\sum_{k=1}^n \phi_k(C) = \sum_{k=1}^n \psi_k^*(C) + n\epsilon > c(N)$ ,  $\phi$  violates EF. ■

To construct the next example we need to define the concept of an *m.c.s.t. partition*.

Given  $C$ , let  $g_N(C)$  be the (unique) m.c.s.t. of  $C$ . Suppose  $g_N(C) = g_{N_1} \cup g_{N_2} \dots \cup g_{N_K}$ , where each  $g_{N_k}$  is the m.c.s.t. on  $N_k$  for the matrix  $C$  restricted to  $N_k^+$ , with  $\cup_{k=1}^K N_k = N$  and  $N_i \cap N_j = \emptyset$ . We will call such a partition the m.c.s.t. partition of  $N$ .

**Example 7**: We now construct a rule  $\mu$  which satisfies EF and TR, but does not satisfy EPM.

Let  $N = [N_1, \dots, N_T]$  be the m.c.s.t. partition and  $\#N_t = n_t$ . Let  $C^t$  be the restriction of  $C$  over  $N_t^+$ . First, calculate  $\psi^*$  separately for each  $C^t$ . Consider any  $N_t$ . If  $n_t = 1$ ,  $\mu_k(C) = c_{k0}$  where  $k \in N_t$ . For  $n_t \geq 2$ ,

$$(i) \mu_k(C) = \psi_k^*(C^t) \forall k \neq b^{n_t-1}, b^{n_t}.$$

$$(ii) \mu_{b^{n_t-1}}(C) = \psi_{b^{n_t-1}}^*(C^t) + \psi_{b^{n_t}}^*(C^t) \text{ and } \mu_{b^{n_t}}(C) = 0.$$

EF is obviously satisfied. If  $n_t > 2$ ,  $\mu$  satisfies TR because  $\psi^*$  satisfies TR. If  $n_t = 2$  then TR is vacuously satisfied as the reduced matrix lies outside  $\mathcal{C}$  because the reduced cost of connecting the singleton node to the source will be 0. But this allocation violates EPM. In order to see the latter, consider the following matrix  $C$ .

$$C = \begin{pmatrix} 0 & 1 & 3 \\ 1 & 0 & 2 \\ 3 & 2 & 0 \end{pmatrix}.$$

Then,  $g_N(C) = \{(01), (12)\}$ . Clearly, 2 is an extreme point of  $C$ . Let  $\bar{C}$  be the restriction of  $C$  over  $\{0,1\}$ . Then,  $\mu_1(C) = 3 > 1 = \mu_1(\bar{C})$  and hence EPM is violated.

We remark in the next theorem that the Bird rule  $B$  satisfies EF and EPM. Since  $B \neq \psi^*$ , it follows that  $B$  does not satisfy TR. Here is an explicit example to show that  $B$  violates TR.

$$C = \begin{pmatrix} 0 & 2 & 3.5 & 3 \\ 2 & 0 & 1.5 & 1 \\ 3.5 & 1.5 & 0 & 2.5 \\ 3 & 1 & 2.5 & 0 \end{pmatrix}$$

Then,  $B_1(C) = 2, B_2(C) = 1.5$  and  $B_3(C) = 1$ . The reduced matrix is  $C_{B_1(C)}^{tr}$  is shown below.

$$C_{B_1(C)}^{tr} = \begin{pmatrix} 0 & 1.5 & 1 \\ 1.5 & 0 & 0.5 \\ 1 & 0.5 & 0 \end{pmatrix}.$$

Then,  $B_2(C_{B_1(C)}^{tr}) = 0.5$  and  $B_3(C_{B_1(C)}^{tr}) = 1$ . Therefore TR is violated.

However,  $B$  does satisfy Source Consistency on the domain  $\mathcal{C}^1$ . In fact, we now show that  $B$  is the only rule satisfying EF, EPM and SR.

**Theorem 3 :** Over the domain  $\mathcal{C}^1$ , a rule  $\phi$  satisfies SR, EF and EPM iff  $\phi = B$ .

We now show that the three axioms used in Theorem 3 are independent.

A rule which violates EF but satisfies SR and EPM can be constructed using example 6,  $\psi^*$  being replaced by  $B$ .

The rule obtained by replacing  $\psi^*$  with  $B$  in example 7, violates EPM but satisfies EF and SR.

Finally,  $\psi^*$  satisfies all the axioms but SR. Here is an example to show that our rule may violates SR.

$$C = \begin{pmatrix} 0 & 2 & 3 & 4 \\ 2 & 0 & 1.5 & 1 \\ 3 & 1.5 & 0 & 3.5 \\ 4 & 1 & 3.5 & 0 \end{pmatrix}$$

Then,  $\psi_1^*(C) = 1$ ,  $\psi_2^*(C) = 2$  and  $\psi_3^*(C) = 1.5$ . The reduced matrix is  $\widehat{C}$ ,

$$\widehat{C} = \begin{pmatrix} 0 & 2.5 & 2 \\ 2.5 & 0 & 3.5 \\ 2 & 3.5 & 0 \end{pmatrix}$$

$\psi_2^*(\widehat{C}) = 2.5$  and  $\psi_3^*(\widehat{C}) = 2$ . Therefore SR is violated.

In Theorem 2, we have restricted attention to matrices in  $\mathcal{C}^2$ . This is because  $\psi^*$  does not satisfy TR outside  $\mathcal{C}^2$ . The next example illustrates.

**Example 8:** Consider

$$C = \begin{pmatrix} 0 & 3 & 4 & 3 \\ 3 & 0 & 2 & 5 \\ 4 & 2 & 0 & 1 \\ 3 & 5 & 1 & 0 \end{pmatrix}$$

Then,  $g_N^1(C) = \{(10), (12), (23)\}$  and  $g_N^2(C) = \{(30), (32), (21)\}$  are the two m.c.s.t.s corresponding to  $C$ . Taking the average of the two cost allocations derived from the algorithm, we get  $\psi^*(C) = (2.5, 1.5, 2)$ . If we remove 1, which is connected to 0 in  $g_N^1$ , the reduced matrix  $\widehat{C}$  is:

$$\widehat{C} = \begin{pmatrix} 0 & 2.5 & 3 \\ 2.5 & 0 & 1 \\ 3 & 1 & 0 \end{pmatrix}$$

Then,  $\psi_2^*(\widehat{C}) = 1$  and  $\psi_3^*(\widehat{C}) = 2.5$ . So, TR is violated.

**Remark 3:** Note that in the previous example  $C$  lies outside  $\mathcal{C}^1$ . If we take a matrix in  $\mathcal{C}^1 \setminus \mathcal{C}^2$ , then Lemma 2 will no longer be valid - the reduced matrix may lie outside  $\mathcal{C}^1$  even when a node connected to the source pays the minimum cost amongst all its links. Thus,  $\psi^*$  will satisfy TR vacuously. But there may exist allocation rules other than  $\psi^*$  which satisfies EF, TR and EPM over  $\mathcal{C}^1$ .

Similarly,  $B$  does not satisfy SR outside  $\mathcal{C}^1$ .

**Example 9:** Consider the same matrix as in Example 8. Recall that  $B(C) = (2.5, 1.5, 2)$ .

If we remove 1, which is connected to 0 in  $g_N^1$ , the reduced matrix  $\widehat{C}$  is:

$$\widehat{C} = \begin{pmatrix} 0 & 2.5 & 3 \\ 2.5 & 0 & 1 \\ 3 & 1 & 0 \end{pmatrix}$$

Then,  $B_2(\widehat{C}) = 2.5$  and  $B_3(\widehat{C}) = 1$ . Therefore SR is violated.

**Remark 4:** An interesting open question is the characterization of  $\psi^*$  using cost monotonicity and other axioms.

## Appendix

**Proof of Theorem 1:** We first show that  $\psi^*$  satisfies Cost Monotonicity.

Fix any  $N \subset \mathcal{N}$ . We give our proof for matrices in  $\mathcal{C}^2$ , and then indicate how the proof can be extended to cover all matrices. Let  $C, \bar{C} \in \mathcal{C}^2$  be such that for some  $i, j \in N^+$ ,  $c_{ij} > \bar{c}_{ij}$ , and  $c_{kl} = \bar{c}_{kl}$  for all other pairs  $(kl)$ . We need to show that  $\psi_k^*(C) \geq \psi_k^*(\bar{C})$  for  $k \in N \cap \{i, j\}$ .

In describing the algorithm which is used in constructing  $\psi^*$ , we fixed a specific matrix, and so did not have to specify the dependence of  $A^k, t^k, a^k, b^k$  etc. on the matrix. But, now we need to distinguish between these entities for the two matrices  $C$  and  $\bar{C}$ . We adopt the following notation in the rest of the proof of the theorem. Let  $A^k, t^k, a^k, b^k, g_N$  etc. refer to the matrix  $C$ , while  $\bar{A}^k, \bar{t}^k, \bar{a}^k, \bar{b}^k, \bar{g}_N$  etc. will denote the entities corresponding to  $\bar{C}$ .

**Case 1:**  $(ij) \notin \bar{g}_N$ .

Then,  $\bar{g}_N = g_N$ . Since the cost of all edges in  $g_N$  remain the same,  $\psi_k^*(\bar{C}) = \psi_k^*(C)$  for all  $k \in N$ .

**Case 2:**  $(ij) \in \bar{g}_N$ .

Without loss of generality, let  $i$  be the immediate predecessor of  $j$  in  $\bar{g}_N$ . Since the source never pays anything, we only consider the case where  $i$  is not the source.

Suppose  $i = \bar{b}^{k-1}$ . As the cost of all other edges remain the same,  $A^{k-1} = \bar{A}^{k-1}$  and  $t^{k-1} = \bar{t}^{k-1}$ . Now,  $\psi_i^*(\bar{C}) = \min(\bar{t}^{k-1}, \bar{c}_{\bar{a}^k \bar{b}^k})$  and  $\psi_i^*(C) = \min(t^{k-1}, c_{a^k b^k})$ . Since  $\bar{c}_{\bar{a}^k \bar{b}^k} \leq c_{a^k b^k}$ ,  $\psi_i^*(\bar{C}) \leq \psi_i^*(C)$ .

We now show that  $\psi_j^*(\bar{C}) \leq \psi_j^*(C)$ . Let  $j = b^l$  and  $j = \bar{b}^m$ . Note that  $l \geq m$ , and that  $\bar{A}^m \subseteq A^l$ , and  $t^l \geq \bar{t}^m$ .

Now,  $\psi_j^*(\bar{C}) = \min(\bar{t}^m, \bar{c}_{\bar{a}^{m+1} \bar{b}^{m+1}})$ , while  $\psi_j^*(C) = \min(t^l, c_{a^{l+1} b^{l+1}})$ . Since  $t^l \geq \bar{t}^m$ , we only need to show that  $\bar{c}_{\bar{a}^{m+1} \bar{b}^{m+1}} \leq c_{a^{l+1} b^{l+1}}$ .

**Case 2(a):** Suppose  $a^{l+1} \in \bar{A}^m$ . Since  $b^{l+1} \in N^+ \setminus \bar{A}^m$ ,  $\bar{c}_{\bar{a}^{m+1} \bar{b}^{m+1}} \leq \bar{c}_{a^{l+1} b^{l+1}} \leq c_{a^{l+1} b^{l+1}}$ .

**Case 2(b):** Suppose  $a^{l+1} \notin \bar{A}^m$ . Then,  $a^{l+1} \neq j$ . Also,  $a^{l+1} \in A^l$ , and so

$$c_{a^{l+1}b^{l+1}} \geq c_{a^l b^l} \quad (9)$$

We need to consider two sub-cases.

**Case 2(bi):**  $a^l \in A^{l-1} \setminus \bar{A}^{m-1}$ .

Then, since  $A^l = A^{l-1} \cup \{j\}$  and  $\bar{A}^m = \bar{A}^{m-1} \cup \{j\}$ ,  $a^l \in A^l \setminus \bar{A}^m$ .

Now since  $j \in \bar{A}^m$  and  $a^l \notin \bar{A}^m$ ,  $\bar{c}_{\bar{a}^{m+1}\bar{b}^{m+1}} \leq \bar{c}_{ja^l} \leq c_{ja^l} = c_{a^l b^l}$ . Using equation 9,  $c_{a^{l+1}b^{l+1}} \geq c_{a^l b^l} \geq \bar{c}_{\bar{a}^{m+1}\bar{b}^{m+1}}$ .

**Case 2(bii):**  $a^l \in \bar{A}^{m-1} = A^{m-1}$ .

Then,  $c_{a^l b^l} \geq c_{a^m b^m}$  since  $m \leq l$ .

Also,  $\bar{A}^m \subseteq A^l$  and  $a^{l+1} \in A^l \setminus \bar{A}^m$  imply that  $\#\bar{A}^m < \#A^l$ . That is,  $l > m$ . So,  $b^m \neq j = b^l$ . This implies  $b^m \notin (\bar{A}^{m-1} \cup \{j\}) = \bar{A}^m$ .

Now,  $a^m \in A^{m-1} = \bar{A}^{m-1}$ . So,  $a^m \in \bar{A}^m$ . But  $a^m \in \bar{A}^m$  and  $b^m \notin \bar{A}^m$  together imply that  $\bar{c}_{\bar{a}^{m+1}\bar{b}^{m+1}} \leq \bar{c}_{a^m b^m} \leq c_{a^m b^m}$ .

So, using equation 9,  $\bar{c}_{\bar{a}^{m+1}\bar{b}^{m+1}} \leq c_{a^m b^m} \leq c_{a^l b^l} \leq c_{a^{l+1}b^{l+1}}$ .

Hence,  $\psi^*$  satisfies cost monotonicity.<sup>16</sup>

We now show that for all  $C \in \mathcal{C}$ ,  $\psi^*(C)$  is an element in the core of the cost game corresponding to  $C$ .

Again, we present the proof for any  $C \in \mathcal{C}^2$  in order to avoid notational complications.<sup>17</sup> We want to show that for all  $S \subseteq N$ ,  $\sum_{i \in S} \psi_i^*(C) \leq c(S)$ .

Without loss of generality, assume that for all  $i \in N$ ,  $b^i = i$  and denote  $c_{a^k b^k} = c^k$ .

**Claim 1:** If  $S = \{1, 2, \dots, K\}$  where  $K \leq \#N$ , then  $\sum_{i \in S} \psi_i^*(C) \leq c(S)$ .

**Proof of Claim:** Clearly,  $g = \cup_{k=1}^K \{a^k k\}$  is a connected graph over  $S \cup \{0\}$ .

Also,  $g$  is in fact the m.c.s.t. over  $S$ .

<sup>16</sup>Suppose  $C \notin \mathcal{C}^2$ . What we have shown above is that the outcome of the algorithm for *each* tie-breaking rule satisfies cost monotonicity. Hence, the average must also satisfy cost monotonicity.

<sup>17</sup>Suppose instead that  $C \notin \mathcal{C}^2$ . Then, our subsequent proof shows that the outcome of the algorithm is in the core for *each*  $\sigma \in \Sigma$ . Since the core is a convex set, the average (that is,  $\psi^*$ ) must be in the core if each  $\psi_\sigma^*$  is in the core.

So,  $c(S) = \sum_{k=1}^K c^k$ . Also,  $\sum_{i \in S} \psi_i^*(C) = \sum_{k=1}^{K+1} c^k - \max_{1 \leq k \leq K+1} c^k \leq \sum_{k=1}^K c^k = c(S)$ . ■

Hence, a blocking coalition cannot consist of an initial set of integers, given our assumption that  $b^k = k$  for all  $k \in N$ .

Now, let  $S$  be a largest blocking coalition. That is,

- (i)  $\sum_{i \in S} \psi_i^*(C) > c(S)$ .
- (ii) If  $S \subset T$ , then  $\sum_{i \in T} \psi_i^*(C) \leq c(T)$ .

There are two possible cases.

**Case 1:**  $1 \notin S$ .

Let  $K = \min_{j \in S} j$ . Consider  $T = \{1, \dots, K-1\}$ . We will show that  $S \cup T$  is also a blocking coalition, contradicting the description of  $S$ .

Now,

$$\sum_{i \in T \cup S} \psi_i^*(C) = \sum_{i \in S} \psi_i^*(C) + \sum_{i \in T} \psi_i^*(C) > c(S) + \sum_{k=1}^K c^k - \max_{1 \leq k \leq K} c^k \geq c(S) + \sum_{k=1}^K c^k - c_{0s},$$
where  $(0s) \in g_S$ , the m.c.s.t. of  $S$ . Note that the last inequality follows from the fact that  $c^k \leq c_{0s}$  for all  $k \in \{1, \dots, K\}$ .

Since  $g = (\cup_{k=1}^K a^k b^k) \cup (g_S \setminus \{(0s)\})$  is a connected graph over  $(T \cup S \cup \{0\})$ ,  $c(S) + \sum_{k=1}^K c^k - c_{0s} \geq c(S \cup T)$ . Hence,  $\sum_{i \in S \cup T} \psi_i^*(C) > c(S \cup T)$ , establishing the contradiction that  $S \cup T$  is a blocking coalition.

**Case 2:**  $1 \in S$ .

From the claim,  $S$  is not an initial segment of the integers. So, we can partition  $S$  into  $\{S_1, \dots, S_K\}$ , where each  $S_k$  consists of consecutive integers, and  $i \in S_k, j \in S_{k+1}$  implies that  $i+1 < j$ . Assume  $m = \max_{j \in S_1} j$  and  $n = \min_{j \in S_2} j$ . Note that  $n > m+1$ . Define  $T = \{m+1, \dots, n-1\}$ . We will show that  $S \cup T$  is a blocking coalition, contradicting the assumption that  $S$  is a largest blocking coalition.

$$\begin{aligned} \sum_{i \in S \cup T} \psi_i^*(C) &= \sum_{i \in S} \psi_i^*(C) + \sum_{i \in T} \psi_i^*(C) \\ &> c(S) + \sum_{i \in S_1 \cup T} \psi_i^*(C) - \sum_{i \in S_1} \psi_i^*(C) \end{aligned}$$

$$\begin{aligned}
&= c(S) + \left( \sum_{i=1}^n c^i - \max_{1 \leq i \leq n} c^i \right) - \left( \sum_{i=1}^{m+1} c^i - \max_{1 \leq i \leq m+1} c^i \right) \\
&= c(S) + \left( \sum_{i=m+2}^n c^i + \max_{1 \leq i \leq m+1} c^i - \max_{1 \leq i \leq n} c^i \right) \\
&\geq c(S) + \sum_{i=m+1}^n c^i - \max_{1 \leq i \leq n} c^i
\end{aligned}$$

Since  $g_S$  is a connected graph over  $S^+$ , there is  $s_2 \in S \setminus S_1$  and  $s_1 \in S_1^+$  such that  $(s_1 s_2) \in g_S$ . Moreover,  $c_{s_1 s_2} \geq \max_{1 \leq i \leq n} c^i$ . Noting that  $\cup_{k=m+1}^n \{(a^k b^k)\} \cup [g_S \setminus \{(s_1 s_2)\}]$  is a connected graph over  $S \cup T \cup \{0\}$ , it follows that

$$\begin{aligned}
\sum_{i \in S \cup T} \psi_i^*(C) &> c(S) + \sum_{i=m+1}^n c^i - c_{s_1 s_2} \\
&\geq c(S \cup T).
\end{aligned}$$

So,  $S \cup T$  is a blocking coalition, establishing the desired contradiction.

This concludes the proof of the theorem. ■

The following lemmas will be used in the proofs of Theorems 2 and 3.

**Lemma 1 :** Let  $C \in \mathcal{C}_N^1$ , and  $i \in N$ . If  $c_{ik} = \min_{l \in N^+ \setminus \{i\}} c_{il}$ , then  $(ik) \in g_N(C)$ .

**Proof :** Suppose  $(ik) \notin g_N(C)$ . As  $g_N(C)$  is a connected graph over  $N^+$ ,  $\exists j \in N^+ \setminus \{i, k\}$  such that  $(ij) \in g_N(C)$  and  $j$  is on the path between  $i$  and  $k$ . But,  $\{g_N \cup (ik)\} \setminus \{(ij)\}$  is still a connected graph which costs no more than  $g_N(C)$ , as  $c_{ik} \leq c_{ij}$ . This is not possible as  $g_N(C)$  is the only m.c.s.t. of  $C$ . ■

**Lemma 2 :** Let  $C \in \mathcal{C}_N^2$ , and  $(01) \in g_N(C)$ . Let  $\psi_1(C) = \min_{k \in N^+ \setminus \{1\}} c_{1k}$ . Then,  $C_{\psi_1(C)}^{tr} \in \mathcal{C}_{N \setminus \{1\}}^2$ .

**Proof :** We will denote  $C_{\psi_1(C)}^{tr}$  by  $\bar{C}$  for the rest of this proof.

Let  $\psi_1(C) = \min_{k \in N^+ \setminus \{1\}} c_{1k} = c_{1k^*}$  (say).

Suppose there exists  $(ij) \in g_N(C)$  such that  $i, j \neq 1$ . Without loss of generality assume  $i$  precedes  $j$  in  $g_N(C)$ . Since  $(01), (ij) \in g_N(C)$ ,  $(1j) \notin g_N(C)$ . Then,  $c_{1j} > c_{ij}$ . As  $\psi_1(C) \leq c_{i1}$ ,  $c_{i1} + c_{1j} - \psi_1(C) \geq c_{1j} > c_{ij}$ . Hence  $\bar{c}_{ij} = c_{ij} \forall (ij) \in g_N(C)$ , such that  $i, j \neq 1$ .

Now, suppose there is  $j \in N^+$  such that  $j \neq k^*$  and  $(1j) \in g_N(C)$ . Since  $(1j), (1k^*) \in g_N(C)$ ,  $(jk^*) \notin g_N(C)$ . Hence,  $c_{1j} < c_{k^*j}$ . Thus,

$$\bar{c}_{k^*j} = \min\{(c_{1j} + c_{1k^*} - \psi_1(C)), c_{k^*j}\} = \min(c_{1j}, c_{k^*j}) = c_{1j}.$$

Next, let  $\bar{g}_{N \setminus \{1\}}$ , be a connected graph over  $N^+ \setminus \{1\}$ , defined as follows.

$$\bar{g}_{N \setminus \{1\}} = \{(ij) \mid \text{either } (ij) \in g_N(C) \text{ s.t. } i, j \neq 1 \text{ or } (ij) = (k^*l) \text{ where } (1l) \in g_N(C)\}.$$

Note that no two edges have equal cost in  $\bar{g}_{N \setminus \{1\}}$ .

Also,

$$\sum_{(ij) \in \bar{g}_{N \setminus \{1\}}} \bar{c}_{ij} = \sum_{(ij) \in g_N(C)} c_{ij} - c_{1k^*}. \quad (10)$$

We prove that  $\bar{C}$  belongs to  $\mathcal{C}_{N \setminus \{1\}}^2$  by showing that  $\bar{g}_{N \setminus \{1\}}$  is the only m.c.s.t. of  $\bar{C}$ .

Suppose this is not true, so that  $g_{N \setminus \{1\}}^*$  is an m.c.s.t. corresponding to  $\bar{C}$ . Then, using 10,

$$\sum_{(ij) \in g_{N \setminus \{1\}}^*} \bar{c}_{ij} \leq \sum_{(ij) \in g_N(C)} c_{ij} - c_{1k^*} \quad (11)$$

Let  $g_{N \setminus \{1\}}^* = g^1 \cup g^2$ , where

$$\begin{aligned} g^1 &= \{(ij) \mid (ij) \in g_{N \setminus \{1\}}^*, c_{ij} = \bar{c}_{ij}\} \\ g^2 &= g_{N \setminus \{1\}}^* \setminus g^1 \end{aligned}$$

If  $(ij) \in g^2$ , then

$$\begin{aligned} \bar{c}_{ij} &= \min(c_{ij}, c_{1i} + c_{1j} - \psi_1(C)) \\ &= c_{1i} + c_{1j} - \psi_1(C) \\ &\geq \max(c_{1i}, c_{1j}) \end{aligned}$$

where the last inequality follows from the assumption that  $\psi_1(C) = \min_{k \in N^+ \setminus \{1\}} c_{1k}$ .

So,

$$\begin{aligned} \bar{c}_{ij} &= c_{ij} \text{ if } (ij) \in g^1 \\ &\geq \max(c_{1i}, c_{1j}) \text{ if } (ij) \in g^2. \end{aligned} \quad (12)$$

Now, extend  $g_{N \setminus \{1\}}^*$  to a connected graph  $g'_N$  over  $N^+$  as follows. Letting  $g = \{(1i) \mid (ij) \in g^2, j \in U(i, k^*, g_{N \setminus \{1\}}^*)\}$ , define

$$g'_N = g^1 \cup (1k^*) \cup g.$$

**Claim:**  $g'_N$  is a connected graph over  $N^+$ .

**Proof of Claim:** It is sufficient to show that every  $i \in N^+ \setminus \{1\}$  is connected to 1 in  $g'_N$ . Clearly, this is true for  $i = k^*$ . Take any  $i \in N^+ \setminus \{1, k^*\}$ . Let  $U(i, k^*, g_{N \setminus \{1\}}^*) = \{m_0, m_1, \dots, m_{p+1}\}$ <sup>18</sup> where  $m_0 = i$  and  $m_{p+1} = k^*$ . If all these edges  $(m_t m_{t+1}) \forall t \leq p$  are in  $g^1$ , then they are also in  $g'_N$ , and there is nothing to prove.

So, suppose there is  $(m_t m_{t+1}) \in g^2$  while all edges in  $\{(m_0 m_1), \dots, (m_{t-1} m_t)\}$  belong to  $g^1$ . In this case,  $(m_t 1)$  as well as all edges in  $\{(m_0 m_1), \dots, (m_{t-1} m_t)\}$  belong to  $g'_N$ . Hence,  $i$  is connected to 1.  $\blacksquare$

To complete the proof of the lemma, note that

$$\sum_{(ij) \in g'_N} c_{ij} = \sum_{(ij) \in g^1} c_{ij} + c_{1k^*} + \sum_{(ij) \in g^2} c_{ij}.$$

Using (12),

$$\sum_{(ij) \in g'_N} c_{ij} \leq \sum_{(ij) \in g^1} \bar{c}_{ij} + c_{1k^*} + \sum_{(ij) \in g^2} \bar{c}_{ij} = \sum_{(ij) \in g_{N \setminus \{1\}}^*} \bar{c}_{ij} + c_{1k^*}$$

Finally, using (11),

$$\sum_{(ij) \in g'_N} c_{ij} \leq \sum_{(ij) \in g_N(C)} c_{ij}.$$

But, this contradicts the assumption that  $g_N(C)$  is the unique m.c.s.t. for  $C$ .  $\blacksquare$

**Lemma 3 :** Let  $C \in \mathcal{C}_N^1$ ,  $(10) \in g_N(C)$ . Suppose  $\psi_1(C) = c_{01}$ . Then  $C_{\psi_1(C)}^{sr} \in \mathcal{C}_{N \setminus \{1\}}^1$ .

**Proof :** Throughout the proof of this lemma, we denote  $C_{\psi_1(C)}^{sr}$  by  $\bar{C}$ .

We know  $\psi_1(C) = c_{01}$ . Suppose  $(ij) \in g_N(C)$  such that  $\{i, j\} \cap \{0, 1\} = \emptyset$ . Then  $\bar{c}_{ij} = c_{ij}$ .

On the other hand if  $(i0) \in g_N(C)$ , and  $i \neq 1$ , then  $\bar{c}_{0i} = \min\{(c_{i1} + c_{10} - \psi_1(C)), c_{0i}\} = \min\{c_{i1}, c_{0i}\} = c_{i0}$ . Note that the last equality follows from the fact that  $(i0) \in g_N(C)$  but  $(i1) \notin g_N(C)$  implies that  $c_{i1} > c_{i0}$ .

<sup>18</sup>This path exists because  $g_{N \setminus \{1\}}^*$  is a connected graph.

If  $(i1) \in g_N(C)$ , then  $\bar{c}_{i0} = \min\{(c_{i1} + c_{10} - \psi_1(C)), c_{i0}\} = \min(c_{i1}, c_{i0}) = c_{i1}$ , as  $(i1) \in g_N(C)$  but  $(i0) \notin g_N(C)$ .

Now we construct  $\bar{g}_{N \setminus \{1\}}$ , a connected graph over  $N^+ \setminus \{1\}$  as follows.

$$\bar{g}_{N \setminus \{1\}} = \{(ij) \mid \text{either } (ij) \in g_N(C) \text{ s.t. } i, j \neq 1 \text{ or } (ij) = (l0) \text{ where } (l1) \in g_N(C)\}$$

Then,  $\bar{g}_{N \setminus \{1\}}$  must be the only m.c.s.t. of  $\bar{C}$ . For if there is another  $g_{N \setminus \{1\}}^*$  which is also an m.c.s.t. of  $\bar{C}$ , then one can show that  $g_N(C)$  cannot be the only m.c.s.t. corresponding to  $C$ .<sup>19</sup> ■

**Lemma 4:** Suppose  $\psi$  satisfies TR, EPM and EF. Let  $C \in \mathcal{C}_N^2$ . If  $(i0) \in g_N(C)$ , then  $\psi_i(C) \geq \min_{k \in N^+ \setminus \{i\}} c_{ik}$ .

**Proof:** Consider any  $C \in \mathcal{C}_N^2$ ,  $(i0) \in g_N(C)$ , and  $\psi$  satisfying TR, EPM, EF. Let  $\psi(C) = x$ , and  $c_{im} = \min_{k \in N^+ \setminus \{i\}} c_{ik}$ . We want to show that  $x_i \geq c_{im}$ .

Choose  $j \notin N^+$ , and define  $\bar{N} = N \cup \{j\}$ . Let  $\bar{C} \in \mathcal{C}_{\bar{N}}^2$  be such that

(i)  $\bar{C}$  coincides with  $C$  on  $N^+$ .

(ii) For all  $k \in N^+ \setminus \{i\}$ ,  $\bar{c}_{jk} > \bar{c}_{ij} > \sum_{(pq) \in g_N(C)} c_{pq}$ .<sup>20</sup>

Hence,  $g_{\bar{N}}(\bar{C}) = g_N(C) \cup \{(ij)\}$ .

Notice that  $j$  is an extreme point of  $\bar{C}$ . Denoting  $\psi(\bar{C}) = \bar{x}$ , EPM implies that

$$x_k \geq \bar{x}_k \forall k \in N \tag{13}$$

We prove the lemma by showing that  $\bar{x}_i \geq \bar{c}_{im} = c_{im}$ .

Let  $\bar{C}_{\bar{x}_i}^{tr} = C'$ , and  $N' = \bar{N} \setminus \{i\}$ ,  $\psi(C') = x'$ . Assume  $\bar{x}_i < \bar{c}_{im}$ .

**Case 1:**  $C' \in \mathcal{C}_{N'}^2$ .

---

<sup>19</sup>The proof of this assertion is analogous to that of the corresponding assertion in Lemma 2, and is hence omitted.

<sup>20</sup>Note that the exact lower bound on  $\bar{c}_{ij}$  will play no role in the subsequent proof. All that we require is that  $\bar{c}_{ij}$  is “high” so that the reduced matrix  $C'$ , to be defined below, has appropriate properties.

Suppose there is some  $k \in N'$  such that  $(ik) \notin g_{\bar{N}}(\bar{C})$ . Let  $l$  be the predecessor of  $k$  in  $g_{\bar{N}}(\bar{C})$ . Since  $(kl) \in g_{\bar{N}}(\bar{C})$  and  $(ik) \notin g_{\bar{N}}(\bar{C})$ ,  $\bar{c}_{kl} < \bar{c}_{ki}$ . Also,  $\bar{c}_{il} \geq \bar{c}_{im} > \bar{x}_i$ . Hence,

$$c'_{kl} = \min(\bar{c}_{kl}, \bar{c}_{ki} + \bar{c}_{li} - \bar{x}_i) = \bar{c}_{kl} \quad (14)$$

Now, consider  $k \in (N' \cup \{0\}) \setminus \{m, j\}$  such that  $(ik) \in g_{\bar{N}}(\bar{C})$ . Note that  $(km) \notin g_{\bar{N}}(\bar{C})$  since  $(im) \in g_{\bar{N}}(\bar{C})$  from lemma 1. Hence,  $\bar{c}_{km} > \bar{c}_{ik}$  since  $(ik) \in g_{\bar{N}}(\bar{C})$  and  $(km) \notin g_{\bar{N}}(\bar{C})$ . So,

$$c'_{km} = \min(\bar{c}_{km}, \bar{c}_{ik} + \bar{c}_{im} - \bar{x}_i) > \bar{c}_{ik} \quad (15)$$

Take any  $(kl) \notin g_{\bar{N}}(\bar{C})$ . Let  $U(k, l, g_{\bar{N}}(\bar{C})) = \{s_1, \dots, s_J\}$ . Then either  $s_j, s_{j+1} \in U(k, i, g_{\bar{N}}(\bar{C}))$  or  $s_j, s_{j+1} \in U(l, i, g_{\bar{N}}(\bar{C}))$ . Without loss of generality, assume  $s_j, s_{j+1} \in U(k, i, g_{\bar{N}}(\bar{C}))$ . Then,  $\bar{c}_{s_j s_{j+1}} < \bar{c}_{kl}$  and  $\bar{c}_{s_j s_{j+1}} \leq \bar{c}_{ki}$ . As  $\bar{x}_i < \bar{c}_{il}$

$$c'_{kl} = \min(\bar{c}_{ki} + \bar{c}_{il} - \bar{x}_i, \bar{c}_{kl}) > \bar{c}_{s_j s_{j+1}} \quad (16)$$

Next, note that since  $\bar{c}_{jm}$  can be chosen arbitrarily high,

$$c'_{jm} = \min(\bar{c}_{jm}, \bar{c}_{ij} + \bar{c}_{im} - \bar{x}_i) = \bar{c}_{ij} + \bar{c}_{im} - \bar{x}_i \quad (17)$$

Since for all  $t \in (N' \cup \{0\}) \setminus \{m, j\}$ ,  $c'_{jt} = \bar{c}_{ij} + \bar{c}_{it} - \bar{x}_i > c'_{jm}$ ,  $(jm) \in g_{N'}(C')$ .

From TR, we have  $x'_k = \bar{x}_k$  for all  $k \in N'$ . Using EF, and equations 14, 15, 16, 17,

$$\sum_{k \in \bar{N} \setminus \{i\}} \bar{x}_k = \sum_{k \in N'} x'_k = c(g_{N'}(C')) > c(g_{\bar{N}}(\bar{C})) - \bar{x}_i \quad (18)$$

But, this violates EF since  $\sum_{k \in \bar{N}} \bar{x}_k > c(g_{\bar{N}}(\bar{C}))$ .

**Case 2:**  $C' \notin \mathcal{C}_{N'}^2$ .

Let  $S_1 = \{(q_1 q_2) \in g_{\bar{N}}(\bar{C}) \mid \{q_1, q_2\} \cap \{i\} = \emptyset\}$ . From equation 14, we have

$$c'_{q_1 q_2} = \bar{c}_{q_1 q_2} \quad (19)$$

Let  $S_2 = \{(kl) \mid k, l \in N'^+, (kl) \notin S_1\}$ . If  $(kl) \in S_2$  then from equation 16 we get

$$c'_{kl} > \bar{c}_{s_j s_{j+1}} \forall s_j, s_{j+1} \in U(k, l, g_{\bar{N}}(\bar{C})) \quad (20)$$

For each  $(kl) \in S_2$ , choose  $p(k, l) \notin \bar{N}^+$  such that  $p(k, l) \neq p(k', l')$  when  $(k, l) \neq (k', l')$ . Let  $\hat{N} = \bar{N} \cup \{p(k, l) | (k, l) \in S_2\}$ . Now define a new matrix  $\hat{C}$  over  $\hat{N}^+$  as follows,

- (i)  $\hat{C}$  coincides with  $\bar{C}$  on  $\bar{N}^+$ .
- (ii)  $\hat{c}_{p(k, l)k} = \min_{t \in \bar{N}^+} \hat{c}_{p(k, l)t}$  and  $\hat{c}_{p(k, l)k} \neq \bar{c}_{q_1 q_2}$  for all  $(q_1 q_2) \in S_1$
- (iii)  $c'_{kl} > \hat{c}_{p(k, l)l} > \bar{c}_{s_j s_{j+1}} \forall s_j, s_{j+1} \in U(k, l, g_{\bar{N}}(\bar{C}))$  and  $\hat{c}_{p(k, l)l} \neq \bar{c}_{q_1 q_2}$  for all  $(q_1 q_2) \in S_1$ .<sup>21</sup>
- (iv) Costs of all the remaining edges are chosen ‘very’ high, for example strictly greater than  $\sum_{(t_1 t_2)} \bar{c}_{t_1 t_2}$ .
- (v) Costs of all the new edges, that is those which were not present in  $\bar{C}$ , are distinct.

Then, from specifications (ii) and (iv) we have  $g_{\hat{N}}(\hat{C}) = g_{\bar{N}}(\bar{C}) \cup \{(p(k, l)k) | (kl) \in S_2\}$ , so that all  $p(k, l)$  are extreme points of  $\hat{C}$ . Let  $\psi(\hat{C}) = \hat{x}$ . From EPM,

$$\bar{x}_i \geq \hat{x}_i \tag{21}$$

Now, consider the reduced matrix  $\tilde{C} \equiv \hat{C}_{\hat{x}_i}^{tr}$ . We want to show that the edges which will be connected in the m.c.s.t. of  $\tilde{C}$  will be contained in  $S_1 \cup \{(p(k, l)k), (p(k, l), l) | (k, l) \in S_2\}$ . Since specifications (i), (ii), (iii) and (v) imply that all these costs are distinct, we will get  $\tilde{C} \in \mathcal{C}_{\hat{N} \setminus \{i\}}^2$ .

Consider any  $(q_1 q_2) \in S_1 \cup S_2$ . Since  $\bar{x}_i \geq \hat{x}_i$ ,  $\bar{c}_{q_1 i} + \bar{c}_{q_2 i} - \bar{x}_i \geq \bar{c}_{q_1 i} + \bar{c}_{q_2 i} - \hat{x}_i$ . Thus,  $\tilde{c}_{q_1 q_2} \geq c'_{q_1 q_2}$ , with equality holding if  $\bar{c}_{q_1 q_2} \leq \bar{c}_{q_1 i} + \bar{c}_{q_2 i} - \bar{x}_i$ . Hence

$$\tilde{c}_{q_1 q_2} = \bar{c}_{q_1 q_2} = c'_{q_1 q_2} \text{ if } (q_1 q_2) \in S_1, \tilde{c}_{q_1 q_2} \geq c'_{q_1 q_2} \text{ otherwise} \tag{22}$$

Since  $\hat{x}_i < \hat{c}_{ik}$ , and  $\hat{c}_{p(k, l)k} < \hat{c}_{p(k, l)i}$ ,

$$\tilde{c}_{p(k, l)k} = \min(\hat{c}_{p(k, l)k}, \hat{c}_{p(k, l)i} + \hat{c}_{ik} - \hat{x}_i) = \hat{c}_{p(k, l)k}.$$

---

<sup>21</sup>Note that this specification of costs is valid because (20) is true.

Also, using specification (iv),

$$\tilde{c}_{p(k,l)l} = \min(\hat{c}_{p(k,l)l}, \hat{c}_{p(k,l)i} + \hat{c}_{il} - \hat{x}_i) = \hat{c}_{p(k,l)l}$$

It is easy to check that all other edges in  $\tilde{C}$  have costs strictly greater than  $\sum_{(t_1 t_2)} \bar{c}_{t_1 t_2}$ .

So, from specification (ii),  $\tilde{c}_{p(k,l)k} = \min_{t \in \hat{N}^+ \setminus \{p(k,l), i\}} \tilde{c}_{p(k,l)t}$ . Hence, from lemma 1,  $(p(k,l)k) \in g_{\hat{N} \setminus i}(\tilde{C})$ .

Finally, specification (iii) implies  $\tilde{c}_{kl} \geq c'_{kl} > \hat{c}_{p(k,l)l} = \tilde{c}_{p(k,l)l}$  for all  $(kl) \in S_2$ .

Thus for all  $(kl) \in S_2$ ,  $(p(k,l)k)$  is connected in a m.c.s.t of  $\tilde{C}$  and  $\tilde{c}_{kl} > \tilde{c}_{p(k,l)l}$ . So,  $(kl)$  cannot be an edge in the m.c.s.t. corresponding to  $\tilde{C}$ . This establishes that the m.c.s.t. can only contain edges from  $S_1 \cup \{(p(k,l)k), (p(k,l)l) | (kl) \in S_2\}$ .

Since  $\tilde{C} \in \mathcal{C}_{\hat{N} \setminus \{i\}}^2$ , we apply the conclusion of Case 1 of the lemma to conclude that  $\hat{x}_i \geq \hat{c}_{im} = \bar{c}_{im}$ . Equation 21 now establishes that  $\bar{x}_i \geq \bar{c}_{im}$ .  $\blacksquare$

**Lemma 5:** Suppose  $\psi$  satisfies SR, EPM and EF. Let  $C \in \mathcal{C}_N^2$ . If  $(i0) \in g_N(C)$ , then  $\psi_i(C) \geq \min_{k \in N^+ \setminus \{i\}} c_{ik}$ .

The proof of Lemma 5 is almost identical to that of Lemma 4, and hence is omitted.

**Proof of Theorem 2 :** First, we prove that  $\psi^*$  satisfies all the three axioms.

Let  $C \in \mathcal{C}^2$ .

Efficiency follows trivially from the algorithm which defines the allocation.

Next, we show that  $\psi^*$  satisfies TR.

Let  $(10) = \operatorname{argmin}_{k \in N} c_{k0}$ . Hence, the algorithm yields  $b^1 = 1$ , and  $\psi_1^*(C) = \min(c_{10}, c_{a^2 b^2})$ . There are two possible choice of  $a^2$ .

*Case 1:*  $a^2 = 1$ . Then, we get  $c_{1b^2} = \min_{k \in N \setminus \{1\}} c_{1k}$ . Therefore  $\psi_1^*(C) = \min(c_{10}, c_{1b^2}) = \min_{k \in N^+ \setminus \{1\}} c_{1k}$ .

*Case 2:*  $a^2 = 0$ . Then,  $c_{b^2 0} \leq c_{1k} \forall k \in N \setminus \{1\}$ . Since  $c_{10} \leq c_{b^2 0}$ , we conclude  $\psi_1^*(C) = \min(c_{10}, c_{b^2 0}) = c_{10} = \min_{k \in N^+ \setminus \{1\}} c_{1k}$ .

So, in either case, 1 pays its minimum cost.

Let  $\psi_1^*(C) = x_1 = \min_{k \in N^+ \setminus \{1\}} c_{1k} = c_{1k^*}$ . Denoting  $C_{x_1}^{tr}$  by  $\bar{C}$ , we know from Lemma 2, that  $\bar{C} \in \mathcal{C}^2$ . Hence, the algorithm is well defined on  $\bar{C}$ .

Let  $\bar{a}^k, \bar{b}^k, \bar{t}^k$ , etc denote the relevant variables of the algorithm corresponding to  $\bar{C}$ .

**Claim:**  $\forall i \in N \setminus \{1\}, \psi_i^*(C) = \psi_i^*(\bar{C})$ . That is,  $\psi^*$  satisfies Tree Consistency.

**Proof of Claim:** From the proof of lemma 2,

- (i)  $\bar{c}_{ij} = c_{ij} \forall (ij) \in g_N(C)$  s.t.  $i, j \neq 1$ .
- (ii)  $\bar{c}_{k^*j} = c_{1j}$  for  $j \in N^+ \setminus \{k^*\}$  s.t.  $(1j) \in g_N(C)$ .

Also,

$$g_{N \setminus \{1\}}(\bar{C}) = \{(ij) | (ij) \in g_N(C) \text{ if } i, j \neq 1 \text{ and } (ij) = (k^*l) \text{ if } (1l) \in g_N(C)\}.$$

Let  $b^2 = i$ . Either  $k^* = 0$  or  $k^* = i$ . In either case,  $\bar{c}_{0i} < \bar{c}_{0j}$  for  $j \notin \{0, 1, i\}$ . Hence,  $\bar{b}^1 = i$ .

$$\text{Now, } t^2 = \max(c_{a^1 b^1}, c_{a^2 b^2}) = \max(c_{10}, c_{a^2 i}) = \bar{c}_{0i} = \bar{t}^1.$$

Also,  $a^3 \in \{0, 1, i\}$ , while  $b^3 \in \{0, 1, i\}_c$ . If  $a^3 \in \{0, i\}$ , then  $\bar{a}^2 = a^3$ . If  $a^3 = 1$ , then  $\bar{a}^2 = k^*$ . In all cases,  $b^3 = \bar{b}^2$ , and  $c_{a^3 b^3} = \bar{c}_{\bar{a}^2 \bar{b}^2}$ . So,

$$\psi_i^*(C) = \min(t^2, c_{a^3 b^3}) = \min(\bar{t}^1, \bar{c}_{\bar{a}^2 \bar{b}^2}) = \psi_i^*(\bar{C}). \quad (23)$$

The claim is established for  $\{b^3, \dots, b^n\}$  by using the structure of  $g_{N \setminus \{1\}}(\bar{C})$ , the definition of  $\bar{C}$  given above, and the following induction hypothesis. The details are left to the reader.

For all  $i = 2, \dots, k-1$ ,

- (i)  $\bar{b}^{i-1} = b^i$ .
- (ii)  $\bar{t}^{i-1} = t^i$ .
- (iii)  $\bar{a}^{i-1} = a^i$  if  $a^i \neq 1$ , and  $\bar{a}^{i-1} = k^*$  if  $a^i = 1$ .

We now have to show that  $\psi^*$  satisfies EPM.

Let  $i \in N$  be an extreme point of  $g_N(C)$ , and  $\hat{C}$  be the restriction of  $C$  over  $N \setminus \{i\}$ . Of course,  $\hat{C} \in \mathcal{C}^2$ .

In order to differentiate between the algorithms on  $C$  and  $\widehat{C}$ , we denote the outcomes corresponding to the latter by  $\widehat{a}^k, \widehat{b}^k, \widehat{t}^k$ , etc.

Suppose  $b^k = i$ . Clearly, the algorithm will produce the same outcomes till step  $(k-1)$ , and so  $\psi_j^*(C) = \psi_j^*(\widehat{C})$  for all  $j \in \{b^1, \dots, b^{k-2}\}$ , and  $t^{k-1} = \widehat{t}^{k-1}$ .

Now, we calculate  $\psi_j^*(C)$  where  $j = b^{k-1}$ . As  $i$  is an extreme point of  $g_N$ , and  $(a^k i) \in g_N$ ,  $a^{k+1} \neq i$ . Also,  $A^k = A^{k-1} \cup \{i\}$ . Hence,  $a^{k+1} \in A^{k-1}$ . This implies  $c_{a^k i} \leq c_{a^{k+1} b^{k+1}}$ . But  $i \notin \widehat{A}_c^{k-1}$ . Hence  $(\widehat{a}^k \widehat{b}^k) = (a^{k+1} b^{k+1})$ . Thus,

$$\psi_j^*(C) = \min(t^{k-1}, c_{a^k i}) \leq \min(\widehat{t}^{k-1}, c_{a^{k+1} b^{k+1}}) = \psi_j^*(\widehat{C}) \quad (24)$$

Also,  $\widehat{t}^k = \max(\widehat{t}^{k-1}, c_{a^{k+1} b^{k+1}}) \geq \max(t^{k-1}, c_{a^k i}, c_{a^{k+1} b^{k+1}}) = t^{k+1}$ . The algorithm on  $C$  determines the cost allocation for  $i$  in step  $(k+1)$ . Since  $i$  is an extreme point of  $g_N$ ,  $i \neq a^s$  for any  $s$ . Hence, the choice of  $a^j$  and  $b^j$  must be the same in  $C$  and  $\widehat{C}$  for  $j \geq k+1$ . So, for all  $j \in \{k+1, \dots, \#N\}$ ,  $a^j = \widehat{a}^{j-1}, b^j = \widehat{b}^{j-1}, t^j \leq \widehat{t}^{j-1}$ . Hence,

$$\psi_{b^j}^*(C) = \min(t^j, c_{a^{j+1} b^{j+1}}) \leq \min(\widehat{t}^{j-1}, c_{\widehat{a}^j \widehat{b}^j}) = \psi_{\widehat{b}^{j-1}}^*(\widehat{C}) \quad (25)$$

Hence, we can conclude that  $\psi^*$  satisfies Extreme Point Monotonicity.

Next, we will prove that only *one* rule over  $\mathcal{C}^2$  satisfies all three axioms. Let  $\psi$  be a rule satisfying all the three axioms. We will show by induction on the cardinality of the set of nodes that  $\psi$  is unique.

Let us start by showing that the result is true for  $|N| = 2$ . There are several cases.

**Case 1:**  $c_{12} > c_{10}, c_{20}$ . From Lemma 4,  $\psi_1(C) \geq c_{10}, \psi_2(C) \geq c_{20}$ . By EF,  $\psi_1(C) + \psi_2(C) = c_{10} + c_{20}$ . Thus  $\psi_1(C) = c_{10}$ , and  $\psi_2(C) = c_{20}$ . So, the allocation is unique.

**Case 2:**  $c_{20} > c_{12} > c_{10}$ . Introduce a third agent 3 and costs  $c_{20} < \bar{c}_{13} < \min(\bar{c}_{32}, \bar{c}_{30})$ . Let the restriction of  $\bar{C}$  on  $\{1, 2\}^+$  coincide with  $C$ . Hence,  $g_{\{1,2,3\}} = \{(01), (12), (13)\}$ . Let  $\psi(\bar{C}) = \bar{x}$ . From Lemma 4,  $\bar{x}_1 \geq \bar{c}_{10} = c_{10}$ .

Denote the reduced matrix  $\bar{C}_{\bar{x}_1}^{tr}$  as  $\widehat{C}$ . Now,  $\widehat{c}_{02} = \min(\bar{c}_{01} + \bar{c}_{12} - \bar{x}_1, \bar{c}_{02}) = \bar{c}_{01} + \bar{c}_{12} - \bar{x}_1$ . Similarly,  $\widehat{c}_{23} = \min(\bar{c}_{13} + \bar{c}_{12} - \bar{x}_1, \bar{c}_{23})$ . Noting that  $\bar{x}_1 \geq \bar{c}_{10}, \bar{c}_{23} > \bar{c}_{12}$

and  $\bar{c}_{13} > \bar{c}_{10}$ , we conclude that

$$\hat{c}_{02} < \hat{c}_{23}.$$

Analogously,  $\hat{c}_{03} = \bar{c}_{01} + \bar{c}_{13} - \bar{x}_1 < \hat{c}_{23}$ .

Hence,  $g_{\{2,3\}}(\hat{C}) = \{(02), (03)\}$ . So,  $\hat{C} \in \mathcal{C}^2$ . Using TR,

$$\psi_2(\hat{C}) = \psi_2(\bar{C}), \psi_3(\hat{C}) = \psi_3(\bar{C}) \quad (26)$$

From Case 1 above,

$$\psi_2(\hat{C}) = \bar{c}_{01} + \bar{c}_{12} - \bar{x}_1, \psi_3(\hat{C}) = \bar{c}_{01} + \bar{c}_{13} - \bar{x}_1 \quad (27)$$

From (26) and (27),

$$\begin{aligned} \psi_2(\bar{C}) + \psi_3(\bar{C}) &= \bar{c}_{01} + \bar{c}_{12} - \bar{x}_1 + \bar{c}_{01} + \bar{c}_{13} - \bar{x}_1 \\ \text{or } \bar{x}_1 + \psi_2(\bar{C}) + \psi_3(\bar{C}) &= \bar{c}_{01} + \bar{c}_{12} + \bar{c}_{13} + (\bar{c}_{01} - \bar{x}_1) \end{aligned}$$

But, from EF,  $\bar{x}_1 + \psi_2(\bar{C}) + \psi_3(\bar{C}) = \bar{c}_{01} + \bar{c}_{12} + \bar{c}_{13}$ . So,  $\bar{x}_1 = \bar{c}_{01}$ . So,  $\psi_2(\hat{C}) = \psi_2(\bar{C}) = \bar{c}_{12} = c_{12}$ .

By EPM,  $\bar{x}_1 \leq \psi_1(C)$ , and  $\psi_2(\bar{C}) \leq \psi_2(C)$ . Using EF, it follows that  $\psi_1(C) = c_{01}$  and  $\psi_2(C) = c_{12}$ . Hence,  $\psi$  is unique.

The case  $c_{10} > c_{12} > c_{20}$  is similar.

**Case 3:**  $c_{20} > c_{10} > c_{12}$ .

We again introduce a third agent (say 3). Consider the matrix  $\bar{C}$ , coinciding with  $C$  on  $\{1, 2\}^+$ , and such that

$$(i) \quad \bar{c}_{32} > \bar{c}_{13} > \bar{c}_{20}.$$

$$(ii) \quad \bar{c}_{30} > \bar{c}_{10} + \bar{c}_{13}.$$

Then,  $\bar{C} \in \mathcal{C}^2$  since it has the unique m.c.s.t.  $g_N(\bar{C}) = \{(01), (12), (13)\}$ , where no two edges have the same cost.

Note that 3 is an extreme point of the m.c.s.t. corresponding to  $\bar{C}$ . Using EPM, we get

$$\psi_1(C) \geq \psi_1(\bar{C}), \psi_2(C) \geq \psi_2(\bar{C}). \quad (28)$$

Consider the reduced matrix  $\bar{C}_{\psi_1(\bar{C})}^{tr}$  on  $\{2, 3\}$ . Denote  $\bar{C}_{\psi_1(\bar{C})}^{tr} = \hat{C}$  for ease of notation. Since  $\psi_1(\bar{C}) \geq \bar{c}_{12}$  from Lemma 4, it follows that  $\bar{c}_{12} + \bar{c}_{10} - \psi_1(\bar{C}) \leq \bar{c}_{10} < \bar{c}_{20}$ , and  $\bar{c}_{12} + \bar{c}_{13} - \psi_1(\bar{C}) \leq \bar{c}_{13} < \bar{c}_{23}$ . Hence,

$$\hat{c}_{20} = \bar{c}_{12} + \bar{c}_{10} - \psi_1(\bar{C}), \hat{c}_{23} = \bar{c}_{12} + \bar{c}_{13} - \psi_1(\bar{C}), \hat{c}_{30} = \bar{c}_{13} + \bar{c}_{10} - \psi_1(\bar{C}) \quad (29)$$

Note that

$$\bar{c}_{12} + \bar{c}_{10} - \psi_1(\bar{C}) < \bar{c}_{12} + \bar{c}_{13} - \psi_1(\bar{C}) < \bar{c}_{10} + \bar{c}_{13} - \psi_1(\bar{C})$$

Hence,  $g_{\{23\}}(\hat{C}) = \{(02), (23)\}$ .

Applying case 2,  $\psi_2(\hat{C}) = \hat{c}_{20} = \bar{c}_{12} + \bar{c}_{10} - \psi_1(\bar{C})$  and  $\psi_3(\hat{C}) = \hat{c}_{23} = \bar{c}_{12} + \bar{c}_{13} - \psi_1(\bar{C})$ . Using TR,  $\psi_2(\hat{C}) = \psi_2(\bar{C})$ ,  $\psi_3(\hat{C}) = \psi_3(\bar{C})$ . Also, EF on  $\bar{C}$  gives,

$$\psi_1(\bar{C}) + \psi_2(\bar{C}) + \psi_3(\bar{C}) = \bar{c}_{10} + \bar{c}_{12} + \bar{c}_{13}$$

$$\text{or } \psi_1(\bar{C}) + (\bar{c}_{12} + \bar{c}_{10} - \psi_1(\bar{C})) + (\bar{c}_{12} + \bar{c}_{13} - \psi_1(\bar{C})) = \bar{c}_{10} + \bar{c}_{12} + \bar{c}_{13}$$

$$\text{or } \psi_1(\bar{C}) = \bar{c}_{12}$$

Hence  $\psi_2(\bar{C}) = \bar{c}_{10}$ ,  $\psi_3(\bar{C}) = \bar{c}_{13}$ . From equation 28,  $\psi_1(C) \geq \bar{c}_{12}$ ,  $\psi_2(C) \geq \bar{c}_{10}$ .

Using EF on  $C$  we can conclude that,  $\psi_1(C) = c_{12}$  and  $\psi_2(C) = c_{10}$ , i.e. the allocation is unique.

The case  $c_{10} > c_{20} > c_{12}$  is similar.

This completes the proof of the case  $|N| = 2$ .<sup>22</sup>

Suppose the theorem is true for all  $C \in \mathcal{C}_N^2$ , where  $|N| < m$ . We will show that the result is true for all  $C \in \mathcal{C}_N^2$  such that  $|N| = m$ .

Let  $C \in \mathcal{C}_N^2$ . Without loss of generality, assume  $c_{10} = \min_{k \in N} c_{k0}$ .<sup>23</sup> Thus  $(10) \in g_N(C)$ . There are two possible cases.

**Case 1:**  $c_{10} = \min_{k \in N \cup \{1\}} c_{1k}$ .

Then choose  $j \in N$  such that  $(j0) \in g_N(C)$  or  $(j1) \in g_N(C)$ .

**Case 2:**  $c_{1j} = \min_{k \in N \cup \{1\}} c_{1k}$ .

Then from Lemma 1,  $(1j) \in g_N(C)$ .

---

<sup>22</sup>Note that these three cases cover all possibilities since equality between different costs will result in the matrix not being in  $\mathcal{C}_N^2$ .

<sup>23</sup>This is unique as  $C \in \mathcal{C}_N^2$ .

In either Case 1 or 2, let  $\bar{C}$  denote the restriction of  $C$  on  $\{1, j\}$ . Then, from the case when  $\#N = 2$ , it follows that  $\psi_1(\bar{C}) = \min_{k \in N^+ \setminus \{1\}} c_{1k}$ .

Now, by iterative elimination of extreme points and repeated application of EPM, it follows that  $\psi_1(C) \leq \psi_1(\bar{C}) = \min_{k \in N^+ \setminus \{1\}} c_{1k}$ . But,  $C \in \mathcal{C}_N^2$ , and  $\psi$  satisfies EF, TR and EPM. So, from lemma 4, it follows that  $\psi_1(C) \geq \min_{k \in N^+ \setminus \{1\}} c_{1k}$ . Hence,  $\psi_1(C) = \min_{k \in N^+ \setminus \{1\}} c_{1k} = x_1$  (say).

We remove 1 to get reduced matrix  $C_{x_1}^{tr}$ . From lemma 2,  $C_{x_1}^{tr} \in \mathcal{C}^2$ . By TR,  $\psi_k(C_{x_1}^{tr}) = \psi_k(C) \forall k \neq 1$ . From the induction hypothesis, the allocation is unique on  $C_{x_1}^{tr}$  and hence on  $C$ .

This completes the proof of the theorem. ■

**Proof of Theorem 3 :** We first show that  $B$  satisfies all the three axioms. EF and EPM follow trivially from the definition. It is only necessary to show that  $B$  satisfies SR.

Let  $(10) \in g_N$ . Then,  $B_1(C) = c_{01}$ . Let us denote the reduced matrix  $C_{B_1}^{sr}$  by  $\bar{C}$ . From Lemma 3,  $\bar{C} \in \mathcal{C}^1$ . Also, the m.c.s.t. over  $N \setminus \{1\}$  corresponding to  $\bar{C}$  is

$$g_{N \setminus \{1\}} = \{(ij) \mid \text{either } (ij) \in g_N(C) \text{ with } i, j \neq 1 \text{ or } (ij) = (l0) \text{ where } (1l) \in g_N(C)\}.$$

Also, for all  $i, j \in N \setminus \{1\}$ ,  $\bar{c}_{ij} = c_{ij}$  if  $(ij) \in g_N$ , and for  $k \in N \setminus \{1\}$ ,  $\bar{c}_{k0} = c_{1k}$  if  $(1k) \in g_N(C)$ . Hence, for all  $k \in N \setminus \{1\}$ ,  $\bar{c}_{k\bar{\alpha}(k)} = c_{k\alpha(k)}$ , where  $\bar{\alpha}(k)$  is the immediate predecessor of  $k$  in  $g_{N \setminus \{1\}}$ . So,  $B_k(\bar{C}) = B_k(C)$  for all  $k \in N \setminus \{1\}$  and  $B$  satisfies Source Consistency.

Next, we show that  $B$  is the only rule over  $\mathcal{C}^1$  which satisfies all the three axioms. This proof is by induction on the cardinality of the set of agents.

We remark that the proof for the case  $|N| = 2$  is virtually identical to that of Theorem 2, with SR replacing TR and Lemma 5 replacing Lemma 4.

Suppose  $B$  is the only rule satisfying the three axioms, for all  $C \in \mathcal{C}^1$ , where  $|N| < m$ . We will show that the result is true for all  $C \in \mathcal{C}^1$  such that  $|N| = m$ .

Let  $C \in \mathcal{C}^1$ . Without loss of generality, assume  $(10) \in g_N(C)$ . There are two possible cases.

**Case 1 :** There are at least two extreme points of  $C$ , say  $m_1$  and  $m_2$ .

First, remove  $m_1$  and consider the matrix  $C^{m_1}$ , which is the restriction of  $C$  over  $(N^+ \setminus \{m_1\})$ . By EPM,  $\psi_i(C) \leq \psi_i(C^{m_1})$  for all  $i \neq m_1$ . As  $C^{m_1}$  has  $(m-1)$  agents, the induction hypothesis gives  $\psi_i(C^{m_1}) = c_{i\alpha(i)}$ . So,  $\psi_i(C) \leq c_{i\alpha(i)} \forall i \neq m_1$ . Similarly by eliminating  $m_2$  and using EPM, we get  $\psi_i(C) \leq c_{i\alpha(i)} \forall i \neq m_2$ . Combining the two, we get  $\psi_i(C) \leq c_{i\alpha(i)} \forall i \in N$ .

But from EF, we know that  $\sum_{i \in N} \psi_i(C) = c(N) = \sum_{i \in N} c_{i\alpha(i)}$ . Therefore  $\psi_i(C) = c_{i\alpha(i)} \forall i \in N$ , and hence the allocation is unique.

**Case 2:** If there is only one extreme point of  $C$ , then  $g_N(C)$  must be a line, i.e. each agent has at most one follower. Without loss of generality, assume 1 is connected to 2 and 0. Let  $\bar{C}$  be the restriction of  $C$  over the set  $\{0, 1, 2\}$ . By iterative elimination of the extreme points and use of EPM we get  $\psi_i(C) \leq \psi_i(\bar{C})$ . Using the induction hypothesis, we get  $\psi_1(C) \leq c_{10}$  and  $\psi_2(C) \leq c_{12}$ .

Suppose  $\psi_1(C) = x_1 = c_{10} - \epsilon$ , where  $\epsilon \geq 0$ . Now consider the reduced matrix  $C_{x_1}^{sr}$ , which will be denoted by  $\hat{C}$ . It can be easily checked that  $g_{N \setminus \{1\}}$  is also a line where 2 is connected to 0. Thus  $\psi_2(\hat{C}) = \hat{c}_{20} = \min(c_{20}, c_{12} + c_{10} - \psi_1(C)) = \min(c_{20}, c_{12} + \epsilon)$ . So,  $\psi_2(\hat{C}) \geq c_{12}$  with equality holding only if  $\epsilon = 0$ . By SR,  $\psi_2(C) = \psi_2(\hat{C})$ . But from EPM  $\psi_2(C) \leq \psi_2(\bar{C}) = c_{12}$ . This is possible only if  $\epsilon = 0$ . Therefore,  $\psi_1(C) = c_{10}$ . Using SR and the induction hypothesis, we can conclude that  $\psi = B$ .

■

## REFERENCES

1. C. G. Bird, On cost allocation for a spanning tree : A game theoretic approach, *Networks* 6 (1976), 335-350.
2. M. Davis and M. Maschler, The core of a cooperative game, *Naval Research Logistics Quarterly* 12 (1965), 223-259.
3. V. Feltkamp, "Cooperation in controlled network structures", Ph.D. thesis, NWO, Netherlands, 1995.
4. R. L. Graham and P. Hell, On the history of the minimum spanning tree problem, *Annals of the History of Computing* 7 (1985), 43-57.
5. D. Granot and F. Granot, On computational complexity of a cost allocation approach to a fixed cost spanning forest problem, *Mathematics of Operations Research* 17 (1993), 765-780.
6. D. Granot and G. Huberman, Minimum cost spanning tree games, *Mathematical Programming* 21 (1981), 1-18.
7. D. Granot and G. Huberman, On the core and Nucleolus of the minimum cost spanning tree games, *Mathematical Programming* 29 (1984), 323-347.
8. D. Granot and M. Maschler, Spanning Network Games, *International Journal of Game Theory* 27 (1998), 467-500.
9. S. Hart and A. Mas-Colell, Potential, value and consistency, *Econometrica* 57 (1989), 589-614.
10. A. Kar, Axiomatization of the Shapley value on minimum cost spanning tree games, *Games and Economic Behavior*, 38 (2002).

11. H. Moulin, Axiomatic cost and surplus-sharing, *in* “Handbook of Social Choice and Welfare” (K.J. Arrow, A.K. Sen, K. Suzumura, Eds.), North-Holland, 1999.
12. B. Peleg, On the reduced game property and its converse, *International Journal of Game Theory* 15 (1986), 187-200.
13. R. C. Prim, Shortest connection networks and some generalizations, *Bell Systems Technology Journal* 36 (1957), 1389-1401.
14. W. W. Sharkey, Network models in economics, *in* “Handbooks in Operation Research and Management Science” (M. O. Ball et al. Eds.), Elsevier Science, 1995.
15. W. Thomson, Consistency and its converse, University of Rochester, Mimeo, (1998).
16. H. P. Young, Cost allocation, *in* “Handbook of Game Theory with Economic Applications” (R. Aumann, S. Hart, Eds.), North Holland, 1994.

