

# A $q$ -analogue of the distance matrix of a tree <sup>1</sup>

R. B. Bapat <sup>2</sup>

A. K. Lal<sup>3</sup>

Sukanta Pati <sup>4</sup>

## Abstract

We consider a  $q$ -analogue of the distance matrix (called the  $q$ -distance matrix) of an unweighted tree and give formulae for the inverse and the determinant, which generalize the existing formulae for the distance matrix. We obtain the Smith normal form of the  $q$ -distance matrix of a tree. The relationship of the  $q$ -distance matrix with the Laplacian matrix leads to  $q$ -analogue of the Laplacian matrix of a tree, some of whose properties are also given. We study another matrix related to the distance matrix (the *exponential distance matrix*) and show its relationship with the  $q$ -Laplacian and the  $q$ -distance matrix. A formula for the determinant of the  $q$ -distance matrix of a weighted tree is also given.

*Keywords:* Tree, Distance Matrix, Laplacian Matrix, Determinant, Block Matrix.

*Mathematics Subject Classification:* Primary: 05C50 Secondary: 15A15.

## 1 Introduction

In this paper, we consider a  $q$ -analogue of the distance matrix of a tree and call it the  $q$ -distance matrix. The inverse and the determinant of the matrix are obtained when the tree is unweighted. We also define some related matrices and study their properties. For a weighted tree, we obtain a formula for the determinant of the  $q$ -distance matrix.

We refer the reader to the book by Harary [6] for basic definitions and terminology in graph theory. We start with some definitions. A *tree* is a simple connected graph without any circuit. A *weighted tree* is a tree in which each edge is assigned a *weight*, which is a positive number. So, an *unweighted tree* is simply a tree with each edge having weight 1.

Let  $\mathbf{e}, \mathbf{0}$  be the column vectors consisting of all ones and all zeros, respectively. Let  $J = \mathbf{e}\mathbf{e}^t$  be the matrix of all ones. For a tree  $T$  on  $n$  vertices, let  $\mathbf{d}^t = (d_1, d_2, \dots, d_n)$ ,  $\delta = 2\mathbf{e} - \mathbf{d}$  and  $\mathbf{z} = \mathbf{d} - \mathbf{e}$ , where  $d_i$  is the degree of the  $i^{\text{th}}$  vertex of  $T$ . Note that  $\delta + \mathbf{z} = \mathbf{e}$ .

---

<sup>1</sup>The paper was written when the last two authors visited the Stat-Math Unit at Indian Statistical Institute, Delhi Centre in June 2005. The two authors take this opportunity to thank the Stat-Math Unit at Indian Statistical Institute, Delhi Centre for the visiting assignment.

<sup>2</sup>Stat-Math Unit, Indian Statistical Institute Delhi, 7-SJSS Marg, New Delhi - 110 016, India;  
e-mail: rbb@isid.ac.in

<sup>3</sup>Corresponding Author: A. K. Lal, Indian Institute of Technology Kanpur, Kanpur - 208 016, India;  
e-mail: arlal@iitk.ac.in.

<sup>4</sup>Department of Mathematics, Indian Institute of Technology, Guwahati, India;  
e-mail: pati@iitg.ac.in

Let  $T$  be a tree on  $n$  vertices. The distance matrix of a tree  $T$  is a  $n \times n$  matrix  $D$  with  $D_{ij} = k$ , if the path from the vertex  $i$  to the vertex  $j$  is of length  $k$ ; and  $D_{ii} = 0$ . The *Laplacian matrix*,  $L$ , of a tree  $T$  is defined by  $L = \text{diag}(\mathbf{d}) - A$ , where  $A$  is the adjacency matrix of  $T$ .

The distance matrix of a tree is extensively investigated in the literature. The first known result concerns the determinant of the matrix  $D$  (see Graham and Pollak [5]), which asserts that if  $T$  is any tree on  $n$  vertices then  $\det(D) = (-1)^{n-1}(n-1)2^{n-2}$ . Thus,  $\det(D)$  is a function dependent on only  $n$ , the number of vertices of the tree. The formula for the inverse of the matrix  $D$  was obtained in a subsequent paper by Graham and Lovasz [4]. Their result was extended for a weighted tree by Bapat *et al* [1]. In Section 2, we extend the result of Graham and Lovasz by considering a new distance matrix, termed the  $q$ -distance matrix, denoted  $\mathcal{D} = (\mathcal{D}_{ij})$  and defined as follows:

Let  $T$  be a tree on  $n$  vertices and  $D = (D_{ij})$  be its classical distance matrix. For an indeterminate  $q$ , we define

$$\mathcal{D}_{ij} = \begin{cases} 1 + q + q^2 + \cdots + q^{k-1} & \text{if } D_{ij} = k \\ 0 & \text{if } i = j \end{cases}.$$




Figure 1: An Unweighted Tree on 6 vertices

For example, the distance matrix  $\mathcal{D}$ , of a tree  $T$  shown in Figure 1 is given by

$$\mathcal{D} = \begin{bmatrix} 0 & 1 & 1+q & 1+q & 1+q+q^2 & 1+q+q^2 \\ 1 & 0 & 1 & 1 & 1+q & 1+q \\ 1+q & 1 & 0 & 1+q & 1+q+q^2 & 1+q+q^2 \\ 1+q & 1 & 1+q & 0 & 1 & 1 \\ 1+q+q^2 & 1+q & 1+q+q^2 & 1 & 0 & 1+q \\ 1+q+q^2 & 1+q & 1+q+q^2 & 1 & 1+q & 0 \end{bmatrix}.$$

Each element of  $\mathcal{D}$  is a polynomial in the indeterminate  $q$ . For convenience we denote the matrix simply by  $\mathcal{D}$  and suppress the dependence on  $q$  in the notation. Observe that  $\mathcal{D}$  is an entrywise nonnegative matrix for all  $q \geq -1$ .

In Section 2, we obtain an expression for  $\mathcal{D}^{-1}$  when  $q \neq -1$ . In Section 3, we use the expression for  $\mathcal{D}^{-1}$  to define a generalization, called the  $q$ -Laplacian, corresponding to the

Laplacian matrix  $L$  of a tree. We also define a related matrix, the exponential distance matrix, and examine its properties in relation to the Laplacian. Section 4 deals with the invariant factors and Smith normal form of the  $q$ -distance matrix. The determinant of the  $q$ -distance matrix for a weighted tree is given in Section 5. The formula contains the classical formula of [5] as a special case.

## 2 $q$ -distance matrix of a tree

In this section, we extend certain results on distance matrices obtained by Graham and Pollak [5] and Graham and Lovasz [4].

Most of the proofs in this paper are based on mathematical induction on the number of vertices of a tree  $T$ . So, in the induction step, we start with a tree  $\bar{T}$  having a pendant vertex  $k+1$  with vertex  $k$  adjacent to it. The tree  $T$  is defined as  $\bar{T} \setminus \{k+1\}$ . Then, using the matrices  $\mathcal{D}, L, \mathbf{z}$  corresponding to the tree  $T$ , we define the corresponding matrices  $\bar{\mathcal{D}}, \bar{L}$  and  $\bar{\mathbf{z}}$  of the tree  $\bar{T}$ . That is, we have

$$\bar{\mathcal{D}} = \begin{bmatrix} \mathcal{D} & \mathbf{e} + q\mathcal{D}\mathbf{e}_k \\ \mathbf{e}^t + q\mathbf{e}_k^t\mathcal{D} & 0 \end{bmatrix}, \bar{L} = \begin{bmatrix} L + \mathbf{e}_k\mathbf{e}_k^t & -\mathbf{e}_k \\ -\mathbf{e}_k^t & 1 \end{bmatrix}, \bar{\mathbf{z}} = \begin{bmatrix} \mathbf{z} + \mathbf{e}_k \\ 0 \end{bmatrix}. \quad (2.1)$$

We start with the main result of this section.

**THEOREM 2.1** *Let  $\mathcal{D}$  be the  $q$ -distance matrix of a tree on  $n$  vertices and  $q \neq -1$ . Then*

$$\mathbf{e} = \frac{1}{n-1}\mathcal{D}(\mathbf{e} - q\mathbf{z}). \quad (2.2)$$

*Also,  $\mathcal{D}$  is invertible, and*

$$\mathcal{D}^{-1} = \frac{1}{(n-1)(1+q)}\mathcal{U} - \frac{1}{1+q}\mathcal{L}, \quad (2.3)$$

*where  $\mathcal{L} = qL - (q-1)I + q(q-1)\text{diag}(\mathbf{z})$  and  $\mathcal{U} = (\mathbf{e} - q\mathbf{z})(\mathbf{e} - q\mathbf{z})^t$ .*

*Proof.* We prove the result by induction on  $n$ . Let  $n = 2$ . In this case, the matrices  $\mathcal{D}, L$  and  $\mathbf{z}$  are defined as follows:

$$\mathcal{D} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, L = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}, \mathbf{z} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

So,  $\frac{1}{n-1}\mathcal{D}(\mathbf{e} - q\mathbf{z}) = \mathcal{D}\mathbf{e} = \mathbf{e}$ . Thus, (2.2) is true for  $n = 2$ . Also, for  $n = 2$  and  $q \neq -1$ , the right hand side of (2.3) reduces to

$$\begin{aligned} \frac{1}{1+q}(qJ - (q-1)J) - \frac{1}{q+1}(qL - (q-1)I) &= \frac{1}{q+1}(-q(I - \mathcal{D}) + (q-1)I + J) \\ &= \frac{1}{q+1}(q\mathcal{D} - I + J) = \frac{1}{q+1}(q\mathcal{D} + \mathcal{D}) \\ &= \mathcal{D} = \mathcal{D}^{-1}. \end{aligned}$$

Hence, (2.3) holds for  $n = 2$ . We now assume that both the results are true for  $n = k$ . Let us prove the result for  $n = k + 1$ .

We first prove (2.2). That is, we need to show that  $\bar{\mathcal{D}}(\mathbf{e} - q\bar{\mathbf{z}}) = k\mathbf{e}$ . From now on, we will use the expressions for  $\bar{\mathcal{D}}$ ,  $\bar{L}$ ,  $\bar{\mathbf{z}}$  from (2.1). In this case, we have,

$$\begin{aligned}\bar{\mathcal{D}}(\mathbf{e} - q\bar{\mathbf{z}}) &= \begin{bmatrix} \mathcal{D} & \mathbf{e} + q\mathcal{D}\mathbf{e}_k \\ \mathbf{e}^t + q\mathbf{e}_k^t\mathcal{D} & 0 \end{bmatrix} \begin{bmatrix} \mathbf{e} - q(\mathbf{z} + \mathbf{e}_k) \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} \mathcal{D}(\mathbf{e} - q\mathbf{z}) - q\mathcal{D}\mathbf{e}_k + \mathbf{e} + q\mathcal{D}\mathbf{e}_k \\ (\mathbf{e}^t + q\mathbf{e}_k^t\mathcal{D})(\mathbf{e} - q(\mathbf{z} + \mathbf{e}_k)) \end{bmatrix}. \end{aligned} \quad (2.4)$$

We calculate the two blocks separately using the induction hypothesis. The first block is given by

$$\mathcal{D}(\mathbf{e} - q\mathbf{z}) - q\mathcal{D}\mathbf{e}_k + \mathbf{e} + q\mathcal{D}\mathbf{e}_k = (k - 1)\mathbf{e} + \mathbf{e} = k\mathbf{e}. \quad (2.5)$$

Note that  $\mathbf{e}^t\mathbf{z} = \mathbf{e}^t(\mathbf{d} - \mathbf{e}) = 2(k - 1) - k = k - 2$ . So,

$$\mathbf{e}^t(\mathbf{e} - q\mathbf{z}) = k - q(k - 2). \quad (2.6)$$

Therefore, using (2.6), the second block reduces to

$$\begin{aligned}(\mathbf{e}^t + q\mathbf{e}_k^t\mathcal{D})(\mathbf{e} - q(\mathbf{z} + \mathbf{e}_k)) &= k - q(k - 2) - q + q\mathbf{e}_k^t\mathcal{D}(\mathbf{e} - q\mathbf{z}) - q^2\mathbf{e}_k^t\mathcal{D}\mathbf{e}_k \\ &= k - q(k - 1) + q\mathbf{e}_k^t(k - 1)\mathbf{e} - q^2 \cdot 0 = k. \end{aligned} \quad (2.7)$$

Therefore, by substituting the results from (2.5) and (2.7) in (2.4), the proof of (2.2) is complete, as

$$\bar{\mathcal{D}}(\mathbf{e} - q\bar{\mathbf{z}}) = \begin{bmatrix} k\mathbf{e} \\ k \end{bmatrix} = k\mathbf{e}.$$

Under the assumption that  $q \neq -1$ , we now prove that the matrix  $\bar{\mathcal{D}}^{-1}$  is indeed given by (2.3). By the induction hypothesis, we assume that  $\mathcal{D}$  is an invertible matrix and use it to show that  $\bar{\mathcal{D}}$  is invertible. From (2.1), note that  $\bar{\mathcal{D}}$  is a block matrix and is given by

$$\bar{\mathcal{D}} = \begin{bmatrix} \mathcal{D} & \mathbf{e} + q\mathcal{D}\mathbf{e}_k \\ \mathbf{e}^t + q\mathbf{e}_k^t\mathcal{D} & 0 \end{bmatrix}.$$

Thus, if  $\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$  is the inverse of  $\bar{\mathcal{D}}$ , then we need to show that

$$A_{11} = \mathcal{D}^{-1} + \mathcal{D}^{-1}(\mathbf{e} + q\mathcal{D}\mathbf{e}_k)W^{-1}(\mathbf{e} + q\mathcal{D}\mathbf{e}_k)^t\mathcal{D}^{-1} \quad (2.8)$$

$$\text{and } A_{12} = -\mathcal{D}^{-1}(\mathbf{e} + q\mathcal{D}\mathbf{e}_k)W^{-1} \quad (2.9)$$

where  $W = 0 - (\mathbf{e} + q\mathcal{D}\mathbf{e}_k)^t\mathcal{D}^{-1}(\mathbf{e} + q\mathcal{D}\mathbf{e}_k) = -(\mathbf{e} + q\mathcal{D}\mathbf{e}_k)^t\mathcal{D}^{-1}(\mathbf{e} + q\mathcal{D}\mathbf{e}_k)$  is a  $1 \times 1$  matrix. From the induction hypothesis and (2.2), observe that  $\mathcal{D}^{-1}\mathbf{e} = \frac{1}{k-1}(\mathbf{e} - q\mathbf{z})$ .

Therefore, using (2.6), we get

$$\begin{aligned}
-W &= (\mathbf{e}^t + q\mathbf{e}_k^t \mathcal{D}) \mathcal{D}^{-1} (\mathbf{e} + q\mathcal{D}\mathbf{e}_k) = \mathbf{e}^t \mathcal{D}^{-1} \mathbf{e} + q\mathbf{e}^t \mathbf{e}_k + q\mathbf{e}_k^t \mathbf{e} + q^2 \mathbf{e}_k^t \mathcal{D}\mathbf{e}_k \\
&= \frac{1}{k-1} (\mathbf{e}^t (\mathbf{e} - q\mathbf{z})) + 2q + q^2 \cdot 0 = \frac{k}{k-1} (1+q).
\end{aligned} \tag{2.10}$$

We will prove (2.8) and (2.9) in two steps.

**Step 1:** Using (2.10) and the induction hypothesis,

$$\begin{aligned}
A_{11} &= \mathcal{D}^{-1} + \mathcal{D}^{-1} (\mathbf{e} + q\mathcal{D}\mathbf{e}_k) W^{-1} (\mathbf{e} + q\mathcal{D}\mathbf{e}_k)^t \mathcal{D}^{-1} \\
&= \mathcal{D}^{-1} - \frac{k-1}{k(1+q)} [(\mathcal{D}^{-1} \mathbf{e} + q\mathbf{e}_k) (\mathbf{e}^t \mathcal{D}^{-1} + q\mathbf{e}_k^t)] \\
&= \frac{\mathcal{U}}{(k-1)(1+q)} - \frac{\mathcal{L}}{1+q} - \frac{k-1}{k(1+q)} \left[ \frac{(\mathbf{e} - q\mathbf{z})(\mathbf{e} - q\mathbf{z})^t}{(k-1)^2} \right. \\
&\quad \left. + \frac{q(\mathbf{e} - q\mathbf{z})\mathbf{e}_k^t}{k-1} + \frac{q\mathbf{e}_k(\mathbf{e} - q\mathbf{z})^t}{k-1} + q^2 \mathbf{e}_k \mathbf{e}_k^t \right] \\
&= \frac{\mathcal{U}}{k(1+q)} - \frac{\mathcal{L}}{1+q} - \frac{q((\mathbf{e} - q\mathbf{z})\mathbf{e}_k^t + \mathbf{e}_k(\mathbf{e} - q\mathbf{z})^t)}{k(1+q)} - \frac{(k-1)q^2 \mathbf{e}_k \mathbf{e}_k^t}{k(1+q)},
\end{aligned} \tag{2.11}$$

and

$$\begin{aligned}
A_{12} &= -\mathcal{D}^{-1} (\mathbf{e} + q\mathcal{D}\mathbf{e}_k) W^{-1} = \left( \frac{1}{k-1} (\mathbf{e} - q\mathbf{z}) + q\mathbf{e}_k \right) \frac{k-1}{k(1+q)} \\
&= \frac{1}{k(1+q)} (\mathbf{e} - q\mathbf{z} + q(k-1)\mathbf{e}_k).
\end{aligned} \tag{2.12}$$

**Step 2:** We now determine the matrices  $\bar{\mathcal{L}}$  and  $\bar{\mathcal{U}}$ . Using (2.1) and (2.3), we have

$$\begin{aligned}
\bar{\mathcal{L}} &= q\bar{\mathcal{L}} - (q-1)\bar{I} + q(q-1)\text{diag}(\bar{\mathbf{z}}) \\
&= q \left[ \begin{array}{c|c} L + \mathbf{e}_k \mathbf{e}_k^t & -\mathbf{e}_k \\ \hline -\mathbf{e}_k^t & 1 \end{array} \right] - \left[ \begin{array}{c|c} I & \mathbf{0} \\ \hline \mathbf{0}^t & 1 \end{array} \right] + q(q-1) \left[ \begin{array}{c|c} \text{diag}(\mathbf{z}) & \mathbf{0} \\ \hline \mathbf{0}^t & 0 \end{array} \right] \\
&= \left[ \begin{array}{c|c} \mathcal{L} + q^2 \mathbf{e}_k \mathbf{e}_k^t & -q\mathbf{e}_k \\ \hline -q\mathbf{e}_k^t & 1 \end{array} \right],
\end{aligned} \tag{2.13}$$

and

$$\begin{aligned}
\bar{\mathcal{U}} &= (\bar{\mathbf{e}} - q\bar{\mathbf{z}})(\bar{\mathbf{e}} - q\bar{\mathbf{z}})^t = \left[ \begin{array}{c} \mathbf{e} - q(\mathbf{z} + \mathbf{e}_k) \\ 1 \end{array} \right] \left[ \begin{array}{c|c} (\mathbf{e} - q(\mathbf{z} + \mathbf{e}_k))^t & 1 \end{array} \right] \\
&= \left[ \begin{array}{c|c} \mathcal{U} - q((\mathbf{e} - q\mathbf{z})\mathbf{e}_k^t + \mathbf{e}_k(\mathbf{e} - q\mathbf{z})^t) + q^2 \mathbf{e}_k \mathbf{e}_k^t & (\mathbf{e} - q\mathbf{z}) - q\mathbf{e}_k \\ \hline ((\mathbf{e} - q\mathbf{z}) - q\mathbf{e}_k)^t & 1 \end{array} \right].
\end{aligned} \tag{2.14}$$

Thus, using (2.13) and (2.14), the first block of the matrix  $\bar{\mathcal{D}}^{-1}$  is given by

$$\begin{aligned}
\frac{1}{k(1+q)} \bar{\mathcal{U}} - \frac{1}{1+q} \bar{\mathcal{L}} &= \frac{\mathcal{U}}{k(1+q)} - \frac{\mathcal{L}}{1+q} \\
&\quad - \frac{q((\mathbf{e} - q\mathbf{z})\mathbf{e}_k^t + \mathbf{e}_k(\mathbf{e} - q\mathbf{z})^t)}{k(1+q)} - \frac{(k-1)q^2 \mathbf{e}_k \mathbf{e}_k^t}{k(1+q)},
\end{aligned} \tag{2.15}$$

and the second block of the matrix  $\bar{D}^{-1}$  is given by

$$\frac{1}{k(1+q)}((\mathbf{e} - q\mathbf{z}) - q\mathbf{e}_k) - \frac{1}{1+q}(-q\mathbf{e}_k) = \frac{1}{k(1+q)}(\mathbf{e} - q\mathbf{z} + q(k-1)\mathbf{e}_k). \quad (2.16)$$

The expressions (2.11) and (2.12) are respectively, equal to the expressions (2.15) and (2.16). Hence, if the two sides of (2.3) are partitioned conformally as in (2.1), then the  $(1, 1)$  and  $(1, 2)$  blocks on both sides are equal. By symmetry, the  $(2, 1)$  block on both sides are also equal. Since a tree has at least two pendant vertices, we can repeat the argument using the second pendant vertex and thus conclude that the  $(2, 2)$  block on both sides of (2.3) are equal. Thus, by the induction hypothesis, we obtain the required result.  $\blacksquare$

For  $q = 1$ , the Theorem 2.1 reduces to the inverse of the distance matrix  $D$ , obtained by Graham and Lovasz [4].

**COROLLARY 2.2** *Let  $T$  be a tree on  $n$  vertices and let  $D$  be its distance matrix. Then*

$$D^{-1} = \frac{1}{2(n-1)}(\mathbf{e} - \mathbf{z})(\mathbf{e} - \mathbf{z})^t - \frac{1}{2}L = \frac{1}{2(n-1)}\delta\delta^t - \frac{1}{2}L.$$

### 3 Exponential distance matrix of a tree

We now define another matrix using the distance matrix of a tree. Let  $T$  be a tree on  $n$  vertices and let  $D = (D_{ij})$  be its distance matrix. We now consider an  $n \times n$  matrix  $F = (F_{ij})$ , called the exponential distance matrix, with  $F_{ij} = \begin{cases} 1 & \text{if } i = j \\ q^{D_{ij}} & \text{if } i \neq j \end{cases}$ .

**PROPOSITION 3.3** *Let  $T$  be a tree on  $n$  vertices and  $F$  be the corresponding exponential distance matrix. If  $q \neq \pm 1$  then*

$$F^{-1} = I - \frac{q}{1-q^2}A + \frac{q^2}{1-q^2} \text{diag}(\mathbf{d}).$$

*Proof.* We will prove the result by induction on  $n$ . The result can be easily verified for  $n = 2$ . Let the result be true for  $n = k$ , and let  $\bar{T}$  be a tree on  $k+1$  vertices with  $k+1$  being a pendant vertex and the vertex  $k$  being adjacent to  $k+1$ . As before, let the tree  $T = \bar{T} \setminus \{k+1\}$ . Suppose  $\bar{F}, F$  respectively, represent the matrices corresponding to the trees  $\bar{T}$  and  $T$ . Then

$$\bar{F} = \begin{bmatrix} F & \mathbf{q} \\ \mathbf{q}^t & 1 \end{bmatrix},$$

where for any  $q \in \mathbb{R}$ ,

$$\mathbf{q}^t = (q^{D_{1,k+1}}, q^{D_{2,k+1}}, \dots, q^{D_{k,k+1}}, q^{D_{k+1,k+1}} = q^0 = 1). \quad (3.17)$$

We are now ready to prove the formula for  $\bar{F}^{-1}$ . Note that by induction hypothesis, for  $q \neq \pm 1$ ,  $F$  is already invertible. So, if  $\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$  is the inverse of  $\bar{F}$ , then we need to show that

$$A_{11} = F^{-1} + F^{-1}\underline{\mathbf{q}}W^{-1}(F^{-1}\underline{\mathbf{q}})^t, \quad \text{and} \quad A_{12} = -F^{-1}\underline{\mathbf{q}}W^{-1}, \quad (3.18)$$

where  $W = 1 - \underline{\mathbf{q}}^t F^{-1} \underline{\mathbf{q}}$ . As  $F\mathbf{e}_k = \frac{1}{q}\underline{\mathbf{q}}$ ,  $W = 1 - q^2$ . Thus,

$$A_{11} = F^{-1} + \frac{1}{1 - q^2}(q\mathbf{e}_k)(q\mathbf{e}_k)^t = F^{-1} + \frac{q^2}{1 - q^2}\mathbf{e}_k\mathbf{e}_k^t \quad \text{and} \quad A_{12} = -\frac{q}{1 - q^2}\mathbf{e}_k. \quad (3.19)$$

Also, from the statement of the proposition and (3.19),

$$\begin{aligned} \bar{F}_{11}^{-1} &= I - \frac{q}{1 - q^2}A + \frac{q^2}{1 - q^2}\text{diag}(\mathbf{d} + \mathbf{e}_k) \\ &= I - \frac{q}{1 - q^2}A + \frac{q^2}{1 - q^2}\text{diag}(\mathbf{d}) + \frac{q^2}{1 - q^2}\mathbf{e}_k\mathbf{e}_k^t = F^{-1} + \frac{q^2}{1 - q^2}\mathbf{e}_k\mathbf{e}_k^t \\ &= A_{11} \end{aligned} \quad (3.20)$$

and

$$\bar{F}_{12}^{-1} = \mathbf{0} - \frac{q}{1 - q^2}\mathbf{e}_k = A_{12}. \quad (3.21)$$

Therefore, from (3.20) and (3.21), if the two sides of  $F^{-1}$  are partitioned conformally as in (2.1), then the (1,1) and (1,2) blocks on both sides are equal. By symmetry, the (2,1) block on both sides are also equal. Since a tree has at least two pendant vertices, we can repeat the argument using the second pendant vertex and thus conclude that the (2,2) block on both sides of (2.3) are equal. So, by the induction hypothesis, the required result follows.  $\blacksquare$

Comparing the expression for  $\mathcal{D}^{-1}$  given in (2.3) with the one given by Bapat *et al* (see (2.1) in [1]), we introduce the  $q$ -Laplacian matrix,  $\mathcal{L}$ , of a tree  $T$  by

$$\mathcal{L} = qL - (q - 1)I + q(q - 1)\text{diag}(\mathbf{z}). \quad (3.22)$$

That is, if  $v_i, v_j$  are any two vertices of the tree  $T$ , then

$$\mathcal{L}_{ij} = \begin{cases} 1 + (\deg(v_i) - 1)q^2 & \text{if } i = j \\ -q & \text{if } i \neq j, (v_i, v_j) \in E(T) \\ 0 & \text{if } i \neq j, (v_i, v_j) \notin E(T) \end{cases}.$$

**REMARK 1** The  $q$ -Laplacian matrix  $\mathcal{L}$  reduces to

1.  $\mathcal{L} = \text{diag}(d) - A = L$ , the Laplacian matrix of a tree whenever  $q = 1$ .
2.  $\mathcal{L} = \text{diag}(d) + A$ , the signless Laplacian matrix of a tree (see [7]), whenever  $q = -1$ .

We now state a few properties of the  $q$ -Laplacian matrix  $\mathcal{L}$ .

**PROPOSITION 3.4** *Let  $T$  be a tree on  $n$  vertices and let  $\mathcal{L}$  be the  $q$ -Laplacian matrix. Then*

1.  $\det(\mathcal{L}) = 1 - q^2$ .
2. *the matrix  $\mathcal{L}$  is positive definite if and only if  $q \in (-1, 1)$ .*

*Proof.* We use induction to prove both parts of the proposition. The result is clearly true for  $n = 2$  as the corresponding matrix is given by  $\mathcal{L} = \begin{bmatrix} 1 & -q \\ -q & 1 \end{bmatrix}$  and

$$\det(\mathcal{L}) = 1 - q^2 > 0 \text{ if and only if } q \in (-1, 1).$$

Let us assume the result to be true for  $n = k$ . We now prove the result for  $n = k + 1$ . As before, let  $\bar{T}$  be a tree on  $k + 1$  vertices. Let  $k + 1$  be a pendant vertex adjacent to vertex  $k$ . Then in the block form,  $\bar{\mathcal{L}}$  is given by

$$\bar{\mathcal{L}} = \begin{bmatrix} q^2 \mathbf{e}_k \mathbf{e}_k^t + \mathcal{L} & -q \mathbf{e}_k \\ -q \mathbf{e}_k^t & 1 \end{bmatrix}.$$

Thus, by the induction hypothesis

$$\det(\bar{\mathcal{L}}) = 1 \cdot \det(q^2 \mathbf{e}_k \mathbf{e}_k^t + \mathcal{L} - (-q \mathbf{e}_k) \cdot 1^{-1} \cdot (-q \mathbf{e}_k^t)) = \det(\mathcal{L}) = 1 - q^2. \quad (3.23)$$

Hence by the induction argument the proof of the first part is complete.

For the proof of the second part, observe that, by the induction hypothesis,  $\mathcal{L}$  is a positive definite matrix. So, the matrix  $q^2 \mathbf{e}_k \mathbf{e}_k^t + \mathcal{L}$  is also a positive definite matrix.

We now suppose that  $\bar{\mathcal{L}}$  is a positive definite matrix. Then  $\det(\bar{\mathcal{L}}) = 1 - q^2$  must be positive. That is, we need  $q \in (-1, 1)$ .

If  $q \in (-1, 1)$ , then  $\det(\bar{\mathcal{L}}) = 1 - q^2 > 0$ . Also, by the induction argument, the matrix  $q^2 \mathbf{e}_k \mathbf{e}_k^t + \mathcal{L}$ , which corresponds to the first block of the matrix  $\bar{\mathcal{L}}$ , is positive definite. Hence, the matrix  $\bar{\mathcal{L}}$  is itself a positive definite matrix.

Therefore, by the induction argument the proof of the second part is also complete.

■

The proof of the following corollary is omitted as it is an immediate consequence of Proposition 3.4 and Remark 1.

**COROLLARY 3.5** *Let  $T$  be a tree on  $n$  vertices. Then the  $q$ -Laplacian matrix  $\mathcal{L}$  of  $T$  is positive semidefinite for  $q = -1, 1$ .*

The next proposition gives a bound on the smallest eigenvalue of the  $q$ -Laplacian matrix  $\mathcal{L}$ .



**PROPOSITION 3.6** *Let  $T$  be a tree and let  $\mathcal{L}$  be the  $q$ -Laplacian matrix. If  $\tau(\mathcal{L})$  denotes the smallest eigenvalue of  $\mathcal{L}$ , then  $\tau(\mathcal{L}) \leq 1$  for all  $q \in \mathbb{R}$ . Also,  $\tau(\mathcal{L}) = 1$  if and only if  $q = 0$ .*

*Proof.* If  $q = 0$  then  $\mathcal{L} = I$  and hence  $\tau(\mathcal{L}) = 1$ . For  $q \neq 0$ , consider a tree  $T$  with 1 as a pendant vertex. Suppose the vertex 2 is adjacent to 1 and has degree  $d$ . Then the  $2 \times 2$  matrix  $M = \begin{bmatrix} 1 & -q \\ -q & 1 + (d-1)q^2 \end{bmatrix}$  is a submatrix of  $\mathcal{L}$ . Note that, the characteristic polynomial of this submatrix is  $p(\lambda) = (\lambda - 1)^2 + q^2(d - 2 - (d-1)\lambda)$ . Note that  $p(\frac{d-2}{d-1}) > 0$  and  $p(1) < 0$  ( $q \neq 0$ ). As  $p(\lambda)$  is a continuous function of  $\lambda$ , by the intermediate value theorem, there exists a real number  $x_0 \in (\frac{d-2}{d-1}, 1)$  such that  $p(x_0) = 0$ . So, by the interlacing eigenvalue theorem  $\tau(\mathcal{L}) < 1$ . Therefore, the required result follows.  $\blacksquare$

We now show that for  $|q| > 1$ ,  $\mathcal{L}$  has exactly one negative eigenvalue.

**PROPOSITION 3.7** *Let  $T$  be a tree and let  $\mathcal{L}$  be the  $q$ -Laplacian matrix. Then for  $|q| > 1$ ,  $\mathcal{L}$  has exactly one negative eigenvalue.*

*Proof.* The result is clearly true for  $n = 2$ , as  $\det(\mathcal{L}) = 1 - q^2 < 0$  for  $|q| > 1$ . So, let us assume the result to be true for  $n = k$ . We now prove the result for  $n = k + 1$ . As before, let  $\bar{T}$  be a tree on  $k + 1$  vertices. Let  $k + 1$  be a pendant vertex adjacent to vertex  $k$ . Then in the block form,  $\bar{\mathcal{L}}$  is given by  $\bar{\mathcal{L}} = \begin{bmatrix} q^2 \mathbf{e}_k \mathbf{e}_k^t + \mathcal{L} & -q \mathbf{e}_k \\ -q \mathbf{e}_k^t & 1 \end{bmatrix}$ . Let  $Q = \begin{bmatrix} I & q \mathbf{e}_k \\ \mathbf{0}^t & 1 \end{bmatrix}$ . Then it is easy to verify that  $Q \bar{\mathcal{L}} Q^t = \begin{bmatrix} \mathcal{L} & \mathbf{0} \\ \mathbf{0}^t & 1 \end{bmatrix} \equiv B(\text{say})$ . Then by Sylvester's inertia theorem, the matrices  $\bar{\mathcal{L}}$  and  $B$  have the same inertia. Therefore, the conclusion follows by appealing to the induction hypothesis.  $\blacksquare$

We now relate the two matrices  $\mathcal{L}$  and  $F$ . By definition,

$$F^{-1} = I - \frac{q}{1 - q^2} A + \frac{q^2}{1 - q^2} \text{diag}(\mathbf{d}) = \frac{1}{1 - q^2} ((1 - q^2)I - qA + q^2 \text{diag}(\mathbf{d})). \quad (3.24)$$

Also,

$$\begin{aligned} \mathcal{L} &= qL - (q - 1)I + q(q - 1)\text{diag}(\mathbf{z}) \\ &= q(\text{diag}(\mathbf{d}) - A) - (q - 1)I + q(q - 1)\text{diag}(\mathbf{d} - \mathbf{e}) \\ &= (1 - q^2)I - qA + q^2 \text{diag}(\mathbf{d}) \end{aligned} \quad (3.25)$$

Thus, from (3.25) and (3.24), we see that  $(1 - q^2)F^{-1} = \mathcal{L}$ . Hence, we arrive at the following lemma.

**LEMMA 3.8** *Let  $T$  be a tree on  $n$  vertices and let  $F$  be the corresponding exponential matrix. If  $\mathcal{L}$  is the  $q$ -Laplacian matrix and  $q \neq \pm 1$ , then*

$$(1 - q^2)F^{-1} = \mathcal{L}.$$

Using the above lemma, we get the following corollary to Proposition 3.4.

**COROLLARY 3.9** *Let  $T$  be a tree on  $n$  vertices and let  $F$  be the corresponding exponential matrix. Then  $F$  is a positive definite matrix for  $q \in (-1, 1)$ .*

*Proof.* Note that a matrix  $A$  is positive definite if and only if  $A^{-1}$  is positive definite. By Proposition 3.4, we know that  $\mathcal{L}$  is positive definite for all  $q \in (-1, 1)$ . Also,  $1 - q^2 > 0$  for all  $q \in (-1, 1)$ . So, by Lemma 3.8,  $F^{-1}$  is a positive definite matrix and hence  $F$  itself is a positive definite matrix. ■

## 4 Invariant factors of the $q$ -distance matrix

We first prove a preliminary result.

**LEMMA 4.10** *Let  $T$  be a tree on  $n \geq 3$  vertices. Then one of the following holds:*

1.  $T$  has a pendant vertex adjacent to a vertex of degree 2.
2.  $T$  has 2 pendant vertices adjacent to the same vertex.

*Proof.* Let  $P = [u_1, u_2, \dots, u_{k-2}, u_{k-1}, u_k]$  be a path corresponding to the diameter of  $T$ . Note that as  $n \geq 3$ ,  $k \geq 3$ . If  $\deg_T(u_{k-1}) = 2$  then the first condition holds.

If  $\deg_T(u_{k-1}) > 2$ , let  $v$  be another vertex adjacent to  $u_{k-1}$ , other than  $u_{k-2}$  and  $u_k$ . Since the diameter of  $T$  is the same as the length of  $P$ , it follows that  $\deg_T(v) = 1$ . Thus Case 2 holds. ■

Recall that a square matrix  $A$  with polynomial entries over  $\mathbb{R}$  is called *unimodular* if  $\det(A)$  is a nonzero real number. For our purpose, we use the word “unimodular” to describe a matrix which satisfies the stronger condition that its determinant is  $\pm 1$ .

**THEOREM 4.11** *Let  $T$  be a tree on  $n \geq 3$  vertices and  $\mathcal{D}$  be the  $q$ -distance matrix of  $T$ . Also, let  $n$  be a pendant vertex. Then there exists a unimodular matrix  $U_n$  such that*

$$U_n \mathcal{D} U_n^t = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \bigoplus_{i=1}^{n-2} \left[ -\frac{i+1}{i} (1+q) \right]$$

and  $U_n \mathbf{e}_n = \mathbf{e}_n$ .

*Proof.* We will prove the result by induction on  $n$ . For  $n = 3$ ,  $\mathcal{D} = \begin{bmatrix} 0 & 1 & 1+q \\ 1 & 0 & 1 \\ 1+q & 1 & 0 \end{bmatrix}$ .

Let  $P_n = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -(1+q) & 1 \end{bmatrix}$ . Then

$$P_n \mathcal{D} P_n^t = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -2(1+q) \end{bmatrix} \quad \text{and} \quad P_n \mathbf{e}_n = \mathbf{e}_n.$$

So, the statement is true for  $n = 3$ . Let the statement be true for  $n = k$  and  $\bar{T}$  be a tree on  $k+1$  vertices with  $k+1$  as a pendant vertex. We will prove the result by considering two cases.

**Case 1:** Suppose that the vertex  $k+1$  is adjacent to the vertex  $k$  of degree 2 (Figure 2).




Figure 2: Figure for Case 1

Let the vertex  $k$  be adjacent to the vertex  $k-1$ . Then the matrix  $\mathcal{D}_{k+1}$  has the form

$$\mathcal{D}_{k+1} = \begin{bmatrix} \mathcal{D}_{k-1} & \mathbf{e} + q\mathcal{D}_{k-1}\mathbf{e}_{k-1} & \mathbf{e} + q\mathbf{e} + q^2\mathcal{D}_{k-1}\mathbf{e}_{k-1} \\ (\mathbf{e} + q\mathcal{D}_{k-1}\mathbf{e}_{k-1})^t & 0 & 1 \\ (\mathbf{e} + q\mathbf{e} + q^2\mathcal{D}_{k-1}\mathbf{e}_{k-1})^t & 1 & 0 \end{bmatrix},$$

where  $\mathcal{D}_{k-1}$  is the polynomial matrix corresponding to the tree  $\bar{T} \setminus \{k, k+1\}$ . Let  $E_{ij} = \mathbf{e}_i \mathbf{e}_j^t$  and define  $P_1 = I - (1+q)E_{k+1,k}$ . Then

$$P_1 \mathcal{D}_{k+1} P_1^t = \begin{bmatrix} \mathcal{D}_{k-1} & \mathbf{e} + q\mathcal{D}_{k-1}\mathbf{e}_{k-1} & -q\mathcal{D}_{k-1}\mathbf{e}_{k-1} \\ (\mathbf{e} + q\mathcal{D}_{k-1}\mathbf{e}_{k-1})^t & 0 & 1 \\ (-q\mathcal{D}_{k-1}\mathbf{e}_{k-1})^t & 1 & -2(1+q) \end{bmatrix}.$$

Now taking  $P_2 = I + qE_{k+1,k-1}$ , we get

$$P_2 P_1 \mathcal{D}_{k+1} P_1^t P_2^t = \begin{bmatrix} \mathcal{D}_{k-1} & \mathbf{e} + q\mathcal{D}_{k-1}\mathbf{e}_{k-1} & \mathbf{0} \\ (\mathbf{e} + q\mathcal{D}_{k-1}\mathbf{e}_{k-1})^t & 0 & 1+q \\ \mathbf{0} & 1+q & -2(1+q) \end{bmatrix}.$$

Note that the upper left  $2 \times 2$  block matrix is nothing but the  $q$ -distance matrix  $\mathcal{D}_k$  of the tree  $\bar{T} \setminus \{k+1\}$ . Observe that for this tree, the vertex  $k$  is a pendant vertex. So, by the induction hypothesis, there exists a unimodular matrix  $U_1$  such that

$$U_1 \mathcal{D}_k U_1^t = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \bigoplus_{i=1}^{k-2} \left[ -\frac{i+1}{i}(1+q) \right] \quad \text{and} \quad U_1 \mathbf{e}_k = \mathbf{e}_k.$$

Thus,

$$\begin{aligned} \begin{bmatrix} U_1 & \mathbf{0} \\ \mathbf{0} & 1 \end{bmatrix} P_2 P_1 \mathcal{D}_{k+1} P_1^t P_2^t \begin{bmatrix} U_1^t & \mathbf{0} \\ \mathbf{0} & 1 \end{bmatrix} &= \begin{bmatrix} U_1 \mathcal{D}_k U_1^t & (1+q)U_1 \mathbf{e}_k \\ (1+q)\mathbf{e}_k^t U_1^t & -2(1+q) \end{bmatrix} \\ &= \left[ \begin{array}{cccccc|c} 0 & 1 & 0 & & 0 & 0 \\ 1 & 0 & 0 & & 0 & 0 \\ 0 & 0 & -2(1+q) & \cdots & 0 & 0 \\ \vdots & \vdots & & \ddots & & \vdots \\ 0 & 0 & 0 & \cdots & -\frac{k-1}{k-2}(1+q) & 1+q \\ \hline 0 & 0 & 0 & \cdots & 1+q & -2(1+q) \end{array} \right]. \end{aligned}$$

So, taking  $P_3 = I + \frac{k-2}{k-1} E_{k+1,k}$ , we have

$$P_3 \begin{bmatrix} U_1 & \mathbf{0} \\ \mathbf{0} & 1 \end{bmatrix} P_2 P_1 \mathcal{D}_{k+1} P_1^t P_2^t \begin{bmatrix} U_1^t & \mathbf{0} \\ \mathbf{0} & 1 \end{bmatrix} P_3^t = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \bigoplus_{i=1}^{k-1} \left[ -\frac{i+1}{i}(1+q) \right].$$

It can be easily verified that

$$\det \left( P_3 \begin{bmatrix} U_1 & \mathbf{0} \\ \mathbf{0} & 1 \end{bmatrix} P_2 P_1 \right) = 1 \quad \text{and} \quad P_3 \begin{bmatrix} U_1 & \mathbf{0} \\ \mathbf{0} & 1 \end{bmatrix} P_2 P_1 \mathbf{e}_{k+1} = \mathbf{e}_{k+1}.$$

**Case 2:** Suppose that the vertices  $k+1$  and  $k$  are both pendant and are adjacent to the vertex  $k-1$  (see Figure 3).




Figure 3: Figure for Case 2

In this case, the matrix  $\mathcal{D}_{k+1}$  has the form

$$\mathcal{D}_{k+1} = \begin{bmatrix} \mathcal{D}_{k-1} & \mathbf{e} + q\mathcal{D}_{k-1}\mathbf{e}_{k-1} & \mathbf{e} + q\mathcal{D}_{k-1}\mathbf{e}_{k-1} \\ (\mathbf{e} + q\mathcal{D}_{k-1}\mathbf{e}_{k-1})^t & 0 & 1+q \\ (\mathbf{e} + q\mathcal{D}_{k-1}\mathbf{e}_{k-1})^t & 1+q & 0 \end{bmatrix}.$$

Let us take  $P_1 = I - E_{k+1,k}$ . Then

$$P_1 \mathcal{D}_{k+1} P_1^t = \begin{bmatrix} \mathcal{D}_{k-1} & \mathbf{e} + q\mathcal{D}_{k-1}\mathbf{e}_{k-1} & \mathbf{0} \\ (\mathbf{e} + q\mathcal{D}_{k-1}\mathbf{e}_{k-1})^t & 0 & 1+q \\ \mathbf{0} & 1+q & -2(1+q) \end{bmatrix}.$$

Note again that the upper left  $2 \times 2$  block matrix is nothing but the  $q$ -distance matrix  $\mathcal{D}_k$ . Observe again that for this tree, the vertex  $k$  is a pendant vertex. So, by the induction hypothesis, there exists a unimodular matrix  $U_1$  such that

$$U_1 \mathcal{D}_k U_1^t = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \bigoplus_{i=1}^{k-2} \left[ -\frac{i+1}{i}(1+q) \right] \quad \text{and} \quad U_1 \mathbf{e}_k = \mathbf{e}_k.$$

So, taking  $P_3 = I + \frac{k-2}{k-1} E_{k+1,k}$ , we have

$$P_3 \begin{bmatrix} U_1 & \mathbf{0} \\ \mathbf{0} & 1 \end{bmatrix} P_1 \mathcal{D}_{k+1} P_1^t \begin{bmatrix} U_1^t & \mathbf{0} \\ \mathbf{0} & 1 \end{bmatrix} P_3^t = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \bigoplus_{i=1}^{k-1} \left[ -\frac{i+1}{i}(1+q) \right].$$

It can be easily verified that

$$\det \left( P_3 \begin{bmatrix} U_1 & \mathbf{0} \\ \mathbf{0} & 1 \end{bmatrix} P_1 \right) = 1 \quad \text{and} \quad P_3 \begin{bmatrix} U_1 & \mathbf{0} \\ \mathbf{0} & 1 \end{bmatrix} P_1 \mathbf{e}_{k+1} = \mathbf{e}_{k+1}.$$

Hence, by the induction hypothesis, the statement holds for all  $n \geq 3$ .  $\blacksquare$

As a corollary to Theorem 4.11, we get the following result about the *inertia* of the matrix  $\mathcal{D}$ . Recall that inertia of a Hermitian matrix  $A$  is defined as the triplet  $(p, n, z)$ , where  $p, n, z$  are the number of positive, negative and zero eigenvalues of  $A$ , respectively.

**COROLLARY 4.12** *Let  $T$  be a tree on  $n$  vertices,  $n \geq 3$ . Also, let  $\mathcal{D}$  be the corresponding  $q$ -distance matrix. Then the following hold:*

1. *if  $q > -1$ , then the inertia of  $\mathcal{D}$  is  $(1, n-1, 0)$ .*
2. *if  $q < -1$ , then the inertia of  $\mathcal{D}$  is  $(n-1, 1, 0)$ .*
3. *if  $q = -1$ , then the inertia of  $\mathcal{D}$  is  $(1, 1, n-2)$ .*

*Proof.* Since the matrices  $\mathcal{D}$  and  $U\mathcal{D}U^t$  are congruent, the result follows from *Sylvester's law of inertia*.  $\blacksquare$

It may be remarked that when  $q > -1$ ,  $\mathcal{D}$  is an elliptic matrix with a zero diagonal in the sense of Fiedler [3]. Also, for  $q = 1$ , the  $q$ -distance matrix is the distance matrix, and one gets the well known result (see Theorem 3, [5]) that the distance matrix has exactly one positive eigenvalue and  $n-1$  negative eigenvalues.

As another application of Theorem 4.11, we obtain the Smith normal form of the  $q$ -distance matrix.

**COROLLARY 4.13** *Let  $T$  be a tree on  $n \geq 3$  vertices and  $\mathcal{D}$  be the  $q$ -distance matrix of  $T$ . Then there exist unimodular matrices  $U, V$  such that*

$$U\mathcal{D}V = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \bigoplus_{i=1}^{n-2} \left[ -\frac{i+1}{i}(1+q) \right],$$

*a diagonal matrix.*

*Proof.* Let  $n$  be a pendant vertex of  $T$ . By Theorem 4.11 there exists a unimodular matrix  $U_n$  such that  $U_n\mathcal{D}U_n^t = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \bigoplus_{i=1}^{n-2} \left[ -\frac{i+1}{i}(1+q) \right]$ . Note that the matrix  $U_n\mathcal{D}^tU_n$  is not a diagonal matrix. This matrix differs from the diagonal matrix only in the first block. Therefore, if we take  $U = \left( \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \oplus I \right) U_n$  and  $V = U_n^t$ , then the new matrix

$$U\mathcal{D}V = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \bigoplus_{i=1}^{n-2} \left[ -\frac{i+1}{i}(1+q) \right]$$

is a diagonal matrix. Also, the matrices  $U$  and  $V$  are unimodular as the matrix  $U_n$  was unimodular. ■

**REMARK 2** *Observe that the matrix  $U_n\mathcal{D}U_n^t$  in Theorem 4.11 is not a diagonal matrix, whereas the matrix  $UDV$  in Corollary 4.13 is a diagonal matrix.*

## 5 $q$ -distance matrix of a weighted tree

We now define the  $q$ -distance matrix of a weighted tree  $T$  on  $n$  vertices. Let  $D = (d_{ij})$  be its distance matrix. Suppose the weights on the  $n-1$  edges of the tree  $T$  are any real numbers  $w_1, w_2, \dots, w_{n-1}$ . Let  $i = i_0, (i_0, i_1), i_1, (i_1, i_2), i_2, \dots, i_{k-1}, (i_{k-1}, i_k), i_k = j$  be a path of length  $k$  from a vertex  $i$  to a vertex  $j$  of  $T$ . If the edge  $(i_t, i_{t+1})$  has weight  $w_t$ , then the  $(i, j)^{\text{th}}$  entry of the  $q$ -distance matrix  $\mathcal{D}$  is set to be  $w_0 + qw_1 + q^2w_2 + \dots + w_{k-1}q^{k-1}$ . Note that the diagonal entries of the matrix  $\mathcal{D}$  are zero and  $\mathcal{D}$  is not a symmetric matrix in general. Also, let  $\sigma_n = \sum_{i=1}^n w_i$ .




Figure 4: A Weighted Tree on 6 Vertices

For example, the distance matrix  $\mathcal{D}$ , for the tree  $T$  shown in Figure 5 is given by

$$\begin{bmatrix} 0 & w_1 & w_1 + qw_2 & w_1 + qw_2 + w_3q^2 & w_1 + qw_2 + w_4q^2 & w_1 + w_5q \\ w_1 & 0 & w_2 & w_2 + w_3q & w_2 + w_4q & w_5 \\ w_2 + qw_1 & w_2 & 0 & w_3 & w_4 & w_2 + w_5q \\ w_3 + qw_2 + q^2w_1 & w_3 + qw_2 & w_3 & 0 & w_3 + w_4q & w_3 + qw_2 + w_5q^2 \\ w_4 + qw_2 + q^2w_1 & w_4 + qw_2 & w_4 & w_4 + w_3q & 0 & w_4 + qw_2 + w_5q^2 \\ w_5 + qw_1 & w_5 & w_5 + qw_2 & w_5 + qw_2 + w_3q^2 & w_5 + qw_2 + w_4q^2 & 0 \end{bmatrix}$$

In the next result we obtain a formula for the determinant of  $\mathcal{D}$ .

**THEOREM 5.1** *Let  $T$  be a weighted tree on  $n$  vertices with edge weights  $w_1, w_2, \dots, w_{n-1}$ . If  $q \neq -1$ , then*

$$\det(\mathcal{D}) = (-1)^{n-1}(1+q)^{n-2}\sigma_n \prod_{i=1}^n w_i.$$

*Proof.* We will prove the result by induction on  $n$ . For  $n = 2$ , we have  $\mathcal{D} = \begin{bmatrix} 0 & w_1 \\ w_1 & 0 \end{bmatrix}$ .

So,  $\det(\mathcal{D}) = -w_1^2 = (-1)^1\sigma_1w_1$ . That is, the result is true for  $n = 2$ .

Let the result be true for  $n = k$ . We now prove the result for  $n = k + 1$ . Suppose  $\bar{T}$  is a tree with edge weights  $w_1, w_2, \dots, w_k$ . Suppose further that  $\bar{T}$  has a pendant vertex  $k + 1$  and is adjacent to the vertex  $k$  with edge weight  $w_k$ . We assume that  $q \neq -1$  and that  $\sum_{i=1}^{k-1} w_i \neq 0$ . This results in no loss of generality since the restrictions can be removed by a continuity argument. Then

$$\bar{\mathcal{D}} = \begin{bmatrix} \mathcal{D} & \mathcal{D}\mathbf{e}_k + w_k\mathbf{q} \\ w_k\mathbf{e}^t + q\mathbf{e}_k^t\mathcal{D} & 0 \end{bmatrix},$$

where for any  $q \in \mathbb{R}$ ,  $\mathbf{q}$  is defined in (3.17).

The proof of the induction part will be done in four steps.

**Step 1:**  $(\mathbf{e} - q\mathbf{z})^t\mathbf{q} = \mathbf{e}^t\mathbf{q} - q\mathbf{z}^t\mathbf{q} = 1 + q$ .

To prove this, suppose that there is a vertex  $i_0$  adjacent to  $t$  vertices, say,  $i_1, i_2, \dots, i_t$ . Also suppose  $i_0$  is at a distance  $d$  from the vertex  $k + 1$ . Then in the expression  $\mathbf{e}^t\mathbf{q}$ , the contribution due to the presence of  $t$  vertices being adjacent to  $i_0$  is  $q^{d-2} + q^{d-1} + (t-1)q^d$ . But in the expression  $q\mathbf{z}^t\mathbf{q}$ , the information that the degree of the vertex  $i_0$  is  $t$ , gives  $q \cdot (t-1)q^{d-1} = (t-1)q^d$ . Thus, in  $(\mathbf{e} - q\mathbf{z})^t\mathbf{q}$ , the contribution at vertex  $i_0$  is  $q^{d-2} + q^{d-1}$ . That is, there is no contribution from the vertices that are at a distance  $d + 1$  from the vertex  $k + 1$ . But then this will be true for all vertices that are at a distance 1 or more. Hence, the only term left out in the expression  $(\mathbf{e} - q\mathbf{z})^t\mathbf{q}$ , is  $1 + q$ .

**Step 2:** In this step, we show that  $\mathbf{e}^t\mathcal{D}^{-1} = \frac{1}{\sigma_{k-1}}(\mathbf{e}^t - q\mathbf{z}^t)$ . That is, we show that  $\sigma_{k-1}\mathbf{e}^t = (\mathbf{e}^t - q\mathbf{z}^t)\mathcal{D}$ .

The result will also be proved by induction. The initial step in the induction argument

can be easily verified. Let the result be true for all trees with  $k$  vertices. We now prove the result for a tree with  $k + 1$  vertices. From (2.1), note that

$$\mathbf{e}^t - q\bar{\mathbf{z}}^t = [\mathbf{e}^t \mid 1] - q[(\mathbf{z} + \mathbf{e}_k)^t \mid 0] = [\mathbf{e}^t - q(\mathbf{z} + \mathbf{e}_k)^t \mid 1].$$

Now, using step 1 and the induction hypothesis, we get

$$\begin{aligned} (\mathbf{e}^t - q\bar{\mathbf{z}}^t)\bar{\mathcal{D}} &= (\mathbf{e}^t - q(\mathbf{z} + \mathbf{e}_k)^t) \begin{bmatrix} \mathcal{D} & \mathcal{D}\mathbf{e}_k + w_k\mathbf{q} \\ w_k\mathbf{e}^t + q\mathbf{e}_k^t\mathcal{D} & 0 \end{bmatrix} \\ &= [\mathbf{e}^t - q(\mathbf{z} + \mathbf{e}_k)^t \mid 1] \begin{bmatrix} \mathcal{D} & \mathcal{D}\mathbf{e}_k + w_k\mathbf{q} \\ w_k\mathbf{e}^t + q\mathbf{e}_k^t\mathcal{D} & 0 \end{bmatrix} \\ &= [\sigma_k\mathbf{e}^t \mid \sigma_k] = \sigma_k\mathbf{e}^t. \end{aligned} \quad (5.1)$$

Thus, by the induction hypothesis, the proof of step 2 is complete.

**Step 3:** We now show that  $(w_k\mathbf{e}^t + q\mathbf{e}_k^t\mathcal{D})\mathcal{D}^{-1}(\mathcal{D}\mathbf{e}_k + w_k\mathbf{q}) = \frac{(1+q)w_k\sigma_k}{\sigma_{k-1}}$ .

We use the results obtained in step 1 and step 2, to prove this step. We have

$$\begin{aligned} & (w_k\mathbf{e}^t + q\mathbf{e}_k^t\mathcal{D})\mathcal{D}^{-1}(\mathcal{D}\mathbf{e}_k + w_k\mathbf{q}) \\ &= w_k\mathbf{e}^t\mathbf{e}_k + w_k^2\mathbf{e}^t\mathcal{D}^{-1}\mathbf{q} + q\mathbf{e}_k^t\mathcal{D}\mathbf{e}_k + qw_k\mathbf{e}_k^t\mathbf{q} \\ &= w_k + w_k^2\mathbf{e}^t\mathcal{D}^{-1}\mathbf{q} + q \cdot 0 + qw_k = w_k(1+q) + \frac{w_k^2}{\sigma_{k-1}}(\mathbf{e}^t - q\bar{\mathbf{z}}^t)\mathbf{q} \\ &= w_k(1+q) + (1+q)\frac{w_k^2}{\sigma_{k-1}} = \frac{(1+q)w_k\sigma_k}{\sigma_{k-1}}. \end{aligned} \quad (5.2)$$

**Step 4:** We now use (5.2) to complete the induction step. By the induction hypothesis and (5.2), we have

$$\begin{aligned} \det(\bar{\mathcal{D}}) &= \det(\mathcal{D}) \cdot (0 - (w_k\mathbf{e}^t + q\mathbf{e}_k^t\mathcal{D})\mathcal{D}^{-1}(\mathcal{D}\mathbf{e}_k + w_k\mathbf{q})) \\ &= -\det(\mathcal{D})\frac{w_k\sigma_k(1+q)}{\sigma_{k-1}} \\ &= -(-1)^{k-2}\sigma_{k-1} \prod_{i=1}^{k-1} w_i(1+q)^{k-2} \times \frac{w_k\sigma_k(1+q)}{\sigma_{k-1}} \\ &= (-1)^{k-1}(1+q)^{k-1}\sigma_k \prod_{i=1}^k w_i. \end{aligned}$$

Thus, by induction, the proof is over. ■

It is tempting to obtain a formula for  $\mathcal{D}^{-1}$ , in the case of a weighted tree. However, it appears that such a formula will be very complicated and we leave it as an open problem. As a consequence of Theorem 5.1, we derive the determinant formula for an unweighted tree.

**COROLLARY 5.2** *Let  $T$  be a tree on  $n$  vertices and let  $\mathcal{D}$  be its  $q$ -distance matrix. Then  $\det(\mathcal{D}) = (-1)^{n-1}(n-1)(1+q)^{n-2}$ .*



*Proof.* In this case, the weight of each edge is 1. So,  $\sigma_{n-1} = 1 + 1 + \cdots + 1 = n - 1$ . Hence, the result follows. ■

For  $q = 1$ , the above result reduces to the result of Graham and Pollak [5] on  $\det(D)$ .

**COROLLARY 5.3** *Let  $T$  be a tree on  $n$  vertices and let  $D$  be its distance matrix. Then  $\det(D) = (-1)^{n-1}(n-1)2^{n-2}$ .*

**ACKNOWLEDGMENT:** We sincerely thank the referee for many helpful suggestions.

## References

- [1] R. B. Bapat, S. J. Kirkland, M. Neumann, *On distance matrices and Laplacians*, *Linear Algebra Appl.*, 401 : 193 - 209, 2005.
- [2] R. B. Bapat and T. E. S. Raghavan, "Nonnegative Matrices and Applications," *Encyclopedia of Mathematics and its Applications* 64, ed. G. C. Rota, Cambridge University Press, Cambridge, London, 1997.
- [3] Miroslav Fiedler, *Some inverse problems for elliptic matrices with zero diagonal*, *Linear Algebra Appl.*, 332-334 : 197 - 204, 2001.
- [4] R. L. Graham and L. Lovasz, *Distance Matrix Polynomials of Trees*, *Adv. in Math.*, 29 (1): 60 - 88, 1978.
- [5] R. L. Graham and H. O. Pollak, *On the addressing problem for loop switching*, *Bell. System Tech. J.*, 50 (1971), 2495 - 2519.
- [6] F. Harary, "Graph Theory," Addison-Wesley, New York.
- [7] L. Hogben, *Spectral Graph Theory and the Inverse Eigenvalue Problem of a Graph*, *Electronic Journal of Linear Algebra*, vol. 14, 12 - 31, January 2005.
- [8] R. A. Horn and C. R. Johnson, "Matrix Analysis," Cambridge University Press, Cambridge, London, 1985.