

AN EXTENSION OF A THEOREM OF EULER

NORIKO HIRATA-KOHNO, SHANTA LAISHRAM, T. N. SHOREY, AND R. TIJDEMAN

ABSTRACT. It is proved that equation (1) with $4 \leq k \leq 109$ does not hold. The paper contains analogous result for $k \leq 100$ for more general equation (2) under certain restrictions.

1. INTRODUCTION

The theorem of Euler ([Eul80], cf. [Mor69, p.21-22], [MS03]) referred in the title of this paper is that a product of four terms in arithmetic progression is never a square. Let $n, d, k \geq 2$ and y be positive integers such that $\gcd(n, d) = 1$. We consider the equation

$$(1) \quad n(n+d) \cdots (n+(k-1)d) = y^2$$

in n, d, k and y . It has infinitely many solutions when $k = 2$ or 3 . A well-known conjecture states that (1) with $k \geq 4$ is not possible. We claim

Theorem 1. *Equation (1) with $4 \leq k \leq 109$ is not possible.*

By Euler, Theorem 1 is valid when $k = 4$. The case when $k = 5$ is due to Obláth [Obl50]. Independently of the authors, Bennett, Bruin, Győry and Hajdu [BBGH06] proved that (1) with $6 \leq k \leq 11$ does not hold. Theorem 1 has been confirmed by Erdős [Erd39] and Rigge [Rig39], independently of each other, when $d = 1$.

Theorem 1 is derived from a more general result and we introduce some notation for stating this. For an integer $\nu > 1$, we denote by $P(\nu)$ the greatest prime factor of ν and we put $P(1) = 1$. Let b be a squarefree positive integer such that $P(b) \leq k$. We consider a more general equation than (1), namely

$$(2) \quad n(n+d) \cdots (n+(k-1)d) = by^2.$$

We write

$$(3) \quad n + id = a_i x_i^2 \quad \text{for } 0 \leq i < k$$

where a_i are squarefree integers such that $P(a_i) \leq \max(P(b), k-1)$ and x_i are positive integers. Every solution to (2) yields a k -tuple $(a_0, a_1, \dots, a_{k-1})$. We re-write (2) as

$$(4) \quad m(m-d) \cdots (m-(k-1)d) = by^2, \quad m = n + (k-1)d.$$

The equation (4) is called the mirror image of (2). The corresponding k -tuple $(a_{k-1}, a_{k-2}, \dots, a_0)$ is called the mirror image of $(a_0, a_1, \dots, a_{k-1})$.

Let $P(b) < k$. Erdős and Selfridge [ES75] proved that (2) with $d = 1$ never holds under the assumption that the left-hand side of (2) is divisible by a prime greater than or equal to k . The result does not hold unconditionally. As mentioned above, equation (2) with $k = 2, 3$ and $b = 1$ has infinitely many solutions. This is also the case when $k = 4$ and $b = 6$, see Tijdeman [Tij89]. On the other hand, equation (2) with $k = 4$ and $b \neq 6$ does not hold. We consider (2) with $d > 1$ and $k \geq 5$. We prove

Theorem 2. *Equation (2) with $d > 1, P(b) < k$ and $5 \leq k \leq 100$ implies that $(a_0, a_1, \dots, a_{k-1})$ is among the following tuples or their mirror images.*

$$(5) \quad \begin{aligned} k = 8 &: (2, 3, 1, 5, 6, 7, 2, 1), (3, 1, 5, 6, 7, 2, 1, 10); \\ k = 9 &: (2, 3, 1, 5, 6, 7, 2, 1, 10); \\ k = 14 &: (3, 1, 5, 6, 7, 2, 1, 10, 11, 3, 13, 14, 15, 1); \\ k = 24 &: (5, 6, 7, 2, 1, 10, 11, 3, 13, 14, 15, 1, 17, 2, 19, 5, 21, 22, 23, 6, 1, 26, 3, 7). \end{aligned}$$

Theorem 2 with $k = 5$ is due to Mukhopadhyay and Shorey [MS03]. Initially, Bennett, Bruin, Győry, Hajdu [BBGH06] and Hirata-Kohno, Shorey (unpublished), independently, proved Theorem 2 with $k = 6$ and $(a_0, a_1, \dots, a_5) \neq (1, 2, 3, 1, 5, 6), (6, 5, 1, 3, 2, 1)$. Next Bennett, Bruin, Győry and Hajdu [BBGH06] removed the assumption on (a_0, a_1, \dots, a_5) in the above result. Thus (2) with $k = 6$ does not hold and we shall refer to it as *the case $k = 6$* . Bennett, Bruin, Győry and Hajdu [BBGH06], independently of us, showed that (2) with $7 \leq k \leq 11$ and $P(b) \leq 5$ is not possible. This is now a special case of Theorem 2.

Let $P(b) = k$. Then we have no new result on (2) with $k = 5$. For $k \geq 7$, we prove

Theorem 3. *Equation (2) with $d > 1, P(b) = k$ and $7 \leq k \leq 100$ implies that $(a_0, a_1, \dots, a_{k-1})$ is among the following tuples or their mirror images.*

$$(6) \quad \begin{aligned} k = 7 &: (2, 3, 1, 5, 6, 7, 2), (3, 1, 5, 6, 7, 2, 1), (1, 5, 6, 7, 2, 1, 10); \\ k = 13 &: (3, 1, 5, 6, 7, 2, 1, 10, 11, 3, 13, 14, 15), \\ & \quad (1, 5, 6, 7, 2, 1, 10, 11, 3, 13, 14, 15, 1); \\ k = 19 &: (1, 5, 6, 7, 2, 1, 10, 11, 3, 13, 14, 15, 1, 17, 2, 19, 5, 21, 22); \\ k = 23 &: (5, 6, 7, 2, 1, 10, 11, 3, 13, 14, 15, 1, 17, 2, 19, 5, 21, 22, 23, 6, 1, 26, 3), \\ & \quad (6, 7, 2, 1, 10, 11, 3, 13, 14, 15, 1, 17, 2, 19, 5, 21, 22, 23, 6, 1, 26, 3, 7). \end{aligned}$$

It has been conjectured that (2) with $k \geq 5$ never holds. Granville (unpublished) showed that k is bounded by an absolute constant whenever *abc*-conjecture holds, see Laishram [Lai04] for a proof. For the convenience of the proofs, we consider Theorems 2 and 3 together. Therefore we formulate

Theorem 4. *Let $d > 1, P(b) \leq k$ and $5 \leq k \leq 100$. Suppose that $k \neq 5$ if $P(b) = k$. Then (2) does not hold except for the $(a_0, a_1, \dots, a_{k-1})$ among (5), (6) and their mirror images.*

It is clear that Theorem 4 implies Theorems 2 and 3. In fact the proof of Theorem 4 provides a method for solving (2) for any given value of k unless $(a_0, a_1, \dots, a_{k-1})$

is given by (5), (6) and their mirror images. This is a new and useful feature of the paper. We have restricted k up to 100 for keeping the computational load under control. It is an open problem to solve (2) for an infinite sequence of values of k . A solution to this problem may be an important contribution towards the Conjecture stated just after Theorem 3. Theorem 4 has been applied in [LS] to show that (2) with $k \geq 6$ implies that $d > 10^{10}$. For more applications, see [LS].

Now we give a sketch of the proof of Theorem 4. Let the assumptions of Theorem 4 be satisfied. Assume (2) such that $(a_0, a_1, \dots, a_{k-1})$ is not among (5), (6) or their mirror images. As already stated, the cases $k = 5$ and $k = 6$ have already been solved in [MS03] and [BBGH06]. Therefore we suppose that $k \geq 7$. Further it suffices to assume that k is prime and we proceed inductively on k . Let k be given. Then we choose a suitable pair (q_1, q_2) of distinct primes $\leq k$ such that

$$\left(\frac{p}{q_1}\right) = \left(\frac{p}{q_2}\right)$$

for small primes p . For example, when $k = 29$, we take $(q_1, q_2) = (19, 29)$ so that the above relation holds with $p = 2, 3, 5, 7$. We show that $q_1 \nmid d$ and $q_2 \nmid d$, see Lemma 8. Assume $q_1 \mid d$ or $q_2 \mid d$. Then we find two primes Q_1 and Q_2 such that $Q_1 \mid d$ or $Q_2 \mid d$ whenever $k \geq 29$, see Lemma 7. Now we arrive at a contradiction by a counting argument using (9) and Lemmas 1, 2. Hence $q_1 \nmid d$ and $q_2 \nmid d$ but this is excluded by Lemma 6, the proof of which depends on Lemma 5. In fact, we need to apply it repeatedly for $k > 11$.

In the case $k = 6$, Bennett, Bruin, Györy and Hajdu [BBGH06] solved the cases $(a_0, a_1, \dots, a_5) \in \{(1, 2, 3, 1, 5, 6), (6, 5, 1, 3, 2, 1)\}$ by using explicit Chabauty techniques due to Bruin and Flynn [BF05]. These cases appear to be similar to our exceptional cases (5) and (6) where we have, in fact, more freedom in the sense that there are at least 7 curves where we may consider applying Chabauty method. Finally we remark that it suffices to solve the cases $k = 7$ in (6) or its mirror images for Theorems 2 and 3 and hence Theorem 4. Further it suffices to solve the cases $k = 8$ in (5) or its mirror images for Theorem 2.

2. NOTATION AND LEMMAS

We define some notation. Let

$$R = \{a_i : 0 \leq i < k\}$$

and for a prime q , we put

$$(7) \quad S = S(q) = \{a \in R : P(a) \leq q\}, \quad S_1 = S_1(q) = \{a \in R : P(a) > q\}.$$

Further we write

$$(8) \quad T = T(q) = \{i : a_i \in S\}, \quad T_1 = T_1(q) = \{i : a_i \in S_1\}.$$

Then we see that

$$(9) \quad |T| + |T_1| = k.$$

For $a \in R$, let

$$\nu(a) = |\{i : a_i = a\}|, \nu_o(a) = |\{i : a_i = a, 2 \nmid x_i\}|, \nu_e(a) = |\{i : a_i = a, 2 \mid x_i\}|.$$

We observe that

$$(10) \quad |T| = \sum_{a \in S} \nu(a).$$

Let

$$\delta = \min(3, \text{ord}_2(d))$$

and

$$\rho = \begin{cases} 3 & \text{if } 3 \mid d, \\ 1 & \text{otherwise.} \end{cases}$$

We have

Lemma 1. For $a \in R$, let $\mathcal{K}_a = \frac{k}{a2^{3-\delta}}$, $\mathcal{K}'_a = \frac{k}{16a}$,

$$f_1(k, a, \delta) = \begin{cases} 1 & \text{if } k \leq a2^{3-\delta} \\ \lceil \mathcal{K}_a \rceil - \lfloor \frac{\lceil \mathcal{K}_a \rceil}{4} \rfloor & \text{if } k > a2^{3-\delta}, 3 \mid d \\ \sum_{i=1}^2 \left(\lceil \frac{\mathcal{K}_a}{3^i} \rceil - \lfloor \frac{\lceil \mathcal{K}_a \rceil}{4} \rfloor \right) & \text{if } k > a2^{3-\delta}, 3 \nmid d \end{cases}$$

and

$$f_2(k, a) = \begin{cases} 1 & \text{if } k \leq 4a \\ \lceil \mathcal{K}'_a \rceil + 1 & \text{if } 4a < k \leq 32a \\ \sum_{i=1}^2 \left(\lceil \frac{\mathcal{K}'_a}{i} \rceil - \lfloor \frac{\lceil \mathcal{K}'_a \rceil}{4} \rfloor \right) & \text{if } k > 32a, 3 \mid d \\ \sum_{i=1}^2 \left(\lceil \frac{\mathcal{K}'_a}{3^i} \rceil - \lfloor \frac{\lceil \mathcal{K}'_a \rceil}{4} \rfloor \right) + \sum_{i=1}^2 \left(\lceil \frac{\mathcal{K}'_a}{2 \cdot 3^i} \rceil - \lfloor \frac{\lceil \mathcal{K}'_a \rceil}{4} \rfloor \right) & \text{if } k > 32a, 3 \nmid d \end{cases}$$

Then we have

$$\nu_o(a) \leq f_1(k, a, \delta), \quad \nu_e(a) \leq f_2(k, a)$$

and

$$\nu(a) \leq F(k, a, \delta) := \begin{cases} 1 & \text{if } k \leq a \\ f_1(k, a, \delta) & \text{if } k > a \text{ and } d \text{ even} \\ f_1(k, a, 0) + f_2(k, a) & \text{if } k > a \text{ and } d \text{ odd.} \end{cases}$$

Proof. Let $I_1 = \{i : a_i = a, x_i \text{ odd}\}$, $I_2 = \{i : a_i = a, 2 \mid x_i\}$ and $I_3 = \{i : a_i = a, 4 \mid x_i\}$. Further for $l = 1, 2, 3$, let

$$I_{l1} := \{i \in I_l : 3 \nmid x_i\}, \quad I_{l2} := \{i \in I_l : 3 \mid x_i\}.$$

Let $\tau := \tau(l, m)$ be defined by $\frac{\tau}{a} = 2^{3-\delta} \cdot 3\rho^{-1}, 2^{3-\delta} \cdot 9, 32 \cdot 3\rho^{-1}, 32 \cdot 9, 16 \cdot 3\rho^{-1}, 16 \cdot 9$ for $(l, m) = (1, 1), (1, 2), (2, 1), (2, 2), (3, 1), (3, 2)$, respectively. Since $x_i^2 \equiv 1 \pmod{8}$ for $i \in I_1$, $(\frac{x_i}{2})^2 \equiv 1 \pmod{8}$ for $i \in I_2$, $16 \mid x_i^2$ for $i \in I_3$ and $x_i^2 \equiv 1 \pmod{3}$

for $i \in I_{l1}$, $9|x_i^2$ for $i \in I_{l2}$ for $l = 1, 2, 3$, we see from $(i - j)d = a(x_i^2 - x_j^2)$ that $\tau|(i - j)$ for $i, j \in I_{lm}$. Since $a|(i - j)$ whenever $a_i = a_j$, we get $\nu(a) = 1$ for $k \leq a$. Thus we suppose that $k > a$. We have $\nu(a) = \nu_o(a) + \nu_e(a)$. It suffices to show $\nu_o(a) \leq f_1(k, a, \delta)$ and $\nu_e(a) \leq f_2(k, a)$ since $\nu_e(a) = 0$ for d even. We observe that $\nu_o(a) = |I_1|$ and $\nu_e(a) = |I_2| + |I_3|$. Since $a2^{3-\delta}|(i - j)$ whenever $i, j \in I_1$, we get $|I_1| \leq 1$ if $k \leq a2^{3-\delta}$. Thus we suppose $k > a2^{3-\delta}$ for proving $|I_1| \leq f_1(k, a, \delta)$. Further from $4a|(i - j)$ for $i, j \in I_2 \cup I_3$, $32a|(i - j)$ for $i, j \in I_2$ and $16a|(i - j)$ for $i, j \in I_3$, we get $|I_2| + |I_3| \leq f_2(k, a)$ for $k \leq 32a$. Hence we suppose that $k > 32a$ for showing $|I_2| + |I_3| \leq f_2(k, a)$.

Let (l, m) be with $1 \leq l \leq 3, 1 \leq m \leq 2$. Let $i_0 = \min_{i \in I_{lm}} i$, $N = \frac{n+i_0d}{a}$ and $D = \frac{\tau}{a}d$. Then we see that ax_i^2 with $i \in I_{lm}$ come from the squares in the set $\{N, N + D, \dots, N + (\lceil \frac{k-i_0}{\tau} \rceil - 1)D\}$. Dividing this set into consecutive intervals of length 4 and using Euler's result, we see that there are at most $\lceil \frac{k-i_0}{\tau} \rceil - \lfloor \frac{\lceil \frac{k-i_0}{\tau} \rceil}{4} \rfloor \leq \lceil \frac{k}{\tau} \rceil - \lfloor \frac{\lceil \frac{k}{\tau} \rceil}{4} \rfloor$ of them which can be squares. Hence $|I_{lm}| \leq \lceil \frac{k}{\tau} \rceil - \lfloor \frac{\lceil \frac{k}{\tau} \rceil}{4} \rfloor$. Now the assertion follows from $|I_l| = \sum_{m=1}^2 |I_{lm}|$ for $l = 1, 2, 3$ since $|I_{l2}| = 0$ for $3|d$. \square

We observe that there are $\frac{p-1}{2}$ distinct quadratic residues and $\frac{p-1}{2}$ distinct quadratic nonresidue modulo an odd prime p . The next lemma follows easily from this fact.

Lemma 2. *Assume (2) holds. Let k be an odd prime. Suppose that $k \nmid d$. Let*

$$T' = \{i : \left(\frac{a_i}{k}\right) = 1, 0 \leq i < k\}, \quad T'' = \{i : \left(\frac{a_i}{k}\right) = -1, 0 \leq i < k\}.$$

Then

$$|T'| = |T''| = \frac{k-1}{2}.$$

Lemma 3. *Assume that (2) with $P(b) \leq k$ has no solution at $k = k_1$ with k_1 prime. Then (2) with $P(b) \leq k$ has no solution at k with $k_1 \leq k < k_2$ where k_2 is the smallest prime larger than k_1 .*

Proof. Let k_1 and k_2 be consecutive primes such that $k_1 \leq k < k_2$. Assume that (2) does not hold at (n, d, k_1) . Suppose

$$n(n+d) \cdots (n+(k-1)d) = by^2.$$

Using (3), we see that

$$n(n+d) \cdots (n+(k_1-1)d) = b'y'^2$$

with $P(b') \leq k_1$. This is not possible. \square

Let q_1, q_2 be distinct primes and

$$\Lambda_1(q_1, q_2) := \left\{p \leq 97 : \left(\frac{p}{q_1}\right) \neq \left(\frac{p}{q_2}\right)\right\}.$$

We write $\Lambda(q_1, q_2) = \Lambda(q_1, q_2, k) := \{p \in \Lambda_1(q_1, q_2) : p \leq k\}$.

Lemma 4. *We have*

(q_1, q_2)	$\Lambda_1(q_1, q_2)$
(5, 11)	{3, 19, 23, 29, 37, 41, 47, 53, 61, 67, 79, 97}
(7, 17)	{11, 13, 19, 23, 29, 37, 47, 59, 71, 79, 83, 89}
(11, 13)	{5, 17, 29, 31, 37, 43, 47, 59, 61, 67, 71, 79, 89, 97}
(11, 59)	{7, 17, 19, 23, 29, 31, 37, 41, 47, 67, 79, 89, 97}
(11, 61)	{13, 19, 23, 31, 37, 41, 53, 59, 67, 71, 73, 83, 89}
(19, 29)	{11, 13, 17, 43, 47, 53, 59, 61, 67, 71, 73}
(23, 73)	{13, 19, 29, 31, 37, 47, 59, 61, 67, 79, 89, 97}
(23, 97)	{11, 13, 29, 41, 43, 53, 59, 61, 71, 79, 89}
(31, 89)	{7, 11, 17, 19, 41, 53, 59, 73, 79}
(37, 83)	{17, 23, 29, 31, 47, 53, 59, 61, 67, 71, 73}
(41, 79)	{11, 13, 19, 37, 43, 59, 61, 67, 89, 97}
(43, 53)	{7, 23, 29, 31, 37, 41, 67, 79, 83, 89}
(43, 67)	{11, 13, 19, 29, 31, 37, 41, 53, 71, 73, 79, 89, 97}
(53, 67)	{7, 11, 13, 19, 23, 43, 71, 73, 83, 97}
(59, 61)	{7, 13, 17, 29, 47, 53, 71, 73, 79, 83, 97}
(73, 97)	{11, 19, 23, 31, 37, 41, 43, 47, 53, 67, 71}
(79, 89)	{13, 17, 19, 23, 31, 47, 53, 71, 83}

Definition: Let \mathcal{P} be a set of primes and $\mathcal{I} \subseteq [0, k) \cap \mathbb{Z}$. We say that \mathcal{I} is covered by \mathcal{P} if, for every $j \in \mathcal{I}$, there exists $p \in \mathcal{P}$ such that $p|a_j$. Further for $i \in \mathcal{I}$, let

$$(11) \quad \mathbf{i}(\mathcal{P}) = |\{p \in \mathcal{P} : p \text{ divides } a_i\}|.$$

For a prime p with $\gcd(p, d) = 1$, let i_p be the smallest $i \geq 0$ such that $p|n + id$. For $\mathcal{I} \subseteq [0, k) \cap \mathbb{Z}$ and primes p_1, p_2 with $\gcd(p_1 p_2, d) = 1$, we write

$$\mathcal{I}' = \mathcal{I}(p_1, p_2) = \mathcal{I} \setminus \cup_{j=1}^2 \{i_{p_j} + p_j i : 0 \leq i < \lceil \frac{k}{p_j} \rceil\}.$$

Lemma 5. *Let \mathcal{P}_0 be a set of primes. Let p_1, p_2 be primes such that $\gcd(p_1 p_2, d) = 1$. Let $(i_1, i_2) = (i_{p_1}, i_{p_2})$, $\mathcal{I} \subseteq [0, k) \cap \mathbb{Z}$ and $\mathcal{I}' = \mathcal{I}(p_1, p_2)$ be such that $\mathbf{i}(\mathcal{P}_0 \cap \Lambda(p_1, p_2))$ is even for each $i \in \mathcal{I}'$. Define*

$$\mathcal{I}_1 = \{i \in \mathcal{I}' : \left(\frac{i - i_1}{p_1}\right) = \left(\frac{i - i_2}{p_2}\right)\} \text{ and } \mathcal{I}_2 = \{i \in \mathcal{I}' : \left(\frac{i - i_1}{p_1}\right) \neq \left(\frac{i - i_2}{p_2}\right)\}.$$

Let $\mathcal{P} = \Lambda(p_1, p_2) \setminus \mathcal{P}_0$. Let ℓ be the number of terms $n + id$ with $i \in \mathcal{I}'$ divisible by primes in \mathcal{P} . Then either

$$|\mathcal{I}_1| \leq \ell, \mathcal{I}_1 \text{ is covered by } \mathcal{P}, \mathcal{I}_2 = \{i \in \mathcal{I}' : \mathbf{i}(\mathcal{P}) \text{ is even}\}$$

or

$$|\mathcal{I}_2| \leq \ell, \mathcal{I}_2 \text{ is covered by } \mathcal{P}, \mathcal{I}_1 = \{i \in \mathcal{I}' : \mathbf{i}(\mathcal{P}) \text{ is even}\}.$$

We observe that $\ell \leq \sum_{p \in \mathcal{P}} \lceil \frac{k}{p} \rceil$.

Proof. Let $i \in \mathcal{I}'$. Let $\mathcal{U}_0 = \{p : p|a_i\}$, $\mathcal{U}_1 = \{p \in \mathcal{U}_0 : p \notin \Lambda(p_1, p_2)\}$, $\mathcal{U}_2 = \{p \in \mathcal{U}_0 : p \in \mathcal{P}_0 \cap \Lambda(p_1, p_2)\}$ and $\mathcal{U}_3 = \{p \in \mathcal{U}_0 : p \in \mathcal{P}\}$. Then we have from $a_i = \prod_{p \in \mathcal{U}_0} p$ that

$$\left(\frac{a_i}{p_1}\right) = \prod_{p \in \mathcal{U}_1} \left(\frac{p}{p_1}\right) \prod_{p \in \mathcal{U}_2} \left(\frac{p}{p_1}\right) \prod_{p \in \mathcal{U}_3} \left(\frac{p}{p_1}\right) = (-1)^{i(\mathcal{P})+|\mathcal{U}_2|} \prod_{p \in \mathcal{U}_0} \left(\frac{p}{p_2}\right) = (-1)^{i(\mathcal{P})} \left(\frac{a_i}{p_2}\right)$$

since $|\mathcal{U}_2| = i(\mathcal{P}_0 \cap \Lambda(p_1, p_2))$ is even. Therefore

$$(12) \quad \mathcal{L} := \left\{i \in \mathcal{I}' : \left(\frac{a_i}{p_1}\right) \neq \left(\frac{a_i}{p_2}\right)\right\} = \{i \in \mathcal{I}' : i(\mathcal{P}) \text{ is odd}\}.$$

In particular \mathcal{L} is covered by \mathcal{P} and hence

$$(13) \quad |\mathcal{L}| \leq \ell.$$

We see that $\left(\frac{a_i}{p_j}\right) = \left(\frac{n+id}{p_j}\right) = \left(\frac{i-i_j}{p_j}\right) \left(\frac{d}{p_j}\right)$ for $i \in \mathcal{I}'$ and $j = 1, 2$. Therefore $\mathcal{L} = \mathcal{I}_1$ or \mathcal{I}_2 according as $\left(\frac{d}{p_1}\right) \neq \left(\frac{d}{p_2}\right)$ or $\left(\frac{d}{p_1}\right) = \left(\frac{d}{p_2}\right)$, respectively. Now the assertion of the Lemma 5 follows from (12) and (13). \square

Remark: Let \mathcal{P} consist of one prime p . We observe that $p|n + id$ if and only if $p|(i - i_p)$. Then \mathcal{I}_1 or \mathcal{I}_2 is contained in one residue class modulo p and $p \nmid a_i$ for i in the other set.

Corollary 1. *Let $p_1, p_2, i_1, i_2, \mathcal{P}_0, \mathcal{P}, \mathcal{I}, \mathcal{I}', \mathcal{I}_1, \mathcal{I}_2$ and ℓ be as in Lemma 5. Assume that*

$$(14) \quad \ell < \frac{1}{2}|\mathcal{I}'|.$$

Then $|\mathcal{I}_1| \neq |\mathcal{I}_2|$. Let

$$(15) \quad \mathcal{M} = \begin{cases} \mathcal{I}_1 & \text{if } |\mathcal{I}_1| < |\mathcal{I}_2| \\ \mathcal{I}_2 & \text{otherwise} \end{cases}$$

and

$$(16) \quad \mathcal{B} = \begin{cases} \mathcal{I}_2 & \text{if } |\mathcal{I}_1| < |\mathcal{I}_2| \\ \mathcal{I}_1 & \text{otherwise.} \end{cases}$$

Then $|\mathcal{M}| \leq \ell$, \mathcal{M} is covered by \mathcal{P} and $\mathcal{B} = \{i \in \mathcal{I}' | i(\mathcal{P}) \text{ is even}\}$.

Proof. We see from Lemma 5 that $\min(|\mathcal{I}_1|, |\mathcal{I}_2|) \leq \ell$ and from (14) that $\max(|\mathcal{I}_1|, |\mathcal{I}_2|) \geq \frac{1}{2}|\mathcal{I}'| > \ell$. Now the assertion follows from Lemma 5. \square

We say that $(\mathcal{M}, \mathcal{B}, \mathcal{P}, \ell)$ has *Property \mathfrak{H}* if $|\mathcal{M}| \leq \ell$, \mathcal{M} is covered by \mathcal{P} and $i(\mathcal{P})$ is even for $i \in \mathcal{B}$.

Lemma 6. *Let k be a prime with $7 \leq k \leq 97$ and assume (2). For $k \geq 11$, assume that Theorem 4 is valid for all primes k_1 with $7 \leq k_1 < k$. For $11 \leq k \leq 29$, assume that $k \nmid d$ and $k \nmid n + id$ for $0 \leq i < k - k'$ and $k' \leq i < k$ where $k' < k$ are consecutive primes. Let $(q_1, q_2) = (5, 7)$ if $k = 7$; $(5, 11)$ if $k = 11$; $(11, 13)$ if $13 \leq k \leq 23$; $(19, 29)$ if $29 \leq k \leq 59$; $(59, 61)$ if $k = 61$; $(43, 67)$ if $k = 67, 71$; $(23, 73)$*

if $k = 73, 79$; $(37, 83)$ if $k = 83$; $(79, 89)$ if $k = 89$ and $(23, 97)$ if $k = 97$. Then $q_1|d$ or $q_2|d$ unless $(a_0, a_1, \dots, a_{k-1})$ is given by the following or their mirror images.

$$\begin{aligned} k = 7 &: (2, 3, 1, 5, 6, 7, 2), (3, 1, 5, 6, 7, 2, 1), (1, 5, 6, 7, 2, 1, 10); \\ k = 13 &: (3, 1, 5, 6, 7, 2, 1, 10, 11, 3, 13, 14, 15), (1, 5, 6, 7, 2, 1, 10, 11, 3, 13, 14, 15, 1); \\ k = 19 &: (1, 5, 6, 7, 2, 1, 10, 11, 3, 13, 14, 15, 1, 17, 2, 19, 5, 21, 22); \\ k = 23 &: (5, 6, 7, 2, 1, 10, 11, 3, 13, 14, 15, 1, 17, 2, 19, 5, 21, 22, 23, 6, 1, 26, 3), \\ & (6, 7, 2, 1, 10, 11, 3, 13, 14, 15, 1, 17, 2, 19, 5, 21, 22, 23, 6, 1, 26, 3, 7). \end{aligned}$$

We shall prove Lemma 6 in section 3.

Lemma 7. *Let k be a prime with $29 \leq k \leq 97$ and Q_0 a prime dividing d . Assume (2) with $k \nmid d$ and $k \nmid n + id$ for $0 \leq i < k - k'$ and $k' \leq i < k$ where $k' < k$ are consecutive primes. Then there are primes Q_1 and Q_2 given in the following table such that either $Q_1|d$ or $Q_2|d$.*

k	Q_0	(Q_1, Q_2)	k	Q_0	(Q_1, Q_2)
$29 \leq k \leq 59$	19	(7, 17)	73, 79	23	(53, 67)
$31 \leq k \leq 59$	29	(7, 17)	79	73	(53, 67)
61	59	(11, 61)	83	37	(23, 73)
67, 71	43	(53, 67)	89	79	(23, 73)
71	67	(43, 53)	97	23	(73, 97), (37, 83)

The proofs of Lemmas 6 and 7 depend on the repeated application of Lemma 5 and Corollary 1. We shall prove Lemma 7 in section 4. Next we apply Lemmas 1, 2 and 7 to prove the following result.

Lemma 8. *Let k be a prime with $7 \leq k \leq 97$. Assume (2) with $k \nmid d$. Further for $k \geq 29$, assume that $k \nmid n + id$ for $0 \leq i < k - k'$ and $k' \leq i < k$ where $k' < k$ are consecutive primes. Let (q_1, q_2) be as in Lemma 6. Then $q_1 \nmid d$ and $q_2 \nmid d$.*

The section 5 contains a proof of Lemma 8. Assume that $3 \nmid d$ and $5 \nmid d$. We define some more notation. For a subset $\mathcal{J} \subseteq [0, k) \cap \mathbb{Z}$, let

$$\begin{aligned} \mathcal{I}_3^0 = \mathcal{I}_3^0(\mathcal{J}) &:= \{i \in \mathcal{J} \mid i \equiv i_3 \pmod{3}\}, \quad \mathcal{I}_3^+ = \mathcal{I}_3^+(\mathcal{J}) := \{i \in \mathcal{J} \mid \left(\frac{i - i_3}{3}\right) = 1\}, \\ \mathcal{I}_3^- = \mathcal{I}_3^-(\mathcal{J}) &:= \{i \in \mathcal{J} \mid \left(\frac{i - i_3}{3}\right) = -1\} \end{aligned}$$

and

$$\mathcal{I}_5^+ = \mathcal{I}_5^+(\mathcal{J}) := \{i \in \mathcal{J} \mid \left(\frac{i - i_5}{5}\right) = 1\}, \quad \mathcal{I}_5^- = \mathcal{I}_5^-(\mathcal{J}) := \{i \in \mathcal{J} \mid \left(\frac{i - i_5}{5}\right) = -1\}.$$

Assume that $a_i \in \{1, 2, 7, 14\}$ for $i \in \mathcal{I}_3^+ \cup \mathcal{I}_3^-$. Then either $a_i \in \{1, 7\}$ for $i \in \mathcal{I}_3^+$, $a_i \in \{2, 14\}$ for $i \in \mathcal{I}_3^-$ or $a_i \in \{2, 14\}$ for $i \in \mathcal{I}_3^+$, $a_i \in \{1, 7\}$ for $i \in \mathcal{I}_3^-$. We define $(\mathcal{I}_3^1, \mathcal{I}_3^2) = (\mathcal{I}_3^+, \mathcal{I}_3^-)$ in the first case and $(\mathcal{I}_3^1, \mathcal{I}_3^2) = (\mathcal{I}_3^-, \mathcal{I}_3^+)$ in the latter. We observe that i 's have the same parity whenever $a_i \in \{2, 14\}$. Thus if i 's have the same parity

in one of \mathcal{I}_3^+ or \mathcal{I}_3^- but not in both, then we see that $(\mathcal{I}_3^1, \mathcal{I}_3^2) = (\mathcal{I}_3^+, \mathcal{I}_3^-)$ or $(\mathcal{I}_3^-, \mathcal{I}_3^+)$ according as i 's have the same parity in \mathcal{I}_3^- or \mathcal{I}_3^+ , respectively. Further we write

$$\mathcal{J}_1 = \mathcal{I}_3^1 \cap \mathcal{I}_5^+, \quad \mathcal{J}_2 = \mathcal{I}_3^1 \cap \mathcal{I}_5^-, \quad \mathcal{J}_3 = \mathcal{I}_3^2 \cap \mathcal{I}_5^+, \quad \mathcal{J}_4 = \mathcal{I}_3^2 \cap \mathcal{I}_5^-$$

and $\mathbf{a}_\mu = \{a_i | i \in \mathcal{J}_\mu\}$ for $1 \leq \mu \leq 4$. Since $\left(\frac{1}{5}\right) = \left(\frac{14}{5}\right) = 1$ and $\left(\frac{2}{5}\right) = \left(\frac{7}{5}\right) = -1$, we see that

$$(17) \quad (\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \mathbf{a}_4) \subseteq (\{1\}, \{7\}, \{14\}, \{2\}) \text{ or } (\{7\}, \{1\}, \{2\}, \{14\})$$

where $(\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \mathbf{a}_4) \subseteq (S_1, S_2, S_3, S_4)$ denotes $\mathbf{a}_\mu \subseteq S_\mu$, $1 \leq \mu \leq 4$. We use $7|(i - i')$ whenever $a_i, a_{i'} \in \{7, 14\}$ to exclude one of the above possibilities.

3. PROOF OF LEMMA 6

Let $k' < k$ be consecutive primes. We may suppose that if (2) holds for some $k > 29$, then $k \nmid d$ and $k \nmid a_i$ for $0 \leq i < k - k'$ and $k' \leq i < k$, otherwise the assertion follows from Theorem 4 with k replaced by k' . The subsections 3.1 to 3.10 will be devoted to the proof of Lemma 6. We may assume that $q_1 \nmid d$ and $q_2 \nmid d$ otherwise the assertion follows.

3.1. The case $k = 7$. Then $5 \nmid d$. By taking mirror images (4) of (2), there is no loss of generality in assuming that $5|n + i_5d, 7|n + i_7d$ for some pair (i_5, i_7) with $0 \leq i_5 < 5, 0 \leq i_7 \leq 3$. Further we may suppose $i_7 \geq 1$, otherwise the assertion follows from the case $k = 6$. We apply Lemma 5 with $\mathcal{P}_0 = \emptyset, p_1 = 5, p_2 = 7, (i_1, i_2) = (i_5, i_7), \mathcal{I} = [0, k] \cap \mathbb{Z}, \mathcal{P} = \Lambda(5, 7) = \{2\}$ and $\ell \leq \ell_1 = \lceil \frac{k}{2} \rceil$ to conclude that either

$$|\mathcal{I}_1| \leq \ell_1, \quad \mathcal{I}_1 \text{ is covered by } \mathcal{P}, \quad \mathcal{I}_2 = \{i \in \mathcal{I}' | \mathbf{i}(\mathcal{P}) \text{ is even}\}$$

or

$$|\mathcal{I}_2| \leq \ell_1, \quad \mathcal{I}_2 \text{ is covered by } \mathcal{P}, \quad \mathcal{I}_1 = \{i \in \mathcal{I}' | \mathbf{i}(\mathcal{P}) \text{ is even}\}.$$

Let $(i_5, i_7) = (3, 1)$. Then $\mathcal{I}_1 = \{0, 2, 6\}$ and $\mathcal{I}_2 = \{4, 5\}$. We see that \mathcal{I}_1 is covered by \mathcal{P} and hence $\mathbf{i}(\mathcal{P})$ is even for $i \in \mathcal{I}_2$. Thus $2 \nmid a_i$ for $i \in \mathcal{I}_2$. Therefore $a_4, a_5 \in \{1, 3\}$ and $a_0, a_2, a_6 \in \{2, 6\}$. If $a_0 = 6$ or $a_6 = 6$, then $3 \nmid a_4 a_5$ so that $a_4 = a_5 = 1$. This is not possible by modulo 3. Thus $a_0 = a_6 = 2$. Since $\left(\frac{a_0}{5}\right) \left(\frac{a_2}{5}\right) = \left(\frac{(-3d)(-d)}{5}\right) = -1$, we get $a_2 = 6$. Hence $a_4 = 1$. Further $a_5 = 3$ since $\left(\frac{a_5}{5}\right) \left(\frac{a_4}{5}\right) = \left(\frac{(2d)(1d)}{5}\right) = -1$. Also $5|a_3$ and $7|a_1$, otherwise the assertion follows from the results [MS03] for $k = 5$ and [BBGH06] for $k = 6$, respectively, stated in section 1. In fact $a_1 = 7, a_3 = 5$ by $\gcd(a_1 a_3, 6) = 1$. Thus $(a_0, a_1, a_2, a_3, a_4, a_5, a_6) = (2, 7, 6, 5, 1, 3, 2)$. The proofs for the other cases of (i_5, i_7) are similar. We get $(a_0, \dots, a_6) = (1, 5, 6, 7, 2, 1, 10)$ when $(i_5, i_7) = (1, 3)$, $(a_0, \dots, a_6) = (1, 2, 7, 6, 5, 1, 3)$ when $(i_5, i_7) = (4, 2)$ and all the other pairs are excluded. Hence Lemma 6 with $k = 7$ follows.

3.2. The case $k = 11$. Then $5 \nmid d$. By taking mirror images (4) of (2), there is no loss of generality in assuming that $5|n + i_5d, 11|n + i_{11}d$ for some pair (i_5, i_{11}) with $0 \leq i_5 < 5, 4 \leq i_{11} \leq 5$. We apply Lemma 5 with $\mathcal{P}_0 = \emptyset, p_1 = 5, p_2 = 11, (i_1, i_2) = (i_5, i_{11}), \mathcal{I} = [0, k] \cap \mathbb{Z}, \mathcal{P} = \Lambda(5, 11) = \{3\}$ and $\ell \leq \ell_1 = \lceil \frac{k}{3} \rceil$ to derive that either

$$|\mathcal{I}_1| \leq \ell_1, \mathcal{I}_1 \text{ is covered by } \mathcal{P}, \mathcal{I}_2 = \{i \in \mathcal{I}' | \mathbf{i}(\mathcal{P}) \text{ is even}\}$$

or

$$|\mathcal{I}_2| \leq \ell_1, \mathcal{I}_2 \text{ is covered by } \mathcal{P}, \mathcal{I}_1 = \{i \in \mathcal{I}' | \mathbf{i}(\mathcal{P}) \text{ is even}\}.$$

We compute $\mathcal{I}_1, \mathcal{I}_2$ and we restrict to those pairs (i_5, i_{11}) for which $\min(|\mathcal{I}_1|, |\mathcal{I}_2|) \leq \ell_1$ and either \mathcal{I}_1 or \mathcal{I}_2 is covered by \mathcal{P} . We find that $(i_5, i_{11}) = (0, 4), (1, 5)$. Let $(i_5, i_{11}) = (0, 4)$. Then $\mathcal{I}_1 = \{3, 9\}$ is covered by \mathcal{P} , $i_3 = 0$ and $\mathbf{i}(\mathcal{P})$ is even for $i \in \mathcal{I}_2 = \{1, 2, 6, 7, 8\}$. Thus $3 \nmid a_i$ for $i \in \mathcal{I}_2$. Further $p \in \{2, 7\}$ whenever $p|a_i$ with $i \in \mathcal{I}_2$. Therefore $a_i \in \{1, 2, 7, 14\}$ for $i \in \mathcal{I}_2$. By taking $\mathcal{J} = \mathcal{I}_2$, we have $\mathcal{I}_2 = \mathcal{I}_3^0 \cup \mathcal{I}_3^+ \cup \mathcal{I}_3^-$ and $\mathcal{I}_1 = \mathcal{I}_5^+ \cup \mathcal{I}_5^-$ with

$$\mathcal{I}_3^0 = \{6\}, \mathcal{I}_3^+ = \{1, 7\}, \mathcal{I}_3^- = \{2, 8\}, \mathcal{I}_5^+ = \{1, 6\}, \mathcal{I}_5^- = \{2, 7, 8\}.$$

Let $(\mathcal{I}_3^1, \mathcal{I}_3^2) = (\mathcal{I}_3^+, \mathcal{I}_3^-)$. Then

$$\mathcal{J}_1 = \{1\}, \mathcal{J}_2 = \{7\}, \mathcal{J}_3 = \emptyset, \mathcal{J}_4 = \{2, 8\}.$$

The possibility $(\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \mathbf{a}_4) \subseteq (\{7\}, \{1\}, \{2\}, \{14\})$ is excluded since $7|(i - i')$ whenever $a_i, a_{i'} \in \{7, 14\}$. Therefore $a_1 = 1, a_7 = 7, a_2 = a_8 = 2$. Further $a_6 = 1$ since $6 \in \mathcal{I}_5^+$ and $a_1 = 1, a_7 = 7$. This is not possible since $1 = \left(\frac{a_6}{7}\right) \left(\frac{a_8}{7}\right) = \left(\frac{(-d)(d)}{7}\right) = -1$.

Let $(\mathcal{I}_3^1, \mathcal{I}_3^2) = (\mathcal{I}_3^-, \mathcal{I}_3^+)$. Then we argue as above to conclude that $a_2 = a_8 = 1, a_1 = 2, a_7 = 14$ which is not possible since $n + 2d$ and $n + 8d$ cannot both be odd squares. The other case $(i_5, i_{11}) = (1, 5)$ is excluded similarly.

3.3. The cases $13 \leq k \leq 23$. Then $11 \nmid d$ and $13 \nmid d$. There is no loss of generality in assuming that $11|n + i_{11}d, 13|n + i_{13}d$ for some pair (i_{11}, i_{13}) with $0 \leq i_{11} < 11, 0 \leq i_{13} \leq \frac{k-1}{2}$ and further $i_{13} \geq 2$ if $k = 13$. We have applied Lemma 5 once in each of cases $k = 7$ and $k = 11$ but we apply it twice for every case $13 \leq k \leq 23$ in this section. Let $\mathcal{P}_0 = \emptyset, p_1 = 11, p_2 = 13, (i_1, i_2) = (i_{11}, i_{13}), \mathcal{I} = [0, k] \cap \mathbb{Z}, \mathcal{P} = \mathcal{P}_1 := \Lambda(11, 13)$ and $\ell \leq \ell_1$ where $\ell_1 = 3$ if $k = 13$; $\ell_1 = \lceil \frac{k}{5} \rceil + \lceil \frac{k}{17} \rceil$ if $k > 13$. Then $\ell_1 < \frac{1}{2}|\mathcal{I}'|$ since $|\mathcal{I}'| \geq k - \lceil \frac{k}{11} \rceil - \lceil \frac{k}{13} \rceil$. By Corollary 1, we derive that \mathcal{I}' is partitioned into $\mathcal{M} =: \mathcal{M}_1$ and $\mathcal{B} =: \mathcal{B}_1$ such that $(\mathcal{M}_1, \mathcal{B}_1, \mathcal{P}_1, \ell_1)$ has *Property* \mathfrak{H} . Now we restrict to all such pairs (i_{11}, i_{13}) satisfying $|\mathcal{M}_1| \leq \ell_1$ and \mathcal{M}_1 is covered by \mathcal{P}_1 . We check that $|\mathcal{M}_1| > 2$. Therefore $5 \nmid d$ since \mathcal{M}_1 is covered by \mathcal{P}_1 . Thus there exists i_5 with $0 \leq i_5 < 5$ such that $5|n + i_5d$.

Now we apply Lemma 5 with $p_1 = 5, p_2 = 11$ and partition $\mathcal{B}_1(5, 11)$ into two subsets. Let $\mathcal{P}_0 = \Lambda(11, 13) \cup \{11, 13\}, (i_1, i_2) = (i_5, i_{11}), \mathcal{I} = \mathcal{B}_1, \mathcal{P} = \mathcal{P}_2 := \Lambda(5, 11) \subseteq \{3, 19, 23\}$ and $\ell \leq \ell_2$ where $\ell_2 = 5, 6, 8, 11$ if $k = 13, 17, 19, 23$, respectively. Hence \mathcal{B}'_1 is partitioned into \mathcal{I}_1 and \mathcal{I}_2 satisfying either

$$|\mathcal{I}_1| \leq \ell_2, \mathcal{I}_1 \text{ is covered by } \mathcal{P}_2, \mathcal{I}_2 = \{i \in \mathcal{I}' | \mathbf{i}(\mathcal{P}_2) \text{ is even}\}$$

or

$$|\mathcal{I}_2| \leq \ell_2, \mathcal{I}_2 \text{ is covered by } \mathcal{P}_2, \mathcal{I}_1 = \{i \in \mathcal{I}' \mid i(\mathcal{P}_2) \text{ is even}\}.$$

We compute $\mathcal{I}_1, \mathcal{I}_2$ and we restrict to those pairs (i_{11}, i_{13}) for which $\min(|\mathcal{I}_1|, |\mathcal{I}_2|) \leq \ell_2$ and either \mathcal{I}_1 or \mathcal{I}_2 is covered by \mathcal{P}_2 . We find that $(i_{11}, i_{13}) = (4, 2), (5, 3)$ if $k = 13$; $(0, 0), (5, 3)$ if $k = 17$; $(0, 0), (0, 9), (7, 5), (7, 9), (8, 6), (9, 7), (10, 8)$ if $k = 19$ and $(0, 0), (0, 9), (1, 10), (2, 11), (4, 0), (5, 1), (5, 7), (6, 2), (6, 8), (7, 9), (8, 10), (9, 11)$ if $k = 23$.

Let (i_{11}, i_{13}) be such a pair. We write M for the one of \mathcal{I}_1 or \mathcal{I}_2 which is covered by \mathcal{P}_2 and B for the other. For $i \in \mathcal{B}'_1$, we see that $p \nmid a_i$ whenever $p \in \mathcal{P}_0$ since $17|a_i$ implies $5|a_i$. Therefore

$$(18) \quad i(\mathcal{P}_2) \text{ is even for } i \in B \text{ and } p \nmid a_i \text{ for } i \in B \text{ whenever } p \in \mathcal{P}_0,$$

since $B \subseteq \mathcal{B}'_1$. Further we check that $|M| > 1$ if $k \neq 23$ and > 3 if $k = 23$ implying $3 \nmid d$.

By taking $\mathcal{J} = B$, we get $B = \mathcal{I}_3^0 \cup \mathcal{I}_3^+ \cup \mathcal{I}_3^-$ and $B = \mathcal{I}_5^+ \cup \mathcal{I}_5^-$. Then $p \in \{2, 7\}$ whenever $p|a_i$ with $i \in \mathcal{I}_3^+ \cup \mathcal{I}_3^-$ by (18). By computing $\mathcal{I}_3^+, \mathcal{I}_3^-$, we find that i 's have the same parity in exactly one of $\mathcal{I}_3^+, \mathcal{I}_3^-$. Therefore we get from (17) that

$$(\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \mathbf{a}_4) \subseteq (\{1\}, \{7\}, \{14\}, \{2\}) \text{ or } (\{7\}, \{1\}, \{2\}, \{14\}).$$

Let $k = 13$ and $(i_{11}, i_{13}) = (4, 2)$. Then we have $\mathcal{M}_1 = \{0, 5, 10\}$, $i_5 = 0$, $M = \{3, 9, 12\}$ and $B = \{1, 6, 7, 8, 11\}$ since the latter set is not covered by $\mathcal{P}_2 = \{3\}$. Further $i_3 = 0$, $\mathcal{I}_3^0 = \{6\}$, $\mathcal{I}_3^+ = \mathcal{I}_3^- = \{8, 11\}$, $\mathcal{I}_3^2 = \mathcal{I}_3^+ = \{1, 7\}$, $\mathcal{I}_5^+ = \{1, 6, 11\}$, $\mathcal{I}_5^- = \{7, 8\}$, $\mathcal{J}_1 = \{11\}$, $\mathcal{J}_2 = \{8\}$, $\mathcal{J}_3 = \{1\}$, $\mathcal{J}_4 = \{7\}$. Therefore $a_{11} = 1, a_8 = 7, a_1 = 14, a_7 = 2$ or $a_{11} = 7, a_8 = 1, a_1 = 2, a_7 = 14$. The second possibility is excluded since $a_{11} = 7, a_7 = 14$ is not possible. Further from (18), we get $a_6 = 1$ since $2 \nmid a_6$ and $7 \nmid a_6$. Since $13|n + 2d$ and $7|n + d$, we get $\left(\frac{i-2}{13}\right) = \left(\frac{a_i a_6}{13}\right) = \left(\frac{a_i}{13}\right)$ and $-\left(\frac{i-1}{7}\right) = \left(\frac{a_i a_6}{7}\right) = \left(\frac{a_i}{7}\right)$. We observe that $13|n + 2d, 11|n + 4d, 7|n + d, 5|n, 3|n, 2|n + d, 5|a_i$ for $i \in \mathcal{M}$ and $3|a_i$ for $i \in \mathcal{M}_1$. Now we see that $a_0 \in \{5, 15\}$ and $a_0 = 5$ is excluded since $\left(\frac{5}{7}\right) \neq -\left(\frac{-1}{7}\right)$. Thus $a_0 = 15$. Next $a_1 = 14, a_2 = 13$ and $a_3 = 3$. Also $a_4 \in \{1, 11\}$ and $a_4 \neq 1$ since $\left(\frac{a_4}{13}\right) = \left(\frac{2}{13}\right) = -1$. Similarly we derive that $a_5 = 10, a_6 = 1, a_7 = 2, a_8 = 7, a_9 = 6, a_{10} = 5, a_{11} = 1$ and $a_{12} = 3$. Thus $(a_0, a_1, \dots, a_{12}) = (15, 14, 13, \dots, 5, 1, 3)$. The other case $(i_{11}, i_{13}) = (5, 3)$ is similar and we get $(a_0, a_1, \dots, a_{12}) = (1, 15, 14, \dots, 5, 1)$.

Let $k = 17$ and $(i_{11}, i_{13}) = (0, 0)$. Then we have $\mathcal{M}_1 = \{5, 10, 15\}$ and $i_5 = 0$. We see from the assumption of Lemma 6 with $k = 17, k' = 13$ that $4 \leq i_{17} < 13$. Hence, from $i_{17} \in \bigcup_{p=5,11,13} \{i_p + pj : 0 \leq j < \lceil \frac{k}{p} \rceil\}$, we get $i_{17} \in \{5, 10, 11\}$. Further $M = \{3, 6, 12\}$, $B = \{1, 2, 4, 7, 8, 9, 14, 16\}$, $i_3 = 0$, $\mathcal{I}_3^0 = \{9\}$, $\mathcal{I}_3^1 = \{1, 4, 7, 16\}$, $\mathcal{I}_3^2 = \{2, 8, 14\}$, $\mathcal{I}_5^+ = \{1, 4, 9, 14, 16\}$, $\mathcal{I}_5^- = \{2, 7, 8\}$, $\mathcal{J}_1 = \{1, 4, 16\}$, $\mathcal{J}_2 = \{7\}$, $\mathcal{J}_3 = \{14\}$ and $\mathcal{J}_4 = \{2, 8\}$. Therefore $a_1 = a_4 = a_{16} = 1, a_7 = 7, a_{14} = 14, a_2 = a_8 = 2$. Thus $a_9 = 1$ by (18) and $2 \nmid a_9, 7 \nmid a_9$. Now we see by Legendre symbol mod 17 that $a_1 = a_4 = a_9 = a_{16} = 1$ is not possible. The case $(i_{11}, i_{13}) = (5, 3)$ is excluded similarly.

Let $k = 19$ and $(i_{11}, i_{13}) = (0, 0)$. Then we have $\mathcal{M}_1 = \{5, 10, 15, 17\}$, $i_5 = 0, i_{17} = 0$, $M = \{3, 6, 12\}$, $B = \{1, 2, 4, 7, 8, 9, 14, 16, 18\}$ and $i_3 = 0$. We see from $i_{19} \in \bigcup_{p=3,5,11,13,17} \{i_p + pj : 0 \leq j < \lceil \frac{k}{p} \rceil\}$ and $2 \leq i_{19} < 17$ that $i_{19} \in \{3, 5, 6, 9, 10, 11, 12, 13, 15\}$. Further $\mathcal{I}_3^0 = \{9, 18\}$, $\mathcal{I}_3^1 = \{1, 4, 7, 16\}$, $\mathcal{I}_3^2 = \{2, 8, 14\}$, $\mathcal{I}_5^+ = \{1, 4, 9, 14, 16\}$, $\mathcal{I}_5^- = \{2, 7, 8, 18\}$, $\mathcal{J}_1 = \{1, 4, 16\}$, $\mathcal{J}_2 = \{7\}$, $\mathcal{J}_3 = \{14\}$ and $\mathcal{J}_4 = \{2, 8\}$. Therefore $a_1 = a_4 = a_{16} = 1$ which is not possible by mod 19. The case $(i_{11}, i_{13}) = (7, 5)$ is excluded similarly. Let $(i_{11}, i_{13}) = (0, 9)$. Then $\mathcal{M}_1 = \{2, 5, 7, 12, 17\}$, $i_5 = 2, i_{17} = 5$, $M = \{1, 3, 10, 16\}$, $B = \{4, 6, 8, 13, 14, 15, 18\}$, $i_3 = 1$ and $i_{19} = 3$. We now consider $(n + 6d)(n + 7d) \cdots (n + 18d) = b'y'^2$. Then $P(b') \leq 13$. By the case $k = 13$, we get $(a_6, a_7, \dots, a_{18}) = (1, 15, \dots, 6, 5, 1)$ since $5|a_7$ and $3|a_{16}$. From $19|n + 3d$, we get $\left(\frac{a_i}{19}\right) = \left(\frac{a_i a_6}{19}\right) = -\left(\frac{i-3}{19}\right)$ which together with $13|n + 9d, 11|n, 7|n + d, 2|n, 5|a_2, 17|a_5, 3|a_1$ implies $a_0 \in \{2, 22\}$, $a_1 \in \{3, 21\}$, $a_2 = 5, a_3 = 19, a_4 = 2$ and $a_5 = 17$. Now from $\left(\frac{a_i}{17}\right) = \left(\frac{a_i a_6}{17}\right) = \left(\frac{i-5}{17}\right)$, we get $a_0 = 22, a_1 = 21$. Thus $(a_0, a_1, \dots, a_{18}) = (22, 21, \dots, 6, 5, 1)$. The case $(i_{11}, i_{13}) = (7, 9)$ is similar and we get $(a_0, a_1, \dots, a_{18}) = (1, 5, 6, \dots, 21, 22)$. For the pair $(i_{11}, i_{13}) = (10, 8)$, we get similarly $(a_0, a_1, \dots, a_{18}) = (21, 5, \dots, 6, 5, 1, 3)$. This is excluded by considering $(n + 3d)(n + 6d) \cdots (n + 18d)$ and $k = 6$. For the pairs $(i_{11}, i_{13}) = (8, 6), (9, 7)$, we get $i_{19} = 0, 1$, respectively, which is not possible since $i_{19} \geq 2$ by the assumption of the Lemma.

Let $k = 23$ and $(i_{11}, i_{13}) = (0, 0)$. Then $\mathcal{M}_1 = \{5, 10, 15, 17, 20\}$, $i_5 = 0, i_{17} = 0$, $M = \{3, 6, 12, 19, 21\}$, $B = \{1, 2, 4, 7, 8, 9, 14, 16, 18\}$, $i_3 = 0$ and $i_{19} = 0$ since $23 \nmid a_{19}$. We have $i_{23} \in \{5, 6, 9, 10, 11, 12, 13, 15, 17, 18\}$ since $4 \leq i_{23} < 19$. Here we observe that $23 \nmid a_{19}$ and $4 \leq i_{23} < 19$ in view of our assumption that $k \nmid a_i$ for $0 \leq i < k - k'$ and $k' \leq i < k$ with $k = 23, k' = 19$. Further $\mathcal{I}_3^0 = \{9, 18\}$, $\mathcal{I}_3^1 = \{1, 4, 7, 16\}$, $\mathcal{I}_3^2 = \{2, 8, 14\}$, $\mathcal{I}_5^+ = \{1, 4, 9, 14, 16\}$, $\mathcal{I}_5^- = \{2, 7, 8, 18\}$, $\mathcal{J}_1 = \{1, 4, 16\}$, $\mathcal{J}_2 = \{7\}$, $\mathcal{J}_3 = \{14\}$ and $\mathcal{J}_4 = \{2, 8\}$. Therefore $a_1 = a_4 = a_{16} = 1, a_7 = 7, a_{14} = 14, a_2 = a_8 = 2$. This is not possible since $\left(\frac{a_1}{23}\right) = \left(\frac{a_4}{23}\right) = \left(\frac{a_{16}}{23}\right) = \left(\frac{a_2}{23}\right) = \left(\frac{a_8}{23}\right) = 1$. The cases $(i_{11}, i_{13}) = (0, 9), (1, 10), (2, 11), (4, 0), (7, 9), (8, 10), (9, 11)$ are excluded similarly. Let $(i_{11}, i_{13}) = (5, 1)$. Then $\mathcal{M}_1 = \{7, 10, 12, 17, 22\}$, $i_5 = 2, i_{17} = 10, M = \{0, 3, 4, 6, 8, 15, 21\}$, $B = \{9, 11, 13, 18, 19, 20\}$ and $i_3 = 0$. This implies either $23|a_4, 19|a_8$ or $23|a_8, 19|a_4$. Further $\mathcal{I}_3^0 = \{9, 18\}$, $\mathcal{I}_3^1 = \{11, 20\}$, $\mathcal{I}_3^2 = \{13, 19\}$, $\mathcal{I}_5^+ = \{11, 13, 18\}$, $\mathcal{I}_5^- = \{9, 19, 20\}$, $\mathcal{J}_1 = \{11\}$, $\mathcal{J}_2 = \{20\}$, $\mathcal{J}_3 = \{13\}$ and $\mathcal{J}_4 = \{19\}$. Therefore $a_{11} = 1, a_{20} = 7, a_{13} = 14, a_{19} = 2$. Further from (18), we get $a_9 \in \{1, 2\}, a_{18} = 1$ since $7 \nmid a_9 a_{18}, 2 \nmid a_{18}$. However $a_9 = 2$ as $9 \in \mathcal{I}_5^-, 18 \in \mathcal{I}_5^+$. Since $\left(\frac{a_{11}}{23}\right) = \left(\frac{a_{18}}{23}\right) = 1$, we see that $23|a_4, 19|a_8$. By using $\left(\frac{a_i}{p}\right) = \left(\frac{a_i a_{11}}{p}\right) = \left(\frac{(i-i_p)(11-i_p)}{p}\right)$, we get $\left(\frac{a_i}{23}\right) = -\left(\frac{i-4}{23}\right), \left(\frac{a_i}{11}\right) = -\left(\frac{i-5}{11}\right), \left(\frac{a_i}{7}\right) = -\left(\frac{i-6}{7}\right)$ and $\left(\frac{a_i}{5}\right) = \left(\frac{i-2}{5}\right)$. Now from $23|a_4, 19|a_8, 17|a_{10}, 13|n + d, 11|n + 5d, 7|n + 6d, 5|n + 2d, 3|n, 2|n + d$, \mathcal{M}_1 is covered by $\{5, 17\}$, M is covered by $\{3, 19, 23\}$, we derive that $(a_0, a_1, \dots, a_{22}) = (3, 26, \dots, 6, 5)$. The pairs $(i_{11}, i_{13}) = (5, 7), (6, 2), (6, 8)$ are similar and we get $(a_0, a_1, \dots, a_{22}) = (6, 7, \dots, 3, 7), (7, 3, \dots, 7, 6), (5, 6, 7, \dots, 3)$, respectively.

3.4. Introductory remarks on the cases $k \geq 29$. Assume $q_1 \nmid d$ and $q_2 \nmid d$. Then, by taking mirror image (4) of (2), there is no loss of generality in assuming that

$q_1|n + i_{q_1}d, q_2|n + i_{q_2}d$ for some pair (i_{q_1}, i_{q_2}) with $0 \leq i_{q_1} < q_1, 0 \leq i_{q_2} \leq \frac{k-1}{2}$ and further $i_{q_2} \geq k - k'$ if $q_2 = k$. For $k = 61$, by taking $(n + 8d) \cdots (n + 60d)$ and $k = 53$, we may assume that $\max(i_{59}, i_{61}) \geq 8$ if $i_{59} \geq 2$. Let $\mathcal{P}_0 = \emptyset, p_1 = q_1, p_2 = q_2, (i_1, i_2) = (i_{q_1}, i_{q_2}), \mathcal{I} = [0, k] \cap \mathbb{Z}, \mathcal{P} = \mathcal{P}_1 := \Lambda(q_1, q_2)$ and $\ell \leq \ell_1 = \sum_{p \in \mathcal{P}_1} \left\lceil \frac{k}{p} \right\rceil$. We check that $\ell_1 < \frac{1}{2}|\mathcal{I}'|$ since $|\mathcal{I}'| \geq k - \left\lceil \frac{k}{q_1} \right\rceil - \left\lceil \frac{k}{q_2} \right\rceil$. By Corollary 1, we get $\mathcal{M} =: \mathcal{M}_1$ and $\mathcal{B} =: \mathcal{B}_1$ with $(\mathcal{M}_1, \mathcal{B}_1, \mathcal{P}_1, \ell_1)$ having *Property* \mathfrak{H} . We now restrict to all such pairs (i_{q_1}, i_{q_2}) for which $|\mathcal{M}_1| \leq \ell_1$ and \mathcal{M}_1 is covered by \mathcal{P}_1 . We find that there is no such pair (i_{q_1}, i_{q_2}) when $k = 97$.

3.5. The cases $29 \leq k \leq 59$. As stated in Lemma 6, we have $q_1 = 19, q_2 = 29$ and $\mathcal{P}_1 = \Lambda(19, 29) \subseteq \{11, 13, 17, 43, 47, 53, 59\}$. Then the pairs (i_{q_1}, i_{q_2}) are given by

$$\begin{aligned} k = 29 &: (0, 9), (1, 10), (2, 11), (3, 12), (4, 13), (15, 5), (16, 6), (17, 7), (18, 8); \\ k = 31 &: (0, 0), (0, 9), (1, 10), (2, 11), (3, 12), (4, 13), (11, 1), \\ &\quad (12, 2), (13, 3), (14, 4), (15, 5), (16, 6), (17, 7), (18, 8); \\ k = 37 &: (0, 0), (0, 9), (1, 10), (2, 11), (3, 12), (4, 13), (17, 7), (18, 8); \\ k = 41 &: (0, 0), (2, 11), (3, 12), (4, 13); \\ k = 43 &: (0, 0), (1, 1), (3, 12), (4, 13), (5, 14), (6, 15), (7, 16), (8, 17); \\ k = 47 &: (0, 0), (1, 1), (7, 16), (8, 17), (9, 18), (10, 19), (11, 20), \\ &\quad (12, 21), (13, 22), (13, 23), (14, 23); \\ k = 53 &: (0, 0), (1, 0), (1, 1), (13, 22), (13, 23), (14, 23), (14, 24), \\ &\quad (15, 24), (15, 25), (16, 25), (16, 26), (17, 26); \\ k = 59 &: (0, 0), (0, 28), (1, 0), (1, 1), (2, 1), (3, 2), (17, 27), (18, 28). \end{aligned}$$

Let $k = 31$ and $(i_{19}, i_{29}) = (0, 9)$. We see that $\mathcal{P}_1 = \{11, 13, 17\}, \mathcal{M}_1 = \{4, 5, 12, 16, 21, 25, 27\}$ and $\mathcal{B}_1 = \{1, 2, 3, 6, 7, 8, 10, 11, 13, 14, 15, 17, 18, 20, 22, 23, 24, 26, 28, 29, 30\}$. Since \mathcal{M}_1 is covered by \mathcal{P}_1 , we get 11 divides a_5, a_{16}, a_{27} ; 13 divides a_{12}, a_{25} and 17 divides a_4, a_{21} so that $i_{11} = 5, i_{13} = 12, i_{17} = 4$. We see that $\gcd(11 \cdot 13 \cdot 17, a_i) = 1$ for $i \in \mathcal{B}_1$. Now we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \{19, 29\}, p_1 = 11, p_2 = 13, (i_1, i_2) := (i_{11}, i_{13}) = (5, 12), \mathcal{I} = \mathcal{B}_1, \mathcal{P} = \mathcal{P}_2 := \Lambda(11, 13) \setminus \mathcal{P}_0 = \{5, 31\}$ and $\ell \leq \ell_2 = \sum_{p \in \mathcal{P}_2} \left\lceil \frac{k}{p} \right\rceil = 8$. Thus $|\mathcal{I}'| = |\mathcal{B}_1| = 21 > 2\ell_2$. Then the condition of Corollary 1 are satisfied and we have $\mathcal{M} =: \mathcal{M}_2, \mathcal{B} =: \mathcal{B}_2$ and $(\mathcal{M}_2, \mathcal{B}_2, \mathcal{P}_2, \ell_2)$ has *Property* \mathfrak{H} . We get $\mathcal{M}_2 = \{1, 3, 7, 8, 18, 23, 28\}$. This is not possible since \mathcal{M}_2 is not covered by \mathcal{P}_2 . Further the following pairs (i_{19}, i_{29}) are excluded similarly:

$$\begin{aligned} k = 29 &: (0, 9), (1, 10), (2, 11), (3, 12), (4, 13), (15, 5), (16, 6), (17, 7), (18, 8); \\ k = 31 &: (1, 10), (2, 11), (3, 12), (4, 13), (18, 8). \end{aligned}$$

Thus $k > 29$.

Let $k = 59$ and $(i_{19}, i_{29}) = (0, 0)$. Then we see that $\mathcal{P}_1 = \{11, 13, 17, 43, 47, 53, 59\}, \mathcal{M}_1 = \{11, 13, 17, 22, 26, 33, 34, 39, 43, 44, 47, 51, 52, 53, 55\}, \mathcal{B}_1 = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 14, 15, 16, 18, 20, 21, 23, 24, 25, 27, 28, 30, 31, 32, 35, 36, 37, 40, 41, 42, 45, 46, 48, 49, 50, 54, 56\}, i_{11} = i_{13} = i_{17} = 0, \{43, 47, 53\}$ is covered by $\{43, 47, 53, 59\} =: \mathcal{P}'_1$. Let

$p|a_i$ for $i \in \mathcal{B}_1$ and $p \in \mathcal{P}_1$. Then we show that $i \in \{4, 6, 10\}$. Let $59|a_{43}$. Then $\{47, 53\}$ is covered by $\{43, 47, 53\}$. Let $43|a_{47}$. If $43|a_i$ with $i \in \mathcal{B}_1$, then $i = 4$ and $43 \cdot p|a_4$ with $p \in \{47, 53\}$ since $i(\mathcal{P}_1)$ is even. This implies either $53|a_{53}, 43 \cdot 47|a_4$ or $47|a_{53}, 43 \cdot 53|a_4$. Similarly we get $i \in \{4, 6, 10\}$ by considering all the cases $59|a_{43}, 59|a_{47}$ and $59 \nmid a_{43}a_{47}a_{53}$. We observe that $59 \nmid a_{53}$ since $6 \leq i_{59} < 53$. Hence we conclude that $p \nmid a_i$ for $i \in \mathcal{B}_1 \setminus \{4, 6, 10\}$ and $p \in \mathcal{P}'_1$. Further we observe that

$$(19) \quad i_{59} \in \mathcal{M}_1 \cup \{19, 29, 38\} \cup \{6, 10\}.$$

Now we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \{19, 29\}$, $p_1 = 11, p_2 = 13$, $(i_1, i_2) := (0, 0)$, $\mathcal{I} = \mathcal{B}_1 \setminus \{4, 6, 10\}$, $\mathcal{P} = \mathcal{P}_2 := \Lambda(11, 13) \setminus \mathcal{P}_0 = \{5, 31, 37\}$ and $\ell \leq \ell_2 = \sum_{p \in \mathcal{P}_2} \left\lceil \frac{k}{p} \right\rceil = 16$. Thus $|\mathcal{I}'| = |\mathcal{B}_1| - 2 > 2\ell_2$. Then the conditions of Corollary 1 are satisfied and we have $\mathcal{M} =: \mathcal{M}_2$, $\mathcal{B} =: \mathcal{B}_2$ with $(\mathcal{M}_2, \mathcal{B}_2, \mathcal{P}_2, \ell_2)$ having *Property* \mathfrak{H} . We get $\mathcal{M}_2 = \{5, 15, 20, 30, 31, 35, 37, 40, 45\}$, $\mathcal{B}_2 = \{1, 2, 3, 7, 8, 9, 12, 14, 16, 18, 21, 23, 24, 25, 27, 28, 32, 36, 41, 42, 46, 48, 49, 50, 54, 56\}$, $i_5 = 0$, $31|a_{31}, 37|a_{37}$ or $31|a_{37}, 37|a_{31}$. Now we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \{19, 29\}$, $p_1 = 5, p_2 = 11$, $(i_1, i_2) := (0, 0)$, $\mathcal{I} = \mathcal{B}_2$, $\mathcal{P} = \mathcal{P}_3 := \Lambda(5, 11) \setminus \mathcal{P}_0 = \{3, 23, 41\}$ and $\ell \leq \ell_3 = \sum_{p \in \mathcal{P}_3} \left\lceil \frac{k}{p} \right\rceil$. Then by Lemma 5, we see that $M = \{3, 6, 12, 21, 23, 24, 27, 41, 42, 46, 48, 54\}$ is covered by \mathcal{P}_3 and $i(\mathcal{P}_3)$ is even for $i \in B = \{1, 2, 7, 8, 9, 14, 16, 18, 28, 32, 36, 49, 56\}$. Thus $i_3 = i_{23} = i_{41} = 0$ and $p \in \{2, 7\}$ whenever $p|a_i$ with $i \in B$. Putting $\mathcal{J} = B$, we have $B = \mathcal{I}_3^0 \cup \mathcal{I}_3^1 \cup \mathcal{I}_3^2$ and $B = \mathcal{I}_5^+ \cup \mathcal{I}_5^-$ with

$$\mathcal{I}_3^0 = \{9, 18, 36\}, \quad \mathcal{I}_3^1 = \{1, 7, 16, 28, 49\}, \quad \mathcal{I}_3^2 = \{2, 8, 14, 32, 56\}$$

and

$$\mathcal{I}_5^+ = \{1, 9, 14, 16, 36, 49, 56\}, \quad \mathcal{I}_5^- = \{2, 7, 8, 18, 28, 32\}.$$

so that

$$\mathcal{J}_1 = \{1, 16, 49\}, \quad \mathcal{J}_2 = \{7, 28\}, \quad \mathcal{J}_3 = \{14, 56\}, \quad \mathcal{J}_4 = \{2, 8, 32\}.$$

Hence $(\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \mathbf{a}_4) \subseteq (\{1\}, \{7\}, \{14\}, \{2\})$ by (17). Thus $a_1 = a_{16} = a_{49} = 1$, $a_7 = a_{28} = 7, a_{14} = a_{56} = 14, a_2 = a_8 = a_{32} = 2$. Further we get $a_9 = a_{36} = 1$ and $a_{18} = 2$ since $9, 36 \in \mathcal{I}_5^+$ and $18 \in \mathcal{I}_5^-$. Since

$$(20) \quad \left(\frac{a_i}{59} \right) = 1 \text{ for } a_i \in \{1, 7\},$$

we see that $\left(\frac{a_i}{59} \right) = 1$ for $i \in \{1, 7, 9, 16, 28, 36, 49\}$ which is not possible by (19).

Let $k = 41$ and $(i_{19}, i_{29}) = (2, 11)$. Then we see that $\mathcal{P}_1 = \{11, 13, 17\}$, $\mathcal{M}_1 = \{1, 6, 7, 14, 18, 23, 27, 29\}$, $\mathcal{B}_1 = \{0, 3, 4, 5, 8, 9, 10, 12, 13, 15, 16, 17, 19, 20, 22, 24, 25, 26, 28, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39\}$, $i_{11} = 7, i_{13} = 1, i_{17} = 6$. Further $\gcd(a_i, 11 \cdot 13 \cdot 17) = 1$ for $i \in \mathcal{B}_1$. Now we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \{19, 29\}$, $p_1 = 11, p_2 = 13$, $(i_1, i_2) := (7, 1)$, $\mathcal{I} = \mathcal{B}_1$, $\mathcal{P} = \mathcal{P}_2 := \Lambda(11, 13) \setminus \mathcal{P}_0 = \{5, 31, 37\}$ and $\ell \leq \ell_2 = \sum_{p \in \mathcal{P}_2} \left\lceil \frac{k}{p} \right\rceil = 13$. Then $|\mathcal{I}'| = |\mathcal{B}_1| > 2\ell_2$. Thus the conditions of Corollary 1 are satisfied and we get $\mathcal{M} =: \mathcal{M}_2$ and $\mathcal{B} =: \mathcal{B}_2$ such that $(\mathcal{M}_2, \mathcal{B}_2, \mathcal{P}_2, \ell_2)$ has *Property* \mathfrak{H} . We have $\mathcal{M}_2 = \{0, 3, 5, 9, 10, 20, 25, 30, 35\}$, $\mathcal{B}_2 = \{4, 8, 12, 13, 15, 16, 17, 19, 22, 24, 26, 28, 31, 32, 33, 34, 36, 37, 38, 39\}$, $i_5 = 0$. Further $31 \cdot 37|a_3a_9$, $31 \nmid a_{34}$. We take $\mathcal{P}_0 = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \{19, 29\}$, $p_1 = 5, p_2 = 11$, $(i_1, i_2) := (0, 7)$, $\mathcal{I} = \mathcal{B}_2$, $\mathcal{P} = \mathcal{P}_3 := \Lambda(5, 11) \setminus \mathcal{P}_0 = \{3, 23, 41\}$, $\ell \leq \sum_{p \in \mathcal{P}_3} \left\lceil \frac{k}{p} \right\rceil$ and apply Lemma 5 to see that $M = \{13, 16, 17, 19, 28, 34, 37\}$ is covered by \mathcal{P}_3 , $i_3 = 1$, $i(\mathcal{P}_3)$ is even for $i \in B = \{4, 8, 12, 22, 24, 26, 31, 32, 33, 36, 38, 39\}$.

Further $i_{23} = 17$, $i_{41} \in \{2, 11, 21\} \cup \mathcal{M}_1 \cup \mathcal{M}_2 \cup M \cup \{4, 22, 31\}$ or vice-versa. Here we observe that i_{41} exists since $41 \nmid d$. Thus $23 \cdot 41 \mid \prod a_i$ where i runs through the set $\{2, 11, 21\} \cup \mathcal{M}_1 \cup \mathcal{M}_2 \cup \{4, 22, 31\}$. Therefore $a_i \in \{1, 2, 7, 14\}$ for $i \in \mathcal{I}_3^1 \cup \mathcal{I}_3^2$ where $B = \mathcal{I}_3^0 \cup \mathcal{I}_3^1 \cup \mathcal{I}_3^2$, $B = \mathcal{I}_5^+ \cup \mathcal{I}_5^-$ with

$$\mathcal{I}_3^0 = \{4, 22, 31\}, \quad \mathcal{I}_3^1 = \{12, 24, 33, 36, 39\}, \quad \mathcal{I}_3^2 = \{8, 26, 32, 38\}$$

and

$$\mathcal{I}_5^+ = \{4, 24, 26, 31, 36, 39\}, \quad \mathcal{I}_5^- = \{8, 12, 22, 32, 33, 38\}$$

by taking $\mathcal{J} = B$. We get

$$\mathcal{J}_1 = \{24, 36, 39\}, \quad \mathcal{J}_2 = \{12, 33\}, \quad \mathcal{J}_3 = \{26\}, \quad \mathcal{J}_4 = \{8, 32, 38\},$$

and $a_{24} = a_{36} = a_{39} = 1$, $a_{12} = a_{33} = 7$, $a_{26} = 14$, $a_8 = a_{32} = a_{38} = 2$ by (17). Since

$$(21) \quad \left(\frac{a_i}{41}\right) = 1 \text{ for } a_i \in \{1, 2\},$$

we see that $\left(\frac{a_i}{41}\right) = 1$ for $i \in \{8, 24, 32, 36, 38, 39\}$ which is not valid by the possibilities of i_{41} .

All other cases are excluded similarly. Analogous to (20) and (21), we use $\left(\frac{a_i}{k}\right) = 1$ for

$$a_i \in \{1, 7\} \text{ if } k = 37, 53, 59; \quad a_i \in \{1, 2\} \text{ if } k = 31, 41, 47; \quad a_i \in \{1, 14\} \text{ if } k = 43$$

to exclude the remaining possibilities.

3.6. The case $k = 61$. We have $q_1 = 59, q_2 = 61$ and $\mathcal{P}_1 = \{7, 13, 17, 29, 47, 53\}$. Then the pairs (i_{q_1}, i_{q_2}) are given by $(8, 6), (9, 7), (10, 8), (11, 9)$, i.e. $(i + 2, i)$ with $6 \leq i \leq 9$.

Let $(i_{59}, i_{61}) = (8, 6)$. Then $\mathcal{P}_1 = \{7, 13, 17, 29, 47, 53\}$, $\mathcal{M}_1 = \{2, 4, 9, 11, 14, 15, 16, 20, 25, 28, 32, 33, 38, 39, 41, 46, 50, 53, 54, 60\}$, $\mathcal{B}_1 = \{0, 1, 3, 5, 7, 10, 12, 13, 17, 18, 19, 21, 22, 23, 24, 26, 27, 29, 30, 31, 34, 35, 36, 37, 40, 42, 43, 44, 45, 47, 48, 49, 51, 52, 55, 56, 57, 58, 59\}$, $i_7 = 4, i_{13} = 2, i_{17} = 16, i_{29} = 9$ and a_{14}, a_{20} are divisible by 47, 53. Further $\gcd(p, a_i) = 1$ for $i \in \mathcal{B}_1$ and $p \in \mathcal{P}_1$. Let $\mathcal{P}_0 = \mathcal{P}_1 \cup \{59, 61\}$, $p_1 = 7, p_2 = 17, (i_1, i_2) := (4, 16)$, $\mathcal{I} = \mathcal{B}_1$, $\mathcal{P} = \mathcal{P}_2 := \Lambda(7, 17) \setminus \mathcal{P}_0 = \{11, 19, 23, 37\}$ and $\ell \leq \ell_2 = \sum_{p \in \mathcal{P}_2} \left\lceil \frac{k}{p} \right\rceil = 15$. Then $2\ell_2 < |\mathcal{I}'| = |\mathcal{B}_1| - 1$. By Corollary 1, we get $\mathcal{M} =: \mathcal{M}_2$, $\mathcal{B} =: \mathcal{B}_2$ and $(\mathcal{M}_2, \mathcal{B}_2, \mathcal{P}_2, \ell_2)$ has *Property \mathfrak{H}* . We find that $\mathcal{M}_2 = \{1, 10, 12, 21, 23, 29, 30, 34, 44, 45, 48, 56\}$, $\mathcal{B}_2 = \{0, 3, 5, 7, 13, 17, 19, 22, 24, 26, 27, 31, 35, 36, 37, 40, 42, 43, 47, 49, 51, 52, 55, 57, 58, 59\}$, $i_{11} = 1, i_{19} = 10, i_{23} = 21, i_{37} = 30$. Now we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \{59, 61\}$, $p_1 = 11, p_2 = 59, (i_1, i_2) := (1, 8)$, $\mathcal{I} = \mathcal{B}_2$, $\mathcal{P} = \mathcal{P}_3 := \Lambda(11, 59) \setminus \mathcal{P}_0 = \{31, 41\}$ and $\ell \leq \ell_3 = \sum_{p \in \mathcal{P}_3} \left\lceil \frac{k}{p} \right\rceil = 4$. Then $2\ell_3 < |\mathcal{I}'| = |\mathcal{B}_2|$. By Corollary 1, we get $\mathcal{M} =: \mathcal{M}_3$ and $\mathcal{B} =: \mathcal{B}_3$ such that $(\mathcal{M}_3, \mathcal{B}_3, \mathcal{P}_3, \ell_3)$ has *Property \mathfrak{H}* . We get $\mathcal{M}_3 = \{0, 5, 26, 36\}$ which cannot be covered by \mathcal{P}_3 . This is a contradiction. The remaining cases are excluded similarly.

3.7. **The cases $k = 67, 71$.** We have $q_1 = 43, q_2 = 67$ and $\mathcal{P}_1 \subseteq \{11, 13, 19, 29, 31, 37, 41, 53, 71\}$. Then the pairs (i_{q_1}, i_{q_2}) are given by

$$k = 67 : (i, i), 6 \leq i \leq 33;$$

$$k = 71 : (i, i), 0 \leq i \leq 35, i \neq 24, 25 \text{ and } (24, 0), (25, 1), (26, 2), (27, 3).$$

Let $k = 71$ and $(i_{43}, i_{67}) = (27, 3)$. We see that $\mathcal{P}_1 = \{11, 13, 19, 29, 31, 37, 41, 53, 71\}$, $\mathcal{M}_1 = \{4, 5, 8, 12, 13, 15, 17, 18, 26, 29, 31, 32, 33, 37, 39, 41, 44, 48, 51, 57, 59\}$, $\mathcal{B}_1 = \{0, 1, 2, 6, 7, 9, 10, 11, 14, 16, 19, 20, 21, 22, 23, 24, 25, 28, 30, 34, 35, 36, 38, 40, 42, 43, 45, 46, 47, 49, 50, 52, 53, 54, 55, 56, 58, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69\}$, $i_{11} = 4, i_{13} = 5, i_{19} = 13$. Therefore $\{8, 12, 17, 29, 33, 39, 41\}$ is covered by $29, 31, 37, 41, 53, 71$ implying either $i_{29} = 12$ or $i_{29} \in \{17, 29, 33\}$, $i_{31} = 8$. Let $i \in \mathcal{B}_1$ and $p|a_i$ with $p \in \mathcal{P}_1$. Then there is a $q \in \mathcal{P}_1$ such that $pq|a_i$ since $i(\mathcal{P}_1)$ is even. Next we consider the case $i_{31} = 8$. Then $\{12, 17, 29, 33, 41\} =: \mathcal{M}'_1$ is covered by $29, 37, 41, 53, 71$ and $i_{29} \neq 12$. For $29 \in \mathcal{M}'_1$, we may suppose that either $29|a_{29}, 41|a_{17}, 29 \cdot 41|a_{58}$ or $29|a_{29}, 41|a_{41}, 29 \cdot 41|a_0$. Thus 0 or 58 in \mathcal{B}_1 correspond to 29. We argue as above that for any other element of \mathcal{M}'_1 , there is no corresponding element in \mathcal{B}_1 . For the first case, we derive similarly that $31|a_{33}, 37|a_{39}, 31 \cdot 37|a_2$ or $37|a_{17}, 37 \cdot 71|a_{54}$ or $37|a_{29}, 37 \cdot 71|a_{63}$ or $41|a_{17}, 37 \cdot 71|a_{58}$. Therefore

$$29 \cdot 31 \cdot 37 \cdot 41 \cdot 53 \cdot 71 \mid \prod (n + id) \text{ for } i \in \mathcal{M}_1 \cup \{3, 27, 70\} \cup \mathcal{B}'_1$$

where $\mathcal{B}'_1 = \{2, 54, 58, 63\}$ if $i_{29} = 12$ and $\{0, 58\}$ otherwise. Further

$$(22) \quad i_{71} \in \mathcal{M}_1 \cup \{27\} \cup \mathcal{B}'_1 \text{ and } i_{71} \neq 32.$$

For each possibility $i_{29} \in \{0, 4, 12, 17\}$, we now take $\mathcal{P}_0 = \mathcal{P}_1 \cup \{43, 67\}$, $p_1 = 19, p_2 = 29$, $(i_1, i_2) := (13, i_{29})$, $\mathcal{I} = \mathcal{B}_1 \setminus \mathcal{B}'_1$, $\mathcal{P} = \mathcal{P}_2 := \Lambda(19, 29) \setminus \mathcal{P}_0 = \{17, 47, 59, 61\}$ and $\ell = \ell_2 = \sum_{p \in \mathcal{P}_2} \left\lceil \frac{k}{p} \right\rceil = 11$. Then $|\mathcal{I}'| = |\mathcal{B}_1| - 4 > 2\ell_2$. Thus the conditions of Corollary 1 are satisfied and we get $\mathcal{M} =: \mathcal{M}_2$ and $\mathcal{B} =: \mathcal{B}_2$ with $(\mathcal{M}_2, \mathcal{B}_2, \mathcal{P}_2, \ell_2)$ having *Property \mathfrak{H}* . We check that $|\mathcal{M}_2| \leq \ell_2$ only at $i_{29} = 12$ in which case we get $\mathcal{M}_2 = \{9, 11, 19, 23, 36, 53\}$, $\mathcal{B}_2 = \{0, 1, 6, 7, 10, 14, 6, 20, 21, 22, 24, 25, 28, 30, 34, 35, 38, 40, 42, 43, 45, 46, 47, 49, 50, 52, 55, 56, 60, 61, 62, 63, 64, 65, 67, 68, 69\}$, $i_{17} = 2$, $\{9, 11, 23\}$ is covered by $47, 59, 61$. Thus $47 \cdot 59 \cdot 61 \mid a_9 a_{11} a_{23}$. Further $p \nmid a_i$ for $i \in \mathcal{B}_2$ and $p \in \mathcal{P}_2$. We now take $\mathcal{P}_0 = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \{43, 67\}$, $p_1 = 11, p_2 = 13$, $(i_1, i_2) := (4, 5)$, $\mathcal{I} = \mathcal{B}_2$, $\mathcal{P} = \mathcal{P}_3 := \Lambda(11, 13) \setminus \mathcal{P}_0 = \{5\}$ and $\ell = \ell_3 = \left\lceil \frac{k}{5} \right\rceil = 15$. Then $|\mathcal{I}'| = |\mathcal{B}_2| > 2\ell_3$. By Corollary 1, we get $\mathcal{M} =: \mathcal{M}_3$ and $\mathcal{B} =: \mathcal{B}_3$ such that $(\mathcal{M}_3, \mathcal{B}_3, \mathcal{P}_3, \ell_3)$ has *Property \mathfrak{H}* . We calculate $\mathcal{M}_3 = \{0, 10, 25, 30, 35, 40, 50, 55, 60, 65\}$, $\mathcal{B}_3 = \{1, 6, 7, 14, 16, 20, 21, 22, 24, 28, 34, 38, 42, 43, 45, 46, 47, 49, 52, 54, 56, 58, 61, 62, 63, 64, 66, 67, 68, 69\}$, $i_5 = 0$ and further $5 \nmid a_{20} a_{45}$. Lastly we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \mathcal{P}_3 \cup \{43, 67\}$, $p_1 = 5, p_2 = 11$, $(i_1, i_2) := (0, 4)$, $\mathcal{I} = \mathcal{B}_3$, $\mathcal{P} = \mathcal{P}_4 := \Lambda(5, 11) \setminus \mathcal{P}_0 = \{3, 23\}$ and $\ell = \ell_4 = \sum_{p \in \mathcal{P}_4} \left\lceil \frac{k}{p} \right\rceil$. By Lemma 5, we see that $M = \{16, 22, 24, 28, 43, 46, 47, 49, 64, 67\}$ is covered by \mathcal{P}_4 , $i_3 = i_{23} = 1$, $B = \{1, 6, 7, 14, 21, 34, 38, 42, 52, 56, 61, 62, 63, 68, 69\}$ and hence $3 \nmid a_7 a_{34} a_{52} a_{61}$ and possibly $3 \cdot 23|a_1$. Therefore $a_i \in \{1, 2, 7, 14\}$ for $i \in B \setminus \{1\}$. By taking $\mathcal{J} = B \setminus \{1\}$, we have $B \setminus \{1\} = \mathcal{I}_3^0 \cup \mathcal{I}_3^1 \cup \mathcal{I}_3^- = \mathcal{I}_5^+ \cup \mathcal{I}_5^-$ with

$$\mathcal{I}_3^0 = \{7, 34, 52, 61\}, \mathcal{I}_3^1 = \{6, 21, 42, 63, 69\}, \mathcal{I}_3^- = \{14, 38, 56, 62, 68\}$$

and

$$\mathcal{I}_5^+ = \{6, 14, 21, 34, 56, 61, 69\}, \quad \mathcal{I}_5^- = \{7, 38, 42, 52, 62, 63, 68\}.$$

Therefore

$$\mathcal{J}_1 = \{6, 21, 69\}, \quad \mathcal{J}_2 = \{42, 63\}, \quad \mathcal{J}_3 = \{14, 56\}, \quad \mathcal{J}_4 = \{38, 62, 68\}.$$

and hence $a_6 = a_{21} = a_{69} = 1, a_{42} = a_{63} = 7, a_{14} = a_{56} = 14, a_{38} = a_{62} = a_{68} = 2$ by (17). Further we get $a_{34} = a_{61} = 1$ and $a_{52} = 2$ by taking residue classes modulo 5. Since $\left(\frac{1}{71}\right) = \left(\frac{2}{71}\right) = 1$, we see that $\left(\frac{a_i}{71}\right) = 1$ for $i \in \{6, 21, 34, 38, 52, 61, 62, 68, 69\}$ which is not valid by the possibilities of i_{71} given by (22).

Let $k = 67$ and $(i_{43}, i_{67}) = (9, 9)$. We see that $\mathcal{P}_1 = \{11, 13, 19, 29, 31, 37, 41, 53\}$, $\mathcal{M}_1 = \{20, 22, 28, 31, 35, 38, 40, 42, 46, 47, 48, 50, 53, 61, 62, 64, 66\}$, $\mathcal{B}_1 = \{0, 1, 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21, 23, 24, 25, 26, 27, 29, 30, 32, 33, 34, 36, 37, 39, 41, 43, 44, 45, 49, 51, 54, 55, 56, 57, 58, 59, 60, 63, 65\}$, $i_{11} = i_{13} = i_{19} = 9$ and $\{38, 40, 46, 50, 62\}$ is covered by $29, 31, 37, 41, 53$. Further $p \nmid a_i$ for $i \in \mathcal{B}_1$ and $p \in \mathcal{P}_1$ except possibly when $29|a_{50}, 41|a_{62}, 29 \cdot 41|a_{21}$. Now we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \{43, 67\}$, $p_1 = 11, p_2 = 13$, $(i_1, i_2) := (9, 9)$, $\mathcal{I} = \mathcal{B}_1 \setminus \{21\}$ and $\mathcal{P} = \mathcal{P}_2 := \Lambda(11, 13) \setminus \mathcal{P}_0 = \{5, 17, 47, 59, 61\}$. If $5 \nmid d$, we observe that there is at least 1 multiple of 5 among $n + (i_{11} + 11i)d$, $0 \leq i \leq 5$ and $\ell \leq \sum_{p \in \mathcal{P}_2} \left\lfloor \frac{k}{p} \right\rfloor - 1 = 23$. Thus we always have $\ell \leq 23 = \ell_2$. Then $|\mathcal{I}'| = |\mathcal{B}_1| - 1 > 2\ell_2$ since $|\mathcal{B}_1| = 48$. Thus the conditions of Corollary 1 are satisfied and we get $\mathcal{M} =: \mathcal{M}_2$, $\mathcal{B} =: \mathcal{B}_2$ and $(\mathcal{M}_2, \mathcal{B}_2, \mathcal{P}_2, \ell_2)$ has *Property 5*. We have $\mathcal{M}_2 = \{0, 1, 2, 3, 5, 6, 7, 8, 14, 19, 24, 26, 29, 39, 43, 44, 49, 54, 56, 60\}$ which cannot be covered by \mathcal{P}_2 . This is a contradiction. The cases $k = 67, (i_{43}, i_{67}) = (i, i)$ with $9 \leq i \leq 28$ and $k = 71, (i_{43}, i_{67}) = (i, i)$ with $13 \leq i \leq 28, i \neq 24, 25$ are excluded similarly as in this paragraph. The remaining cases are excluded similarly as $k = 71, (i_{43}, i_{67}) = (27, 3)$ given in the preceding paragraph.

3.8. The cases $k = 73, 79$. We have $q_1 = 23, q_2 = 73$ and $\mathcal{P}_1 \subseteq \{13, 19, 29, 31, 37, 47, 59, 61, 67, 79\}$. Then the pairs (i_{q_1}, i_{q_2}) are given by

$$\begin{aligned} k = 73 &: (6, 2), (7, 3), (8, 4), (9, 5); \\ k = 79 &: (0, 0), (1, 1), (2, 2), (7, 3), (8, 4), (9, 5), (10, 6), (11, 7), (12, 8), \\ & (13, 9), (14, 10), (15, 11), (16, 12), (17, 13), (18, 14), (19, 15). \end{aligned}$$

These pairs are of the form $(i + 4, i)$ except for $(0, 0), (1, 1), (2, 2)$ in the case $k = 79$.

Let $k = 79$ and $(i_{23}, i_{73}) = (8, 4)$. We see that $\mathcal{P}_1 = \{13, 19, 29, 31, 37, 47, 59, 61, 67, 79\}$, $\mathcal{M}_1 = \{1, 3, 10, 12, 15, 16, 18, 19, 20, 25, 30, 38, 39, 40, 46, 48, 51, 58, 64, 78\}$, $\mathcal{B}_1 = \{0, 2, 5, 6, 7, 9, 11, 13, 14, 17, 21, 22, 23, 24, 26, 27, 28, 29, 32, 33, 34, 35, 36, 37, 41, 42, 43, 44, 45, 47, 49, 50, 52, 53, 55, 56, 57, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76\}$, $i_{13} = 12, i_{19} = 1$ and $\{3, 10, 15, 16, 18, 19, 30, 40, 46, 48, 78\}$ is covered by $29, 31, 37, 47, 59, 61, 67, 79$. Thus

$$29 \cdot 31 \cdot 37 \cdot 47 \cdot 59 \cdot 61 \cdot 67 \cdot 79 \mid \prod (n + id) \text{ for } i \in \{3, 10, 15, 16, 18, 19, 30, 40, 46, 48, 78\}.$$

Further we have

$$(23) \quad i_{79} \in \{10, 15, 16, 18, 19, 30, 40, 46, 48\}$$

and either $i_{29} = 19$ or $i_{29} \in \{1, 10, 16, 18\}$, $i_{31} = 15$, $i_{37} = 3$, $i_{59} = 19$. Also for $p \in \mathcal{P}_1$, we have $p \nmid a_i$ for $i \in \mathcal{B}_1$ since $i(\mathcal{P}_1)$ is even for $i \in \mathcal{B}_1$. For each possibility $i_{29} \in \{1, 10, 16, 18, 19\}$, we now take $\mathcal{P}_0 = \mathcal{P}_1 \cup \{23, 73\}$, $p_1 = 19$, $p_2 = 29$, $(i_1, i_2) := (1, i_{29})$, $\mathcal{I} = \mathcal{B}_1$, $\mathcal{P} = \mathcal{P}_2 := \Lambda(19, 29) \setminus \mathcal{P}_0 = \{11, 17, 43, 53, 71\}$ and $\ell = \ell_2 = \sum_{p \in \mathcal{P}_2} \left\lceil \frac{k}{p} \right\rceil = 19$. Then $|\mathcal{I}'| \geq |\mathcal{B}_1| - 2 > 2\ell_2$. Thus the conditions of Corollary 1 are satisfied and we have $\mathcal{M} =: \mathcal{M}_2$, $\mathcal{B} =: \mathcal{B}_2$ and $(\mathcal{M}_2, \mathcal{B}_2, \mathcal{P}_2, \ell_2)$ has *Property* \mathfrak{H} implying $i_{29} = 19$ in which case we get $\mathcal{M}_2 = \{0, 6, 9, 11, 22, 24, 26, 33, 34, 43, 44, 55, 60, 66\}$, $\mathcal{B}_2 = \{2, 5, 7, 13, 14, 17, 21, 23, 27, 28, 29, 32, 35, 36, 37, 41, 42, 45, 47, 49, 50, 52, 53, 56, 57, 59, 61, 62, 63, 65, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76\}$, $i_{11} = 0$, $i_{17} = 9$, $\{6, 24, 34\}$ is covered by $43, 53, 71$. Thus $43 \cdot 53 \cdot 71 \mid a_6 a_{24} a_{34}$. Further $p \nmid a_i$ for $i \in \mathcal{B}_2$ and $p \in \mathcal{P}_2$. We now take $\mathcal{P}_0 = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \{23, 73\}$, $p_1 = 11$, $p_2 = 13$, $(i_1, i_2) := (0, 12)$, $\mathcal{I} = \mathcal{B}_2$, $\mathcal{P} = \mathcal{P}_3 := \Lambda(11, 13) \setminus \mathcal{P}_0 = \{5\}$ and $\ell = \ell_3 = \left\lceil \frac{k}{5} \right\rceil = 16$. Then $|\mathcal{I}'| = |\mathcal{B}_2| > 2\ell_3$. By Corollary 1, we get $\mathcal{M} =: \mathcal{M}_3$ and $\mathcal{B} =: \mathcal{B}_3$ with $(\mathcal{M}_3, \mathcal{B}_3, \mathcal{P}_3, \ell_3)$ having *Property* \mathfrak{H} . We calculate $\mathcal{M}_3 = \{7, 17, 32, 37, 42, 47, 57, 62, 67, 72\}$, $\mathcal{B}_3 = \{2, 5, 13, 14, 21, 23, 27, 28, 29, 35, 36, 41, 45, 49, 50, 52, 53, 56, 59, 61, 63, 65, 68, 69, 70, 71, 73, 74, 75, 76\}$, $i_5 = 2$ and $5 \nmid a_i$ for $i \in \mathcal{B}_3$. Lastly we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \mathcal{P}_3 \cup \{23, 73\}$, $p_1 = 5$, $p_2 = 11$, $(i_1, i_2) := (2, 0)$, $\mathcal{I} = \mathcal{B}_3$, $\mathcal{P} = \mathcal{P}_4 := \Lambda(5, 11) \setminus \mathcal{P}_0 = \{3, 41\}$ and $\ell = \ell_4 = \sum_{p \in \mathcal{P}_4} \left\lceil \frac{k}{p} \right\rceil$. By Lemma 5, we see that $M = \{23, 29, 35, 36, 50, 53, 56, 65, 71, 74\}$ is covered by \mathcal{P}_4 , $i_3 = 2$, $i_{41} = 36$, $B = \{5, 13, 14, 21, 28, 41, 45, 49, 59, 61, 63, 68, 69, 70, 73, 75, 76\}$ and hence $a_i \in \{1, 2, 7, 14\}$ for $i \in B$. By taking $\mathcal{J} = B$, we have $B = \mathcal{I}_3^0 \cup \mathcal{I}_3^1 \cup \mathcal{I}_3^2 = \mathcal{I}_5^+ \cup \mathcal{I}_5^-$ with

$$\mathcal{I}_3^0 = \{5, 14, 41, 59, 68\}, \quad \mathcal{I}_3^1 = \{13, 28, 49, 61, 70, 73, 76\}, \quad \mathcal{I}_3^2 = \{21, 45, 63, 69, 75\}$$

and

$$\mathcal{I}_5^+ = \{13, 21, 28, 41, 61, 63, 68, 73, 76\}, \quad \mathcal{I}_5^- = \{5, 14, 45, 49, 59, 69, 70, 75\}.$$

Thus

$$\mathcal{J}_1 = \{13, 28, 61, 73, 76\}, \quad \mathcal{J}_2 = \{49, 70\}, \quad \mathcal{J}_3 = \{21, 63\}, \quad \mathcal{J}_4 = \{45, 69, 75\}.$$

and hence $a_{13} = a_{28} = a_{61} = a_{73} = a_{76} = 1$, $a_{49} = a_{70} = 7$, $a_{21} = a_{63} = 14$, $a_{45} = a_{69} = a_{75} = 2$ by (17). Further we get $a_{41} = a_{68} = 1$ and $a_5 = a_{59} = 2$ by residue modulo 5. Since $\left(\frac{1}{79}\right) = \left(\frac{2}{79}\right) = 1$, we see that $\left(\frac{a_i}{71}\right) = 1$ for $i \in \{5, 13, 28, 41, 45, 59, 61, 68, 69, 75, 76\}$ which is not valid by the possibilities of i_{79} given by (23). The other cases are excluded similarly.

3.9. The case $k = 83$. We have $q_1 = 37$, $q_2 = 83$ and $\mathcal{P}_1 = \{17, 23, 29, 31, 47, 53, 59, 61, 67, 71, 73\}$. Then the pairs (i_{q_1}, i_{q_2}) are given by

$$(13, 4), (14, 5), (15, 6), (16, 7), (17, 8), (18, 9), (19, 10), \\ (20, 11), (21, 12), (22, 13), (23, 14), (24, 15), (25, 16), (26, 17).$$

These pairs are of the form $(i + 9, i)$ with $4 \leq i \leq 17$.

Let $(i_{37}, i_{83}) = (13, 4)$. We see that $\mathcal{P}_1 = \{17, 23, 29, 31, 47, 53, 59, 61, 67, 71, 73\}$, $\mathcal{M}_1 = \{0, 2, 14, 16, 18, 19, 20, 25, 26, 28, 29, 34, 36, 40, 41, 53, 56, 58, 64, 70\}$, $\mathcal{B}_1 = \{1, 3, 5, 6, 7, 8, 9, 10, 11, 12, 15, 17, 21, 22, 23, 24, 27, 30, 31, 32, 33, 35, 37, 38, 39, 42, 43, 44, 45, 46, 47, 48, 49, 51, 52, 54, 55, 57, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82\}$, $i_{17} = 2$, $i_{23} = 18$, $i_{29} = 0$, $i_{31} = 25$ and $\{14, 16, 20, 26, 28, 34, 40\}$

is covered by 47, 53, 59, 61, 67, 71, 73. Further $p \nmid a_i$ for $i \in \mathcal{B}_1$ and $p \in \mathcal{P}_1$. For each possibility $i_{73} \in \{14, 16, 20, 26, 28, 34, 40\}$, we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \{37, 83\}$, $p_1 = 23, p_2 = 73$, $(i_1, i_2) := (18, i_{73})$, $\mathcal{I} = \mathcal{B}_1$, $\mathcal{P} = \mathcal{P}_2 := \Lambda(23, 73) \setminus \mathcal{P}_0 = \{13, 19, 79\}$ and $\ell = \ell_2 = \sum_{p \in \mathcal{P}_2} \left\lceil \frac{k}{p} \right\rceil = 14$. Then $|\mathcal{I}'| = |\mathcal{B}_1| > 2\ell_2$. Thus the conditions of Corollary 1 are satisfied and we get $\mathcal{M} =: \mathcal{M}_2$, $\mathcal{B} =: \mathcal{B}_2$ and $(\mathcal{M}_2, \mathcal{B}_2, \mathcal{P}_2, \ell_2)$ has *Property* \mathfrak{H} which is possible only if $i_{73} = 14$. Then $\mathcal{M}_2 = \{8, 9, 11, 22, 30, 35, 48, 49, 61, 68, 74\}$. Therefore $i_{13} = 9, i_{19} = 11$ and $i_{79} = 8$. This is not possible by applying the case $k = 73$ to $(n + 9d) \cdots (n + 81d)$. Similarly for $(i_{37}, i_{83}) = (14, 5)$, we get $i_{73} = 15, i_{79} = 9$ and this is excluded by applying the case $k = 73$ to $(n + 10d) \cdots (n + 82d)$. For all the remaining cases, we continue similarly to find that \mathcal{M}_2 is not covered by \mathcal{P}_2 for possible choices of i_{73} and hence they are excluded.

3.10. The case $k = 89$. We have $q_1 = 79, q_2 = 89$ and $\mathcal{P}_1 = \{13, 17, 19, 23, 31, 47, 53, 71, 83\}$. Then the pairs (i_{q_1}, i_{q_2}) are given by $(16, 6), (17, 7), (18, 8), (19, 9), (20, 10), (21, 11)$. These pairs are of the form $(i + 10, i)$ with $6 \leq i \leq 11$.

Let $(i_{79}, i_{89}) = (16, 6)$. We see that $\mathcal{P}_1 = \{13, 17, 19, 23, 31, 47, 53, 71, 83\}$, $\mathcal{M}_1 = \{0, 1, 2, 3, 4, 10, 12, 17, 19, 24, 26, 27, 30, 33, 38, 42, 43, 44, 48, 49, 56, 57, 61, 64, 69, 72, 76, 78, 82\}$, $\mathcal{B}_1 = \{5, 7, 8, 9, 11, 13, 14, 15, 18, 20, 21, 22, 23, 25, 28, 29, 31, 32, 34, 35, 36, 37, 39, 40, 41, 45, 46, 47, 50, 51, 52, 53, 54, 55, 58, 59, 60, 62, 63, 65, 66, 67, 68, 70, 71, 73, 74, 75, 77, 79, 80, 81, 83, 84, 85, 86, 87, 88\}$, $i_{13} = 4, i_{17} = 10, i_{19} = 0, i_{23} = 3, i_{31} = 2, i_{47} = 1$ and $\{12, 24, 42\}$ is covered by 53, 71, 83. Further $p \nmid a_i$ for $i \in \mathcal{B}_1$ and $p \in \mathcal{P}_1$. Now we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \{79, 89\}$, $p_1 = 31, p_2 = 89$, $(i_1, i_2) := (2, 6)$, $\mathcal{I} = \mathcal{B}_1$ and $\mathcal{P} = \mathcal{P}_2 := \Lambda(31, 89) \setminus \mathcal{P}_0 = \{7, 11, 41, 59, 73\}$. If $7 \nmid d$, we observe that there is at least 1 multiple of 7 among $n + (i_{13} + 13i)d$, $0 \leq i \leq 6$ and $\ell \leq \ell_2 = \sum_{p \in \mathcal{P}_2} \left\lceil \frac{k}{p} \right\rceil - 1 = 28$. Thus in all cases, we have $\ell \leq \ell_2$ and $|\mathcal{I}'| = |\mathcal{B}_1| > 2\ell_2$. Therefore the conditions of Corollary 1 are satisfied and we get $\mathcal{M} =: \mathcal{M}_2$ and $\mathcal{B} =: \mathcal{B}_2$ with $(\mathcal{M}_2, \mathcal{B}_2, \mathcal{P}_2, \ell_2)$ having *Property* \mathfrak{H} . We find $\mathcal{M}_2 = \{7, 11, 13, 22, 25, 29, 32, 36, 39, 40, 51, 53, 54, 60, 62, 67, 73, 74, 81, 84, 88\}$, $\mathcal{B}_2 = \{5, 8, 9, 14, 15, 18, 20, 21, 23, 28, 31, 34, 35, 37, 41, 45, 46, 47, 50, 52, 55, 58, 59, 63, 65, 66, 68, 70, 71, 75, 77, 79, 80, 83, 85, 86, 87\}$, $i_7 = 4, i_{11} = 7, i_{41} = 13$ and $\{22, 36\}$ is covered by 59, 73. Further for $p \in \mathcal{P}_2$, $p \nmid a_i$ for $i \in \mathcal{B}_2 \setminus \{18\}$. We take $\mathcal{P}_0 = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \{79, 89\}$, $p_1 = 41, p_2 = 79$, $(i_1, i_2) := (13, 16)$, $\mathcal{I} = \mathcal{B}_2 \setminus \{18\}$, $\mathcal{P} = \mathcal{P}_3 := \Lambda(41, 79) \setminus \mathcal{P}_0 = \{37, 43, 61, 67\}$ and $\ell = \ell_3 = \sum_{p \in \mathcal{P}_3} \left\lceil \frac{k}{p} \right\rceil = 10$. Then $|\mathcal{I}'| = |\mathcal{I}| = |\mathcal{B}_2| - 1 > 2\ell_3$. Thus the conditions of Corollary 1 are satisfied and we have $\mathcal{M} =: \mathcal{M}_3$, $\mathcal{B} =: \mathcal{B}_3$ and $(\mathcal{M}_3, \mathcal{B}_3, \mathcal{P}_3, \ell_3)$ has *Property* \mathfrak{H} . We get $\mathcal{M}_3 = \{9, 21, 28, 34, 52, 58\}$, $\mathcal{B}_3 = \{5, 8, 14, 15, 20, 23, 31, 35, 37, 41, 45, 46, 47, 50, 55, 59, 63, 65, 66, 68, 70, 71, 75, 77, 79, 80, 83, 85, 86, 87\}$, $i_{37} = 21, i_{43} = 9$ and $\{28, 34\}$ is covered by 61, 67. Therefore $p \in \{2, 3, 5, 29\}$ whenever $p|a_i$ for $i \in \mathcal{B}_3$. Now we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \mathcal{P}_3 \cup \{79, 89\}$, $p_1 = 7, p_2 = 17$, $(i_1, i_2) := (4, 10)$, $\mathcal{I} = \mathcal{B}_3$, $\mathcal{P} = \mathcal{P}_4 := \Lambda(7, 17) \setminus \mathcal{P}_0 = \{29\}$ and $\ell = \ell_4 = \left\lceil \frac{k}{29} \right\rceil = 4$. Then $|\mathcal{I}'| = |\mathcal{B}_3| - 1$ since $46 \in \mathcal{B}_3$ and $|\mathcal{B}_3| - 1 > 2\ell_4$. By Corollary 1, we get $\mathcal{M} =: \mathcal{M}_4$ and $\mathcal{B} =: \mathcal{B}_4$ with $(\mathcal{M}_4, \mathcal{B}_4, \mathcal{P}_4, \ell_4)$ having *Property* \mathfrak{H} . We find $\mathcal{M}_4 = \{8, 37, 66\}$, $\mathcal{B}_4 = \{5, 14, 15, 20, 23, 31, 35, 41, 45, 47, 50, 55, 59, 63, 65, 68, 70, 71, 75, 77, 79, 80, 83, 85, 86, 87\}$, $i_{29} = 8$ and $P(a_i) \leq 5$ for $i \in \mathcal{B}_4$. Now we get a contradiction by taking $k = 6$ and $(n + 47d)(n + 55d)(n + 63d)(n + 71d)(n + 79d)(n + 87d) = b'y'^2$. Similarly the pair $(i_{79}, i_{89}) = (17, 7)$ is excluded by applying $k = 6$ to $(n + 48d)(n +$

$56d)(n + 64d)(n + 72d)(n + 80d)(n + 88d)$. For all the remaining cases, we continue similarly to find that \mathcal{M}_3 is not covered by \mathcal{P}_3 and hence they are excluded.

4. PROOF OF LEMMA 7

Assume that $Q_1 \nmid d$ and $Q_2 \nmid d$. Then, by taking mirror image (4) of (2), there is no loss of generality in assuming that $0 \leq i_{Q_1} < Q_1, 0 \leq i_{Q_2} \leq \min(Q_2 - 1, \frac{k-1}{2})$. Further $i_{Q_2} \geq k - k'$ if $Q_2 = k$. Let $\mathcal{P}_0 = \{Q_0\}, p_1 = Q_1, p_2 = Q_2, (i_1, i_2) := (i_{Q_1}, i_{Q_2}), \mathcal{I} = [0, k) \cap \mathbb{Z}$ and $\mathcal{P} = \mathcal{P}_1 := \Lambda(Q_1, Q_2) \setminus \mathcal{P}_0$. Then $|\mathcal{I}'| \geq k - \lceil \frac{k}{Q_1} \rceil - \lceil \frac{k}{Q_2} \rceil$ and $\ell \leq \ell_1$ where $\ell_1 = \sum_{p \in \mathcal{P}_1} \lceil \frac{k}{p} \rceil$. In fact we can take $\ell_1 = \sum_{p \in \mathcal{P}_1} \lceil \frac{k}{p} \rceil - 1$ if $(k, Q_0) = (79, 23)$ or $(k, Q_0) = (59, 29)$ with $i_7 \leq 2$ by considering multiples of 13, 11 or 19, 7, 11, respectively.

Let $(k, Q_0) \neq (79, 73)$. Then $\ell_1 < \frac{1}{2}|\mathcal{I}'|$. We observe that $i(\mathcal{P}_0) = 0$ for $i \in \mathcal{I}'$ since $Q_0 \mid d$ and by Corollary 1, we get $\mathcal{M} =: \mathcal{M}_1, \mathcal{B} =: \mathcal{B}_1$ and $(\mathcal{M}_1, \mathcal{B}_1, \mathcal{P}_1, \ell_1)$ has *Property* \mathfrak{H} . We now restrict to all such pairs (i_{Q_1}, i_{Q_2}) with $|\mathcal{M}_1| \leq \ell_1$ and \mathcal{M}_1 is covered by \mathcal{P}_1 . These pairs are given by

k	Q_0	(Q_1, Q_2)	(i_{Q_1}, i_{Q_2})
29	19	(7, 17)	(0, 0), (0, 11)
37	19 or 29	(7, 17)	(0, 0), (1, 2)
47	29	(7, 17)	(0, 0), (4, 12)
59	29	(7, 17)	(1, 1), (1, 6)
71	43	(53, 67)	(0, 0)
89	79	(23, 73)	(0, 0), (19, 15)

Let $(k, Q_0) = (79, 73)$ and $(Q_1, Q_2) = (53, 67)$. We apply Lemma 5 to derive that either $|\mathcal{I}_1| \leq \ell_1, \mathcal{I}_1$ is covered by $\mathcal{P}_1, i(\mathcal{P}_1)$ is even for $i \in \mathcal{I}_2$ or $|\mathcal{I}_2| \leq \ell_1, \mathcal{I}_2$ is covered by $\mathcal{P}_1, i(\mathcal{P}_1)$ is even for $i \in \mathcal{I}_1$. We compute $\mathcal{I}_1, \mathcal{I}_2$ and we find that both \mathcal{I}_1 and \mathcal{I}_2 are not covered by \mathcal{P}_1 for each pair (i_{53}, i_{67}) with $0 \leq i_{53} < 53, 0 \leq i_{67} \leq \frac{k-1}{2}$.

Let $(k, Q_0) = (37, 29), (Q_1, Q_2) = (7, 17)$ and $(i_7, i_{17}) = (1, 2)$. Then $\mathcal{P}_1 = \{11, 13, 19, 23, 37\}$. We find that $\mathcal{M}_1 = \{3, 7, 10, 13, 14, 17, 23, 25\}, \mathcal{B}_1 = \{0, 4, 5, 6, 9, 11, 12, 16, 18, 20, 21, 24, 26, 27, 28, 30, 31, 32, 33, 34, 35\}, i_{11} = 3, i_{13} = 10$ and $\{7, 13, 17\}$ is covered by 19, 23, 37. Further $p \nmid a_i$ for $p \in \mathcal{P}_1, i \in \mathcal{B}_1$. Now we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \{7, 17, 29\}, p_1 = 11, p_2 = 13, (i_1, i_2) := (3, 10), \mathcal{I} = \mathcal{B}_1, \mathcal{P} = \mathcal{P}_2 := \Lambda(11, 13) \setminus \mathcal{P}_0 = \{5, 31\}$ and $\ell = \ell_2 = \sum_{p \in \mathcal{P}_2} \lceil \frac{k}{p} \rceil = 10$. Thus $|\mathcal{I}'| = |\mathcal{I}| = |\mathcal{B}_1| = 21 > 2\ell_2$. Then the conditions of Corollary 1 are satisfied and we have $\mathcal{M} =: \mathcal{M}_2, \mathcal{B} =: \mathcal{B}_2$ and $(\mathcal{M}_2, \mathcal{B}_2, \mathcal{P}_2, \ell_2)$ has *Property* \mathfrak{H} . We get $\mathcal{M}_2 = \{5, 6, 16, 21, 26, 31\}, \mathcal{B}_2 = \{0, 4, 9, 11, 12, 18, 20, 24, 27, 28, 30, 32, 33, 34, 35\}, i_5 = 1, 31 \mid a_5$ and $5 \nmid a_{11}$. Also $P(a_i) \leq 3$ for $i \in \mathcal{B}_2$ and $P(a_{31}) = 5$. Thus $P(a_{30}a_{31} \cdots a_{35}) \leq 5$ and this is excluded by the case $k = 6$. The other cases for $k = 29, 37, 47$ are excluded similarly. Each possibility is excluded by the case $k = 6$ after showing $P(a_1a_2 \cdots a_6) \leq 5$ when $(k, Q_0) \in \{(29, 19), (37, 19), (37, 29), (47, 29)\}, (i_7, i_{17}) = (0, 0); P(a_{22}a_{23} \cdots a_{27}) \leq 5$ when $(k, Q_0) = (29, 19), (i_7, i_{17}) = (0, 11); P(a_{30}a_{31} \cdots a_{35}) \leq 5$ when $(k, Q_0) =$

$(37, 19)$, $(i_7, i_{17}) = (1, 2)$ and $P(a_{40}a_{41} \cdots a_{45}) \leq 5$ when $(k, Q_0) = (47, 29)$, $(i_7, i_{17}) = (4, 12)$.

Let $(k, Q_0) = (59, 29)$, $(Q_1, Q_2) = (7, 17)$ and $(i_7, i_{17}) = (1, 1)$. Then $\mathcal{P}_1 = \{11, 13, 19, 23, 37, 47, 59\}$. We find that $\mathcal{M}_1 = \{0, 12, 14, 20, 23, 24, 27, 30, 34, 38, 39, 40, 45, 47, 48, 53, 56, 58\}$, $\mathcal{B}_1 = \{2, 3, 4, 5, 6, 7, 9, 10, 11, 13, 16, 17, 19, 21, 25, 26, 28, 31, 32, 33, 37, 41, 42, 44, 46, 49, 51, 54, 55\}$, $i_{11} = i_{13} = i_{19} = i_{23} = 1$, $\{30, 38, 48\}$ is covered by $37, 47, 59$. Further $p \nmid a_i$ for $p \in \mathcal{P}_1$, $i \in \mathcal{B}_1$. Now we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \{7, 17, 29\}$, $p_1 = 11, p_2 = 13$, $(i_1, i_2) := (1, 1)$, $\mathcal{I} = \mathcal{B}_1$, $\mathcal{P} = \mathcal{P}_2 := \Lambda(11, 13) \setminus \mathcal{P}_0 = \{5, 31, 43\}$ and $\ell = \ell_2 = \sum_{p \in \mathcal{P}_2} \left[\frac{k}{p} \right]$. By Lemma 5, we get $M = \{6, 11, 16, 21, 31, 32, 41, 44, 46\}$, $i_5 = 1$, $31 \cdot 43 | a_{32} a_{44}$ and $i(\mathcal{P}_2)$ is even for $i \in B = \{2, 3, 4, 5, 7, 9, 10, 13, 17, 19, 25, 26, 28, 33, 37, 42, 49, 51, 54, 55\}$. Further for $p \in \mathcal{P}_2$, $p \nmid a_i$ for $i \in B$. Finally we apply Lemma 5 with $\mathcal{P}_0 = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \{7, 17, 29\}$, $p_1 = 5, p_2 = 11$, $(i_1, i_2) := (1, 1)$, $\mathcal{I} = B$ and $\mathcal{P} = \mathcal{P}_3 := \Lambda(5, 11) \setminus \mathcal{P}_0 = \{3, 41, 53\}$. We get $M_1 = \{4, 7, 13, 25, 28, 42, 49, 54, 55\}$ which is covered by \mathcal{P}_3 , $i_3 = 1$, $\{42, 54\}$ is covered by $\{41, 53\}$ and $i(\mathcal{P}_3)$ is even for $i \in B_1 = \{2, 3, 5, 9, 10, 17, 19, 33, 37\}$. Hence $P(a_i) \leq 2$ for $i \in B_1$. Since $\left(\frac{a_i}{29}\right) = \left(\frac{n}{29}\right)$ and $\left(\frac{2}{29}\right) \neq 1$, we see that $a_i = 1$ for $i \in B_1$. By taking $\mathcal{J} = B_1$, we derive that either $\mathcal{I}_5^+ = \emptyset$ or $\mathcal{I}_5^- = \emptyset$ which is a contradiction. The other case $(i_7, i_{17}) = (1, 6)$ is excluded similarly.

Let $(k, Q_0) = (71, 43)$, $(Q_1, Q_2) = (53, 67)$, $(i_{53}, i_{67}) = (0, 0)$. Then $\mathcal{P}_1 = \{7, 11, 13, 19, 23, 71\}$. We get $\mathcal{M}_1 = \{7, 11, 13, 14, 19, 21, 22, 23, 26, 28, 33, 35, 38, 39, 42, 43, 44, 46, 52, 55, 56, 57, 63, 65, 66, 69, 70\}$, $\mathcal{B}_1 = \{1, 2, 3, 4, 5, 6, 8, 9, 10, 12, 15, 16, 17, 18, 20, 24, 25, 27, 29, 30, 31, 32, 34, 36, 37, 40, 41, 45, 47, 48, 49, 50, 51, 54, 58, 59, 60, 61, 62, 64, 68\}$, $i_7 = i_{11} = i_{13} = i_{19} = i_{23} = 0$, $i_{71} = 43$. Further, for $p \in \mathcal{P}_1$, $p \nmid a_i$ for $i \in \mathcal{B}_1$. Now we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \{43, 53, 67\}$, $p_1 = 11, p_2 = 13$, $(i_1, i_2) := (0, 0)$, $\mathcal{I} = \mathcal{B}_1$, $\mathcal{P} = \mathcal{P}_2 := \Lambda(11, 13) \setminus \mathcal{P}_0 = \{5, 17, 29, 31, 37, 47, 59, 61\}$ and $\ell = \ell_2 = \sum_{p \in \mathcal{P}_2} \left[\frac{k}{p} \right]$. By Lemma 5, we see that $M = \{5, 10, 15, 17, 20, 29, 30, 31, 34, 37, 40, 45, 47, 51, 58, 59, 60, 61, 62, 68\}$ is covered by \mathcal{P}_2 , $i(\mathcal{P}_2)$ is even for $i \in B = \{1, 2, 3, 4, 6, 8, 9, 12, 16, 18, 24, 25, 27, 32, 36, 41, 48, 49, 50, 54, 64\}$. We get $i_5 = i_{17} = i_{29} = i_{31} = 0$, and $\{37, 47, 59, 61\}$ is covered by $37, 47, 59, 61$. Thus $37 \cdot 47 \cdot 59 \cdot 61 | a_{37} a_{47} a_{59} a_{61}$. Further $p \nmid a_i$ for $i \in B$ and $p \in \mathcal{P}_2$. We take $\mathcal{P}_0 = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \{43, 53, 67\}$, $p_1 = 5, p_2 = 11$, $(i_1, i_2) := (0, 0)$, $\mathcal{I} = \mathcal{B}_2$, $\mathcal{P} = \mathcal{P}_3 := \Lambda(5, 11) \setminus \mathcal{P}_0 = \{3, 41\}$ and $\ell = \ell_3 = \sum_{p \in \mathcal{P}_3} \left[\frac{k}{p} \right]$. By Lemma 5, we see that $M_1 = \{3, 6, 12, 24, 27, 41, 48, 54\}$ is covered by \mathcal{P}_3 , $i(\mathcal{P}_3)$ is even for $i \in B_1 = \{1, 2, 4, 8, 9, 16, 18, 32, 36, 49, 64\}$. Thus $i_3 = 0$ implying $i_{41} = 0$ and $p = 2$ whenever $p | a_i$ for $i \in B_1$. By taking $\mathcal{J} = B_1$, we have $B_1 = \mathcal{I}_5^+ \cup \mathcal{I}_5^-$ with

$$\mathcal{I}_5^+ = \{1, 4, 9, 16, 36, 49, 64\}, \quad \mathcal{I}_5^- = \{2, 8, 18, 32\}.$$

Thus $a_i = 1$ for $i \in \mathcal{I}_5^+$ and $a_i = 2$ for $i \in \mathcal{I}_5^-$ since $a_i \in \{1, 2\}$ for $i \in B_1$. This is a contradiction since $43 | d$, $\left(\frac{a_i}{43}\right) = \left(\frac{n}{43}\right)$ and $\left(\frac{1}{43}\right) \neq \left(\frac{2}{43}\right)$.

Let $k = 89$, $Q_0 = 79$, $(Q_1, Q_2) = (23, 73)$, $(i_{23}, i_{73}) = (19, 15)$. Then $\mathcal{P}_1 = \{13, 19, 29, 31, 37, 47, 59, 61, 67, 79, 89\}$. We find that $\mathcal{M}_1 = \{1, 9, 10, 12, 14, 21, 23, 26, 27, 29, 30, 31, 36, 41, 49, 50, 51, 57, 59, 62, 69, 75\}$, $\mathcal{B}_1 = \{0, 2, 3, 4, 5, 6, 7, 8, 11, 13, 16, 17, 18, 20, 22, 24, 25, 28, 32, 33, 34, 35, 37, 38, 39, 40, 43, 44, 45, 46, 47, 48, 52, 53, 54, 55, 56, 58, 60, 61, 63, 64, 66, 67, 68, 70, 71, 72, 73, 74, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87\}$, $i_{13} =$

10, $i_{19} = 12, i_{29} = 1, i_{31} = 26, i_{37} = 14$ and $\{9, 21, 27, 29, 41\}$ is covered by 47, 59, 61, 67, 89. Thus $i_{89} \in \{9, 21, 27, 29, 41\}$. Further for $p \in \mathcal{P}_1$, $p \nmid a_i$ for $i \in \mathcal{B}_1$. Now we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \{23, 73, 79\}$, $p_1 = 19, p_2 = 29$, $(i_1, i_2) := (12, 1), \mathcal{I} = \mathcal{B}_1$, $\mathcal{P} = \mathcal{P}_2 := \Lambda(19, 29) \setminus \mathcal{P}_0 = \{11, 17, 43, 53, 71\}$ and $\ell = \ell_2 = \sum_{p \in \mathcal{P}_2} \left\lfloor \frac{k}{p} \right\rfloor = 22$. Thus $|\mathcal{I}'| = |\mathcal{I}| = |\mathcal{B}_1| > 2\ell_2$. By Corollary 1, we have $\mathcal{M} =: \mathcal{M}_2$, $\mathcal{B} =: \mathcal{B}_2$ and $(\mathcal{M}_2, \mathcal{B}_2, \mathcal{P}_2, \ell_2)$ has *Property* \mathfrak{H} . We get $\mathcal{M}_2 = \{0, 2, 3, 11, 17, 20, 22, 33, 35, 37, 44, 45, 54, 55, 66, 71, 77\}$, $\mathcal{B}_2 = \{4, 5, 6, 7, 8, 13, 16, 18, 24, 25, 28, 32, 34, 38, 39, 40, 43, 46, 47, 48, 52, 53, 56, 58, 60, 61, 63, 64, 67, 68, 70, 72, 73, 74, 76, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87\}$, $i_{11} = 0, i_{17} = 3, i_{43} = 2$ and $\{17, 35\}$ is covered by 53, 71. Further $p \nmid a_i$ for $i \in \mathcal{B}_2$ and $p \in \mathcal{P}_2$. We take $\mathcal{P}_0 = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \{23, 73, 79\}$, $p_1 = 11, p_2 = 13$, $(i_1, i_2) := (0, 10), \mathcal{I} = \mathcal{B}_2$, $\mathcal{P} = \mathcal{P}_3 := \Lambda(11, 13) \setminus \mathcal{P}_0 = \{5\}$ and $\ell = \ell_3 = \sum_{p \in \mathcal{P}_3} \left\lfloor \frac{k}{p} \right\rfloor = 18$. Thus $|\mathcal{I}'| = |\mathcal{I}| = |\mathcal{B}_2| > 2\ell_3$. Then the conditions of Corollary 1 are satisfied and we have $\mathcal{M} =: \mathcal{M}_3$, $\mathcal{B} =: \mathcal{B}_3$ with $(\mathcal{M}_3, \mathcal{B}_3, \mathcal{P}_3, \ell_3)$ having *Property* \mathfrak{H} . We get $\mathcal{M}_3 = \{8, 18, 28, 43, 48, 53, 58, 68, 73, 78, 83\}$, $\mathcal{B}_3 = \{4, 5, 6, 7, 13, 16, 24, 25, 32, 34, 38, 39, 40, 46, 47, 52, 56, 60, 61, 63, 64, 67, 70, 72, 74, 76, 79, 80, 81, 82, 84, 85, 86, 87\}$, $i_5 = 3$. Lastly we take $\mathcal{P}_0 = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \mathcal{P}_3 \cup \{23, 73, 79\}$, $p_1 = 5, p_2 = 11$, $(i_1, i_2) := (3, 0), \mathcal{I} = \mathcal{B}_3$, $\mathcal{P} = \mathcal{P}_4 := \Lambda(5, 11) \setminus \mathcal{P}_0 = \{3, 41\}$ and $\ell = \ell_4 = \sum_{p \in \mathcal{P}_4} \left\lfloor \frac{k}{p} \right\rfloor$. By Lemma 5, we see that $M = \{4, 6, 34, 40, 46, 47, 61, 64, 67, 76, 82, 85\}$ is covered by \mathcal{P}_4 , $\mathbf{i}(\mathcal{P}_4)$ is even for $i \in B = \{5, 7, 16, 24, 25, 32, 39, 52, 56, 60, 70, 72, 74, 79, 80, 81, 84, 86, 87\}$. Thus $i_3 = 1, i_{41} = 6$ and $p \in \{2, 7, 83\}$ whenever $p|a_i$ for $i \in B$. Since $79|d$, we see that $a_i \in \{1, 2, 83, 2 \cdot 83\}$ or $a_i \in \{7, 14, 7 \cdot 83, 14 \cdot 83\}$ for $i \in B$. The latter possibility is excluded since $7 \nmid (i - i')$ for all $i, i' \in B$. By taking $\mathcal{J} = B$, we have $B = \mathcal{I}_5^+ \cup \mathcal{I}_5^-$ with

$$\mathcal{I}_5^+ = \{7, 24, 32, 39, 52, 72, 74, 79, 84, 87\}, \quad \mathcal{I}_5^- = \{5, 16, 25, 56, 60, 70, 80, 81, 86\}.$$

Then we observe that either $a_i \in \{1, 2 \cdot 83\}$ for $i \in \mathcal{I}_5^+$ and $a_i \in \{2, 83\}$ for $i \in \mathcal{I}_5^-$ or vice-versa. This is not possible by parity argument. The other case $(i_{23}, i_{73}) = (0, 0)$ is excluded similarly.

5. PROOF OF LEMMA 8

Let $7 \leq k \leq 97$ be primes. Suppose that the assumptions of Lemma 8 are satisfied. Assume that $q_1|d$ or $q_2|d$ and we shall arrive at a contradiction. We divide the proof in subsections 5.1 and 5.2

5.1. The cases $7 \leq k \leq 23$. We take $q = 5$ in (7) and (8). We may suppose that $5|d$ if $k = 7, 11$ and $11|d$ if $k = 13$. Let $5|d$. Then

$$(24) \quad S \subseteq \{1, 6\} \text{ or } S \subseteq \{2, 3\}$$

according as $\left(\frac{n}{5}\right) = 1$ or -1 , respectively. Thus (24) holds if $k = 7, 11$. Let $11|d$. Then

$$(25) \quad S \subseteq \{1, 3, 5, 15\} \text{ or } S \subseteq \{2, 6, 10, 30\}$$

according as $\left(\frac{n}{11}\right) = 1$ or -1 , respectively. Let $13|d$. Then

$$(26) \quad S \subseteq \{1, 3, 10, 30\} \text{ or } S \subseteq \{2, 5, 6, 15\}$$

according as $\left(\frac{n}{13}\right) = 1$ or -1 , respectively. Thus either (25) or (26) holds if $13 \leq k \leq 23$.

By observing that a_i 's divisible by a prime p can occur in at most $\left\lceil \frac{k}{p} \right\rceil$ terms, we have

$$(27) \quad |T_1| \leq t'_1 := \begin{cases} \sum_{p>5} \left\lceil \frac{k}{p} \right\rceil & \text{if } k = 7, 11 \\ \sum_{p>5} \left\lceil \frac{k}{p} \right\rceil - 2 & \text{if } 13 \leq k < 23 \\ \sum_{p>5} \left\lceil \frac{k}{p} \right\rceil - 3 & \text{if } k = 23 \end{cases}$$

where the sum is taken over all $p \leq k$. For the last sum, we observe that 7 and 11 together divide at most six a_i 's when $k = 23$. We divide the proof into 4 cases.

Case I. Let $2 \nmid d$ and $3 \nmid d$. From (24), (25), (26), (10) and Lemma 1, we get

$$|T| \leq t_1 := \begin{cases} \max(f_1(k, 1, 0) + f_1(k, 6, 0), f_1(k, 2, 0) + f_1(k, 3, 0)) + \left\lceil \frac{k}{4} \right\rceil & \text{if } k = 7, 11, \\ f_1(k, 1, 0) + f_1(k, 3, 0) + f_1(k, 5, 0) + f_1(k, 15, 0) + \left\lceil \frac{k}{4} \right\rceil & \text{if } k > 11 \end{cases}$$

since $f_1(k, a, \delta)$ is non-increasing function of a and $\sum_{a \in R} \nu_e(a) \leq \left\lceil \frac{k}{4} \right\rceil$. We check that $k = |T| + |T_1| \leq t_1 + t'_1 < k$, a contradiction.

Thus we have either $2|d$ or $3|d$. Let $k = 7, 11$. If $2|d$, then $S \subseteq \{1\}$ or $S \subseteq \{3\}$. If $3|d$, we have $S \subseteq \{1\}$ or $S \subseteq \{2\}$. By Lemma 2, we get $|T| \leq \frac{k-1}{2}$. We check that $k = |T| + |T_1| \leq \frac{k-1}{2} + t'_1 < k$ by (27). This is a contradiction. From now on, we may also that suppose that $13 \leq k \leq 23$.

Case II. Let $2|d$ and $3 \nmid d$. Then $S \subseteq \{1, 3, 5, 15\}$ if $11|d$ and $S \subseteq \{1, 3\}$ or $S \subseteq \{5, 15\}$ if $13|d$. Let $2||d$. From (10) and Lemma 1 with $\delta = 1$, we get

$$|T| \leq F(k, 1, 1) + F(k, 3, 1) + F(k, 5, 1) + F(k, 15, 1) =: t_2.$$

Let $4||d$. From $a_i \equiv n \pmod{4}$, we see that $S \subseteq \{1, 5\}$ or $S \subseteq \{3, 15\}$ if $11|d$ and either $S = \emptyset$ or $S = \{1\}, \{3\}, \{5\}$ or $\{15\}$ if $13|d$. Therefore

$$|T| \leq F(k, 1, 2) + F(k, 5, 2) =: t_3.$$

by Lemma 1 with $\delta = 2$. Let $8|d$. Then $a_i \equiv n \pmod{8}$ and Lemma 1 with $\delta = 3$ imply

$$|T| \leq F(k, 1, 3) =: t_4.$$

Thus $|T| \leq \max(t_2, t_3, t_4)$. This with (27) contradicts (9).

Case III. Let $2 \nmid d$ and $3|d$. From $a_i \equiv n \pmod{3}$, we see that either $S = \emptyset$ or $S = \{1\}, \{2\}, \{5\}$ or $\{10\}$ if $11|d$ and $S \subseteq \{1, 10\}$ or $S \subseteq \{2, 5\}$ if $13|d$. By (10) and Lemma 1, we get

$$|T| \leq F(k, 1, 0) + F(k, 5, 0),$$

which together with (27) contradicts (9).

Case IV. Let $2|d$ and $3|d$. Then $S \subseteq \{1\}, \{5\}$. By Lemma 2, we get $|T| \leq \frac{k-1}{2}$. We check that $k = |T| + |T_1| \leq \frac{k-1}{2} + t'_1 < k$, a contradiction.

5.2. **The cases $k \geq 29$.** Let $29 \leq k \leq 59$ and $19|d$. Then by Lemma 7 with $Q_0 = 19$, we get $7|d$ or $17|d$. Thus we get a prime pair $(Q, Q') = (7, 19)$ or $(Q, Q') = (17, 19)$ such that $QQ'|d$. Similarly we get $(Q, Q') = (7, 29)$ or $(Q, Q') = (17, 29)$ with $QQ'|d$ when $31 \leq k \leq 59$ and $29|d$. Let $k = 71$. Then we have either $43|d, 67|d$ or $43|d, 67 \nmid d$ or $43 \nmid d, 67|d$. We get prime pair $(Q, Q') = (43, 67)$ with $QQ'|d$ if $43|d, 67|d$. If $43|d, 67 \nmid d$, we get from Lemma 7 with $Q_0 = 43$ that $53|d$ and we take $(Q, Q') = (43, 53)$ such that $QQ'|d$. If $43 \nmid d, 67|d$, we get from Lemma 7 with $Q_0 = 67$ that $53|d$ and we take $(Q, Q') = (53, 67)$ such that $QQ'|d$. Similarly we get prime pairs (Q, Q') with $QQ'|d$ for each $61 \leq k \leq 97$ are given in the table below. For $q \leq 17$, we see that

$$(28) \quad |T_1| \leq \sum_{\substack{p>q \\ p \neq Q, Q'}} \left[\frac{k}{p} \right] \leq t'_2 := \begin{cases} \sum_{p>q} \left[\frac{k}{p} \right] - 2 & \text{if } 29 \leq k \leq 61 \\ \sum_{p>q} \left[\frac{k}{p} \right] - 4 & \text{if } 61 < k < 97 \\ \sum_{p>q} \left[\frac{k}{p} \right] - 7 & \text{if } k = 97 \end{cases}$$

where the sum is taken over primes $\leq k$.

Case I. Let $2 \nmid d$ and $3 \nmid d$. We take $q = 11$ if $k = 71$, $(Q, Q') = (43, 67)$ and $q = 7$ otherwise, in (7) and (8). From $\binom{a_i}{Q} = \binom{n}{Q}$ and $\binom{a_i}{Q'} = \binom{n}{Q'}$, we get $S \subseteq S' = \{s : s \text{ squarefree, } P(s) \leq q, \binom{s}{Q} = \binom{n}{Q}, \binom{s}{Q'} = \binom{n}{Q'}\}$. By considering $\left(\binom{n}{Q}, \binom{n}{Q'}\right) = (1, 1), (1, -1), (-1, 1)$ and $(-1, -1)$, we get four possibilities of S' . For each value of k , we give below a table for (Q, Q') and S' .

k	(Q, Q')	$S \subseteq S'$ with S' given by one of
$29 \leq k \leq 59$	$(7, 19), (7, 29)$	$\{1, 30\}, \{2, 15\}, \{3, 10\}, \{5, 6\}$
$29 \leq k \leq 59$	$(17, 19), (17, 29)$	$\{1, 30, 35, 42\}, \{2, 15, 21, 70\}, \{3, 10, 14, 105\}, \{5, 6, 7, 210\}$
61	$(11, 59)$	$\{1, 3, 5, 15\}, \{2, 6, 10, 30\}, \{7, 21, 35, 105\}, \{14, 42, 70, 210\}$
67, 71	$(43, 53)$	$\{1, 6, 10, 15\}, \{2, 3, 5, 30\}, \{7, 42, 70, 105\}, \{14, 21, 35, 210\}$
71	$(43, 67)$	See (29)
71	$(53, 67)$	$\{1, 6, 10, 15\}, \{2, 3, 5, 30\}, \{7, 42, 70, 105\}, \{14, 21, 35, 210\}$
73	$(23, 53)$	$\{1, 6, 70, 105\}, \{2, 3, 35, 210\}, \{5, 14, 21, 30\}, \{7, 10, 15, 42\}$
73	$(23, 67)$	$\{1, 6, 35, 210\}, \{2, 3, 70, 105\}, \{5, 7, 30, 42\}, \{10, 14, 15, 21\}$
79	$(23, 53), (53, 73)$	$\{1, 6, 70, 105\}, \{2, 3, 35, 210\}, \{5, 14, 21, 30\}, \{7, 10, 15, 42\}$
79	$(23, 67), (67, 73)$	$\{1, 6, 35, 210\}, \{2, 3, 70, 105\}, \{5, 7, 30, 42\}, \{10, 14, 15, 21\}$
83	$(23, 37), (37, 73)$	$\{1, 3, 70, 210\}, \{2, 6, 35, 105\}, \{5, 14, 15, 42\}, \{7, 10, 21, 30\}$
89	$(23, 79), (73, 79)$	$\{1, 2, 105, 210\}, \{3, 6, 35, 70\}, \{5, 10, 21, 42\}, \{7, 14, 15, 30\}$
97	$(23, 37), (23, 83)$	$\{1, 3, 70, 210\}, \{2, 6, 35, 105\}, \{5, 14, 15, 42\}, \{7, 10, 21, 30\}$

For $k = 71$, $(Q, Q') = (43, 67)$, we get $S \subseteq S'$ with S' given by one of

$$(29) \quad \begin{aligned} & \{1, 6, 10, 14, 15, 21, 35, 210\}, \{2, 3, 5, 7, 30, 42, 70, 105\} \\ & \{11, 66, 110, 154, 165, 231, 385, 2310\}, \{22, 33, 55, 77, 330, 462, 770, 1155\}. \end{aligned}$$

From the possibilities of $S \subseteq S'$ given by the above table, (10) and Lemma 1, we get

$$|T| \leq t_5 := \max \sum_{s \in S'} F(k, s, 0)$$

where the maximum is taken over all the four choices of S' . This with (28) gives $|T| + |T_1| \leq t_5 + t'_2 < k$ a contradicting (9).

Case II. Let $2|d$ and $3 \nmid d$. We take $q = 7$ for $2||d, 4||d$ and $q = 11$ for $8|d$. Let $2||d$. Then $S \subseteq \{1, 3, 5, 7, 15, 21, 35, 105\} =: S_2$. From (10) and Lemma 1 with $\delta = 1$, we get

$$|T| \leq \sum_{s \in S_2} F(k, s, 1) =: t_6$$

Let $4||d$. Then we see that either $S \subseteq \{1, 5, 21, 105\} =: S_{41}$ or $S \subseteq \{3, 7, 15, 35\} =: S_{42}$. From (10) and Lemma 1 with $\delta = 2$, we get

$$|T| \leq \max_{i=1,2} \sum_{s \in S_{4i}} F(k, s, 2) =: t_7.$$

Hence, if $8 \nmid d$, then $|T| \leq \max(t_6, t_7)$. This with (28) implies $|T| + |T_1| \leq \max(t_6, t_7) + t'_2 < k$, contradicting (9).

Let $8|d$. Then we see from $a_i \equiv n \pmod{8}$ that $S \subseteq \{1, 33, 105, 385\} =: S_{81}$ or $S \subseteq \{3, 11, 35, 1155\} =: S_{82}$ or $S \subseteq \{5, 21, 77, 165\} =: S_{83}$ or $S \subseteq \{7, 15, 55, 231\} =: S_{84}$. Then

$$|T| \leq \max_{1 \leq i \leq 4} \sum_{s \in S_{8i}} F(k, s, 3) =: t_8.$$

by Lemma 1 with $\delta = 3$. This with (28) implies $|T| + |T_1| \leq t_8 + t'_2 < k$, a contradiction.

Case III. Let $2 \nmid d$ and $3|d$. We take $q = 11$. Then by modulo 3, we get either $S \subseteq \{1, 7, 10, 22, 55, 70, 154, 385\} =: S_{31}$ or $S \subseteq \{2, 5, 11, 14, 35, 77, 110, 770\} =: S_{32}$. By (10) and Lemma 1, we get

$$|T| \leq \max_{i=1,2} \sum_{s \in S_{3i}} F(k, s, 0) =: t_9.$$

This together with (28) contradicts (9).

Case IV. Let $2|d$ and $3|d$. Let $2||d$. We take $q = 7$. Then we see that either $S \subseteq \{1, 7\}$ or $S \subseteq \{5, 35\}$. By (10) and Lemma 1, we get $|T| \leq F(k, 1, 1) + F(k, 7, 1)$ which together with (28) contradicts (9).

Let $4||d$. We take $q = 13$. From $a_i \equiv n \pmod{12}$, we see that $S \subseteq S' \in \mathfrak{S} := \{\{1, 13, 385, 5005\}, \{5, 65, 77, 1001\}, \{7, 55, 91, 715\}, \{11, 35, 143, 455\}\}$. Then

$$|T| \leq \max_{S' \in \mathfrak{S}} \sum_{s \in S'} F(k, s, 2)$$

which together with (28) contradicts (9).

Let $8|d$. We take $q = 17$. From $a_i \equiv n \pmod{24}$, we see that $S \subseteq S' = \{1, 385, 1105, 17017\}$ or $S \subseteq S'' \in \mathfrak{S}_1$ where \mathfrak{S}_1 is the union of sets

$$\{5, 77, 221, 85085\}, \{7, 55, 2431, 7735\}, \{11, 35, 1547, 12155\}, \{13, 85, 1309, 5005\}, \\ \{17, 65, 1001, 6545\}, \{91, 187, 595, 715\}, \{119, 143, 455, 935\}.$$

Let $S \subseteq S'' \in \mathfrak{S}_1$. Then

$$|T| \leq \max_{S'' \in \mathfrak{S}_1} \sum_{s \in S''} F(k, s, 3) =: t_{10}.$$

Let $S \subseteq S'$. By Lemma 2, we get $\nu(1) \leq \frac{k-1}{2}$. This together with $\nu(1105) + \nu(17017) \leq 1$ by $13 \cdot 17 | \gcd(1105, 17017)$ and $\nu(385) \leq 1$ by Lemma 1 gives $|T| \leq \frac{k-1}{2} + 2$. Therefore $|T| \leq \max(t_{10}, \frac{k-1}{2} + 2)$. This with (28) contradicts (9). \square

6. PROOF OF THEOREM 4

Let $k = 7$. By the case $k = 6$, we may assume that $7 \nmid d$. Now the assertion follows from Lemmas 8 and 6. Let $k = 8$. Then by applying the case $k = 7$ twice to $n(n+d) \cdots (n+6d) = b'y'^2$ and $(n+d) \cdots (n+7d) = b''y''^2$, we get

$$(a_0, \dots, a_6), (a_1, \dots, a_7) \in \{(2, 3, 1, 5, 6, 7, 2), (3, 1, 5, 6, 7, 2, 1), (1, 5, 6, 7, 2, 1, 10), \\ (2, 7, 6, 5, 1, 3, 2), (1, 2, 7, 6, 5, 1, 3), (10, 1, 2, 7, 6, 5, 1)\}.$$

This gives $(a_0, \dots, a_7) = (2, 3, 1, 5, 6, 7, 2, 1), (3, 1, 5, 6, 7, 2, 1, 10)$ or their mirror images and the assertion follows. Let $k = 9$. By applying the case $k = 8$ twice to $n(n+d) \cdots (n+7d) = b'y'^2$ and $(n+d) \cdots (n+8d) = b''y''^2$, we get the result. Let $k = 10$. By applying $k = 9$ twice, we get $(a_0, a_1, \dots, a_8), (a_1, a_2, \dots, a_8, a_9) \in \{(2, 3, \dots, 1, 10), (10, 1, \dots, 3, 2)\}$ which is not possible.

Let $k \geq 11$ and $k' < k$ be consecutive primes. We suppose that Theorem 4 is valid with k replaced by k' . Let $k|d$. Then $\binom{a_i}{k} = \binom{n}{k}$ for all $0 \leq i < k$. By applying the case $k = k'$ to $n(n+d) \cdots (n+(k'-1)d) = b'y'^2$ with $P(b') \leq k'$, we get $k' \leq 23$ and $1, 2, 3, 5 \in \{a_0, a_1, a_2, \dots, a_{k'-1}\}$ in view of (5) and (6). Therefore $\binom{2}{k} = \binom{3}{k} = \binom{5}{k} = 1$ which is not possible.

Thus we may assume that $k \nmid d$ and $k|n+id$ for some $0 \leq i \leq \frac{k-1}{2}$ by considering the mirror image (4) of (2) whenever Theorem 4 holds at k' . We shall use this assertion without reference in the proof of Theorem 4.

Let $k = 11$. By Lemmas 8 and 6, we see that $11|n+id$ for $0 \leq i \leq 3$. If $11|n$, the assertion follows by the case $k = 10$. Let $11|n+d$. We consider $(n+2d) \cdots (n+10d) = b'y'^2$ with $P(b') \leq 7$ and the case $k = 9$ to get $(a_2, a_3, \dots, a_{10}) \in \{(2, 3, 1, 5, 6, 7, 2, 1, 10), (10, 1, 2, 7, 6, 5, 1, 3, 2)\}$. The first possibility is excluded since $1 = \binom{14}{11} = \binom{a_2 a_7}{11} = \binom{1 \cdot 6}{11} = -1$. For the second possibility, we observe $P(a_0) \leq 5$ since $\gcd(a_0, 7 \cdot 11) = 1$ and this is excluded by the case $k = 6$ applied to $n(n+2d)(n+4d)(n+6d)(n+8d)(n+10d)$. Let $11|n+2d$. Then by the case $k = 8$, we have $(a_3, a_4, \dots, a_{10}) \in \{(2, 3, 1, 5, 6, 7, 2, 1), (3, 1, 5, 6, 7, 2, 1, 10), (1, 2, 7, 6, 5, 1, 3, 2), (10, 1, 2, 7, 6, 5, 1, 3)\}$. The first three possibilities are excluded by

considering the values of Legendre symbol mod 11 at a_3, a_8 ; a_3, a_4 and a_3, a_5 , respectively. If the last possibility holds, then $a_0 = 1$ since $\gcd(a_0, 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11) = 1$ and this is not possible since $1 = \left(\frac{a_0 a_4}{11}\right) = \left(\frac{(-2)^2}{11}\right) = -1$. Let $11|n + 3d$. We consider $(n + 4d) \cdots (n + 10d) = b'y'^2$ with $P(b') \leq 7$ and the case $k = 7$ to get $(a_4, \dots, a_{10}) \in \{(2, 3, 1, 5, 6, 7, 2), (3, 1, 5, 6, 7, 2, 1), (1, 5, 6, 7, 2, 1, 10), (2, 7, 6, 5, 1, 3, 2), (1, 2, 7, 6, 5, 1, 3), (10, 1, 2, 7, 6, 5, 1)\}$ which is not possible as above. This completes the proof for $k = 11$. The assertion for $k = 12$ follows from that of $k = 11$.

Let $k = 13$. Then the assertion follows from Lemmas 8, 6 and the case $k = 11$. Let $k = 14$. By applying $k = 13$ to $n(n + d) \cdots (n + 12d) = b'y'^2$ and $(n + d) \cdots (n + 13) = b''y''^2$, we get the assertion. Let $k = 15$. Then applying $k = 14$ both to $n(n + d) \cdots (n + 13d)$ and $(n + d) \cdots (n + 14d)$ gives the result. Now $k = 16$ follows from the case $k = 15$.

Let $k = 17$. Then $17|n + 2d$ or $17|n + 3d$ by Lemmas 8, 6 and the case $k = 15$. Let $17|n + 2d$. Then by applying the case $k = 14$ to $(n + 3d) \cdots (n + 16d) = b'y'^2$ with $P(b') \leq 13$, we get $(a_3, a_4, \dots, a_{16}) \in \{(3, 1, \dots, 15, 1), (1, 15, \dots, 1, 3)\}$. The first possibility is excluded by Legendre symbol mod 17 at a_3, a_4 . For the second, we observe that $\gcd(a_1, 7 \cdot 11 \cdot 13 \cdot 17) = 1$ which is not possible by the case $k = 6$ applied to $(n + d)(n + 4d)(n + 7d)(n + 10d)(n + 13d)(n + 16d)$. Let $17|n + 3d$. By considering $(n + 4d) \cdots (n + 16d) = b'y'^2$ with $P(b') \leq 13$, it follows from the case $k = 13$ that $(a_4, \dots, a_{16}) \in \{(3, 1, \dots, 14, 15), (1, 5, \dots, 15, 1), (15, 14, \dots, 1, 3), (1, 15, \dots, 5, 1)\}$. The first three possibilities are excluded by considering Legendre symbol mod 17 at a_4, a_5 . If the last possibility holds, we observe that $a_1 = 1$ since $\gcd(a_1, \prod_{p \leq 17} p) = 1$ and then $1 = \left(\frac{a_1 a_4}{17}\right) = \left(\frac{(-6)(-3)}{17}\right) = -1$, a contradiction. The assertion for $k = 18$ follows from that of $k = 17$.

Let $k = 19$. Then the assertion follows from Lemmas 8, 6 and the case $k = 17$. By applying $k = 19$ twice to $n(n + d) \cdots (n + 18d)$ and $(n + d) \cdots (n + 18d)(n + 19d)$, the assertion for $k = 20$ follows and this implies the cases $k = 21, 22$.

Let $k = 23$. We see from Lemmas 8, 6 and the case $k = 20$ that $23|n + 3d$. We consider $k = 19$ and $(n + 4d) \cdots (n + 22d) = b'y'^2$ with $P(b') \leq 19$ to get $(a_4, a_5, \dots, a_{22}) = (1, 5, \dots, 21, 22)$ or $(22, 21, \dots, 5, 1)$. By considering the values of Legendre symbol mod 23 at a_4 and a_5 , we may assume the second possibility. Now $P(a_2) \leq 11$ and this is not possible by the case $k = 11$ applied to $(n + 2d)(n + 4d) \cdots (n + 22d)$. Let $k = 24$. We get $(a_0, a_1, \dots, a_{23}) = (5, 6, \dots, 3, 7), (7, 3, \dots, 6, 5)$ by considering $k = 23$ both to $n(n + d) \cdots (n + 22d)$ and $(n + d) \cdots (n + 23d)$. Further the assertion for $25 \leq k \leq 28$ follows from $k = 24$.

Let $k \geq 29$. First we consider $k = 29$. We see from Lemmas 8, 6 and the case $k = 25$ that $29|n + 4d$ or $29|n + 5d$. Let $29|n + 4d$. Then considering $k = 24$ and $(n + 5d)(n + 6d) \cdots (n + 28d)$, we get $(a_5, a_6, \dots, a_{28}) = (5, 6, \dots, 3, 7)$ or $(7, 3, \dots, 6, 5)$. By observing $1 = \left(\frac{30}{29}\right) = \left(\frac{a_5 a_6}{29}\right) = \left(\frac{1 \cdot 2}{29}\right) = -1$, we may assume the second possibility. Then $a_1 = 1$ implying $1 = \left(\frac{a_2 a_8}{29}\right) = \left(\frac{(-2)^4}{29}\right) = -1$, a contradiction. Let $29|n + 5d$. Now by considering $k = 23$ and $(n + 6d) \cdots (n + 28d)$, we get

$(a_6, a_7, \dots, a_{28}) \in \{(5, 6, \dots, 26, 3), (6, 7, \dots, 3, 7), (3, 26, \dots, 6, 5), (7, 3, \dots, 7, 6)\}$. Then we may restrict to the last possibility by considering the Legendre symbol mod 29 at the first two entries in the remaining possibilities. It follows that $a_3 = 1$ implying $1 = \left(\frac{a_3 a_9}{29}\right) = \left(\frac{(-2)^4}{29}\right) = -1$, a contradiction. This completes the proof for $k = 29$. We now proceed by induction. By Lemmas 8 and 6, the assertion follows for all primes k . Now Lemma 3 completes the proof of Theorem 4. \square

7. PROOF OF THEOREM 1

Observe that for all tuples in (5) and (6), the product of the a_i 's is not a square. Hence, by Theorem 4, we may assume that $101 \leq k \leq 109$. Assume (1). Then $\text{ord}_p(a_0 a_1 \cdots a_{k-1})$ is even for each prime p . Let $101 \leq k \leq 105$. Then $P(a_4 a_5 \cdots a_{100}) \leq 97$. Now the assertion follows from Theorem 4 by considering $(n + 4d) \cdots (n + 100d)$ and $k = 97$. Let $k = 106, 107$. Then $P(a_4 a_5 \cdots a_{102}) \leq 101$. We may suppose that $P(a_4 a_5) = 101$ or $P(a_{101} a_{102}) = 101$ otherwise the assertion follows by the case $k = 99$ in Theorem 4. Let $P(a_4 a_5) = 101$. Then $P(a_6 \cdots a_{102}) \leq 97$ and the assertion follows by $k = 97$ in Theorem 4. This is also the case when $P(a_{101} a_{102}) = 101$ since $P(a_4 \cdots a_{100}) \leq 97$ in this case. Let $k = 108, 109$. Then $P(a_6 \cdots a_{102}) \leq 101$. Thus either $P(a_6 a_7) = 101$ or $P(a_{101} a_{102}) = 101$. Let $P(a_6 a_7) = 101$. Then $P(a_8 \cdots a_{102}) \leq 97$. We may assume that $97 | a_8 a_9 a_{10} a_{11}$ or $97 | a_{97} \cdots a_{101} a_{102}$. Let $97 | a_8 a_9 a_{10} a_{11}$. Then $P(a_{12} a_{13} \cdots a_{102}) \leq 89$ and the assertion follows by the case $k = 91$ of Theorem 4. Let $97 | a_{97} \cdots a_{102}$. Then $P(a_8 a_9 \cdots a_{96}) \leq 89$ and the assertion follows from the case $k = 89$ of Theorem 4. When $P(a_{101} a_{102}) = 101$, we argue as above to get the assertion. \square

REFERENCES

- [BBGH06] M.A. Bennett, N. Bruin, K. Györy and L. Hajdu, *Powers from products of consecutive terms in arithmetic progression*, Proc. London Math. Soc. **92** (2006), 273-306.
- [BF05] N. Bruin and E. V. Flynn, *Towers of 2-covers of hyperelliptic curves*, Trans. Amer. Math. Soc. **357** (2005), 4329-4347.
- [Erd39] P. Erdős, *Note on the product of consecutive integers* (II), Jour. London Math. Soc. **14** (1939), 245-249.
- [ES75] P. Erdős and J.L. Selfridge, *The product of consecutive integers is never a power*, Illinois Jour. Math. **19** (1975), 292-301.
- [Eul80] L. Euler, *Mém. Acad. Sc. St. Peters.* **8** 1817-1818 (1820), 3; *Comm. Arith.* **II** (1849), 411-413.
- [Lai04] S. Laishram, *Topics in Diophantine equations*, M.Sc. Thesis, Mumbai University, 2004, online at <http://www.math.tifr.res.in/~shanta/MScthesis.pdf>.
- [LS] S. Laishram and T. N. Shorey, *The equation $n(n+d) \cdots (n+(k-1)d) = by^2$ with $\omega(d) \leq 6$ or $d \leq 10^{10}$* , Acta Arith., accepted for publication.
- [Mor69] L.J. Mordell, *Diophantine equations*, Academic Press, New York (1969).
- [MS03] A. Mukhopadhyay and T.N. Shorey, *Almost squares in arithmetic progression* (II), Acta Arith. **110** (2003), 1-14.
- [Obl50] R. Obläth, *Über das Produkt fünf aufeinander folgender Zahlen in einer arithmetischen Reihe*, Publ. Math. Debrecen **1** (1950), 222-226.

- [Rig39] O. Rigge, *Über ein diophantisches problem*, In: 9th Congress Math. Scand. Helsingfors, 1938, Mercator (1939), 155-160.
- [Tij89] R. Tijdeman, *Diophantine equations and diophantine approximations*, Number Theory and Applications, Kluwer Acad. Press (1989), 215-243.

DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE AND TECHNOLOGY, NIHON UNIVERSITY, TOKYO 101-8308, JAPAN

E-mail address: `hirata@math.cst.nihon-u.ac.jp`

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, MUMBAI 400005, INDIA

E-mail address: `shanta@math.tifr.res.in`

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, MUMBAI 400005, INDIA

E-mail address: `shorey@math.tifr.res.in`

MATHEMATICAL INSTITUTE, LEIDEN UNIVERSITY, 2300 RA LEIDEN, THE NETHERLANDS

E-mail address: `tijdeman@math.leidenuniv.nl`