

isid/ms/2016/05

May 23, 2016

<http://www.isid.ac.in/~statmath/index.php?module=Preprint>

On the Kolmogorov - Wiener - Masani spectrum of a multi-mode weakly stationary quantum process

K. R. PARTHASARATHY AND RITABRATA SENGUPTA

Indian Statistical Institute, Delhi Centre
7, SJSS Marg, New Delhi-110 016, India

ON THE KOLMOGOROV - WIENER - MASANI SPECTRUM OF A MULTI-MODE WEAKLY STATIONARY QUANTUM PROCESS

K. R. PARTHASARATHY AND RITABRATA SENGUPTA

ABSTRACT. We introduce the notion of a k -mode weakly stationary quantum process ϱ based on the canonical Schrödinger pairs of position and momentum observables in copies of $L^2(\mathbb{R}^k)$, indexed by an additive abelian group D of countable cardinality. Such observables admit an autocovariance map \tilde{K} from D into the space of real $2k \times 2k$ matrices. The map \tilde{K} on the discrete group D admits a spectral representation as the Fourier transform of a $2k \times 2k$ complex Hermitian matrix-valued totally finite measure Φ on the compact character group \hat{D} , called the Kolmogorov-Wiener-Masani (KWM) *spectrum* of the process ϱ . Necessary and sufficient conditions on a $2k \times 2k$ complex Hermitian matrix-valued measure Φ on \hat{D} to be the KWM spectrum of a process ϱ are obtained. This enables the construction of examples. Our theorem reveals the dramatic influence of the uncertainty relations among the position and momentum observables on the KWM spectrum of the process ϱ . In particular, KWM spectrum cannot admit a gap of positive Haar measure in \hat{D} .

The relationship between the number of photons in a particular mode at any site of the process and its KWM spectrum needs further investigation.

Keywords. Weakly stationary quantum process, Kolmogorov-Wiener-Masani spectrum, autocovariance map, spectral representation, uncertainty relations.

Mathematics Subject Classification (2010): 81S25, 60G25, 60G10.

1. INTRODUCTION

In his celebrated little book “Osnovnye ponyatiya teorii veroyatnostei” [Kol74], A. N. Kolmogorov introduced the notion of a stochastic process as a consistent family of finite dimensional probability distributions in \mathbb{R}^n , $n = 1, 2, \dots$. In the same spirit a quantum process can be described as a consistent family of density operators or, equivalently, states in tensor products $\mathcal{H}_1 \otimes \dots \otimes \mathcal{H}_n$ of Hilbert spaces with $n = 1, 2, \dots$. One can replace the ‘time set’ $\{1, 2, \dots\}$ by an abstract countable set D with the discrete topology and a family $\{\mathcal{H}_a : a \in D\}$ of Hilbert spaces. Then a quantum process yields a density operator $\rho_{a_1, a_2, \dots, a_n}$ in $\mathcal{H}_{a_1} \otimes \dots \otimes \mathcal{H}_{a_n}$ for every finite sequence (a_1, \dots, a_n) with distinct elements from D . All these density operators will obey natural consistency conditions. For example, the relative trace of $\rho_{a_1, a_2, \dots, a_n}$ over \mathcal{H}_{a_n} is $\rho_{a_1, a_2, \dots, a_{n-1}}$. If (b_1, \dots, b_n) is a permutation of a_1, \dots, a_n then $\rho_{b_1, \dots, b_n} = U \rho_{a_1, \dots, a_n} U^{-1}$ where U is the corresponding Hilbert space isomorphism from $\mathcal{H}_{a_1} \otimes \dots \otimes \mathcal{H}_{a_n}$ onto $\mathcal{H}_{b_1} \otimes \dots \otimes \mathcal{H}_{b_n}$ induced by the permutation. We denote the quantum process over D by

$$(1.1) \quad \varrho = \{(\mathcal{H}_{a_1, \dots, a_n}, \rho_{a_1, \dots, a_n}) : (a_1, \dots, a_n) \in \mathcal{S}_D\}$$

where \mathcal{S}_D denotes the set of all finite length sequences of distinct elements from the countable set D .

RS acknowledges financial support from the National Board for Higher Mathematics, Govt. of India.

In this paper we are interested in the special case where $\mathcal{H}_a = L^2(\mathbb{R}^k)$ for all a in D , k being a fixed positive integer, called the *number of modes* of the process. Each \mathcal{H}_a admits Schrödinger canonical pairs q_{ar}, p_{ar} , $r = 1, 2, \dots, k$ of position and momentum observables obeying the Heisenberg canonical commutation relations (CCR). We can look upon q_{ar}, p_{ar} , $r = 1, 2, \dots, k$ as observables in $\mathcal{H}_{a_1} \otimes \dots \otimes \mathcal{H}_{a_n}$ whenever the sequence (a_1, a_2, \dots, a_n) from \mathcal{S}_D contains the element a and denote such amplified observables by the same respective symbols. With such a convention one obtains the algebra of all polynomials of all q_{ar}, p_{ar} , $r = 1, 2, \dots, k$, $a \in D$. Using the finite-partite states ρ_{a_1, \dots, a_n} , $(a_1, \dots, a_n) \in \mathcal{S}_D$ one can compute the expectations of the polynomials whenever they exist. Write

$$(X_{a_1}, X_{a_2}, \dots, X_{a_{(2k-1)}}, X_{a_{2k}}) = (q_{a_1}, p_{a_1}, \dots, q_{a_k}, p_{a_k})$$

and define the covariances

$$(1.2) \quad \kappa_{rs}(a, b) = \left\langle \frac{1}{2} (X_{ar} X_{bs} + X_{bs} X_{ar}) \right\rangle - \langle X_{ar} \rangle \langle X_{bs} \rangle$$

where $\langle \cdot \rangle$ denotes expectation. To compute these quantities we need a knowledge of only the ‘bipartite’ states $\rho_{a,b}$, $(a, b) \in \mathcal{S}_D$. Thus we obtain a $2k \times 2k$ real matrix-valued covariance kernel $\mathcal{K} = [[K(a, b)]]$ defined by

$$(1.3) \quad K(a, b) = [[\kappa_{rs}(a, b)]], \quad r, s, \in \{1, 2, \dots, 2k\},$$

for $a, b \in D$.

Suppose D is a countable discrete additive abelian group with addition operation $+$ and null element 0 . Let the covariance kernel \mathcal{K} of a k -mode quantum process over D be translation invariant in the sense that

$$(1.4) \quad K(a+x, b+x) = K(a, b) \quad \forall a, b, x \in D.$$

Then we say that the quantum process is second order *weakly stationary*, or, simply, weakly stationary. For such a process there exists a map \tilde{K} from D into the space of $2k \times 2k$ real matrices such that

$$(1.5) \quad K(a, b) = \tilde{K}(b-a), \quad a, b \in D.$$

The map \tilde{K} is called the *autocovariance map* of the weakly stationary quantum process.

Owing to the matrix-positivity properties enjoyed by covariances between observables the autocovariance map \tilde{K} satisfies the matrix inequalities

$$(1.6) \quad \sum_{i,j} \alpha_i \alpha_j \tilde{K}(a_j - a_i) \geq 0$$

for any $a_1, a_2, \dots, a_n \in D$ and real scalars $\alpha_1, \alpha_2, \dots, \alpha_n$, $n = 1, 2, \dots$. Thanks to Bochner’s theorem for locally compact abelian groups there exists a complex-Hermitian and positive $2k \times 2k$ matrix-valued measure Φ on the compact dual character group \hat{D} of D such that

$$(1.7) \quad \tilde{K}(a) = \int_{\hat{D}} \chi(a) \Phi(d\chi), \quad \text{for all } a \text{ in } D.$$

The matrix-valued measure Φ satisfies the conjugate symmetry property

$$(1.8) \quad \Phi(S^{-1}) = \overline{\Phi(S)}$$

for any Borel set $S \subset \widehat{D}$. Furthermore, the Heisenberg uncertainty relations prevailing among the various position and momentum observables of the quantum process reveal their dramatic influence on the measure Φ through the matrix inequalities

$$(1.9) \quad \Phi(S) + \frac{i}{2}\lambda(S)J_{2k} \geq 0$$

for all Borel sets $S \subset \widehat{D}$, where λ is the normalised Haar measure of the compact group \widehat{D} and J_{2k} is the fundamental symplectic matrix given by

$$(1.10) \quad J_{2k} = \bigoplus_{k\text{-copies}} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

which is a diagonal block matrix with each diagonal block equal to J_2 . The inequality (1.9) implies, in particular, that whenever $\Phi(S) = 0$, $\lambda(S)$ is also zero.

Borrowing from the extensive theory of linear least square prediction of real valued weakly stationary processes pioneered by A. N. Kolmogorov [Kol41a, Kol41b] and N. Wiener [Wie49], and multivariate weakly stationary processes by N. Wiener and P. Masani [WM57, WM58] we call (1.7) the spectral representation of $\widetilde{\mathcal{K}}$ in \widehat{D} and the matrix-valued positive measure Φ the *Kolmogorov - Wiener - Masani spectrum* (or *KWM spectrum*) of the autocovariance map $\widetilde{\mathcal{K}}$ of the underlying quantum process.

Inequality (1.9) implies that whenever $\Phi(S) = 0$ for some Borel set $S \subset \widehat{D}$, then $\lambda(S) = 0$. In other words, the KWM spectrum does not admit a ‘Haar gap’.

Conversely, given a complex Hermitian positive $2k \times 2k$ matrix-valued measure Φ on the Borel σ -algebra of \widehat{D} satisfying the conjugate symmetry condition (1.8), the spectral uncertainty relations (1.9), and the condition $\Phi(\widehat{D}) = M$, there exists a weakly stationary k -mode quantum process over D with KWM spectrum Φ . Indeed, such a process can be realized as a mean zero quantum Gaussian process in the sense that all its finite-partite states ρ_{a_1, \dots, a_n} , $(a_1, \dots, a_n) \in \mathcal{S}_D$ are mean zero Gaussian states.

The spectral representation of the autocovariance function and its converse enable us to construct interesting examples of weakly stationary quantum processes.

2. QUANTUM PROCESSES

A quantum system in its most elementary form is determined by a pair (\mathcal{H}, ρ) where \mathcal{H} is a complex separable Hilbert space and ρ is a density operator in \mathcal{H} , i.e., a positive operator with unit trace. The operator ρ is called the state of the system. We shall deal with several quantum systems and assume that all the Hilbert spaces in this paper are complex and separable. Scalar products in Hilbert spaces will be expressed in the Dirac notation and adjoints of operators as well as matrices will be indicated by the symbol \dagger . By a positive operator X in a Hilbert space \mathcal{H} we mean that $\langle u|X|u \rangle \geq 0$ for all $u \in \mathcal{H}$. By a positive $n \times n$ matrix we mean an $n \times n$ Hermitian matrix which is positive semidefinite.

If $\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \dots \otimes \mathcal{H}_n$ is the tensor product of Hilbert spaces \mathcal{H}_i , $1 \leq i \leq n$, ρ is a state in \mathcal{H} and $F \subset \{1, 2, \dots, n\}$ is the subset $\{i_1 < i_2 < \dots < i_k\}$ then we write $\mathcal{H}_F = \mathcal{H}_{i_1} \otimes \dots \otimes \mathcal{H}_{i_k}$. One obtains a state ρ_F in \mathcal{H}_F by taking the relative trace of ρ successively in \mathcal{H}_i , $i \notin F$ in some

order. The resulting state ρ_F is independent of the order in which the traces are taken. The system (\mathcal{H}_F, ρ_F) is called the F -marginal of (\mathcal{H}, ρ) .

In the Hilbert space of any quantum system a bounded or unbounded self-adjoint operator X is called an *observable* of the system. Suppose $\mathcal{F}_{\mathbb{R}}$ is the Borel σ -algebra of \mathbb{R} and $P^X(\cdot)$ is the spectral measure of X on $\mathcal{F}_{\mathbb{R}}$. Then the quantity $\text{Tr } \rho P^X(E)$, $E \in \mathcal{F}_{\mathbb{R}}$ is interpreted as the probability that the observable takes a value in E in the state ρ . Thus $\text{Tr } \rho P^X(\cdot)$ is the distribution of X in the state ρ . Such an interpretation enables the computation of all moments of X . Indeed, the n -th moment of X , if it exists, is denoted by $\langle X^n \rangle$ and is given by

$$\langle X^n \rangle = \text{Tr } X^n \rho.$$

If X, Y are two observables such that $XY + YX$ is also an observable then the covariance between X and Y in the state ρ is denoted by $\text{Cov}(X, Y)$ and is defined as

$$\text{Cov}(X, Y) = \left\langle \frac{1}{2}(XY + YX) \right\rangle - \langle X \rangle \langle Y \rangle.$$

The quantity $\text{Cov}(X, X)$ is called the variance of X . If X_1, X_2, \dots, X_n are observables with well-defined covariance between X_i and X_j for all i, j then the $n \times n$ positive matrix

$$\Sigma_n = \Sigma_n(X_1, \dots, X_n) = [[\text{Cov}(X_i, X_j)]]$$

is called the *covariance matrix* of the observables (X_1, X_2, \dots, X_n) in the state ρ .

Consider a composite quantum system (\mathcal{H}, ρ) where $\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \dots \otimes \mathcal{H}_n$. If the set $\{1, 2, \dots, n\} = E \cup F$ with $E \cap F = \emptyset$, $E \neq \emptyset$, $F \neq \emptyset$ then \mathcal{H} can be viewed as the tensor product

$$\mathcal{H} = \mathcal{H}_E \otimes \mathcal{H}_F$$

and an observable in \mathcal{H}_E can be looked upon as the observable $X_E \otimes I_F$ in \mathcal{H} with I_F being the identity operator in \mathcal{H}_F . We call $X_E \otimes I_F$ the *ampliation* of X_E in \mathcal{H} and denote it by the same symbol X_E . If ρ_E is the E -marginal of ρ in \mathcal{H}_E then

$$\langle X_E \rangle = \text{Tr } X_E \rho_E = \text{Tr } X_E \rho = \langle X_E \otimes I_F \rangle.$$

We now introduce the notion of a quantum process over a countable index set D . Let $\{\mathcal{H}_a : a \in D\}$ be a family of Hilbert spaces. Denote by \mathcal{S}_D the set of all finite sequences of distinct elements from D . Suppose $\rho_{a_1, a_2, \dots, a_n}$ is a density operator in $\mathcal{H}_{a_1, a_2, \dots, a_n}$ as in (1.1) for each (a_1, a_2, \dots, a_n) in \mathcal{S}_D satisfying the following properties:

- (1) If $\{a_1, a_2, \dots, a_n\} = \{b_1, b_2, \dots, b_n\}$ as sets and π is a permutation of $\{1, 2, \dots, n\}$ such that $a_{\pi(j)} = b_j$, $\forall j$ and

$$U_\pi : \mathcal{H}_{a_1, a_2, \dots, a_n} \rightarrow \mathcal{H}_{b_1, b_2, \dots, b_n}$$

is the natural Hilbert space isomorphism induced by π then

$$\rho_{b_1, b_2, \dots, b_n} = U_\pi \rho_{a_1, a_2, \dots, a_n} U_\pi^{-1}.$$

- (2) The $\{a_{n+1}\}$ -marginal of $\rho_{a_1, a_2, \dots, a_{n+1}}$ is equal to $\rho_{a_1, a_2, \dots, a_n}$ for all $(a_1, a_2, \dots, a_{n+1}) \in \mathcal{S}_D$, $n = 1, 2, \dots$.

Then we say that $\{\rho_{a_1, a_2, \dots, a_n}, (a_1, \dots, a_n) \in \mathcal{S}_D\}$ is a consistent family of states. The family $\boldsymbol{\rho} = \{(\mathcal{H}_{a_1, \dots, a_n}, \rho_{a_1, \dots, a_n}), (a_1, \dots, a_n) \in \mathcal{S}_D\}$ of finite-partite quantum systems is called a *quantum process* over D .

One obtains interesting examples of discrete ‘time’ quantum processes with $D = \mathbb{Z}, \mathbb{Z}^d$ or a general discrete abelian group. When D is \mathbb{Z} , the element a in \mathbb{Z} can be interpreted as time. In general, a in D is interpreted as site.

Suppose $D = \{0, 1, 2, \dots\}$ and $\mathcal{H}_{[n]} = \mathcal{H}_0 \otimes \mathcal{H}_1 \otimes \dots \otimes \mathcal{H}_n$. Let $\rho_{[n]}$ be a density operator in $\mathcal{H}_{[n]}$ such that $\rho_{[n-1]}$ is the marginal in $\mathcal{H}_{[n-1]}$ obtained by tracing out $\rho_{[n]}$ over \mathcal{H}_n for each n . Then $\{(\mathcal{H}_{[n]}, \rho_{[n]}) : n = 0, 1, 2, \dots\}$ yields a quantum process. Denote by $\mathcal{B}_{[n]}$ the C* algebra of all bounded operators in $\mathcal{H}_{[n]}$. Then there is a natural C* embedding $\phi_n : \mathcal{B}_{[n]} \rightarrow \mathcal{B}_{[n+1]}$ with the property

$$\phi_n(X) = X \otimes I, \quad X \in \mathcal{B}_{[n]},$$

where I is the identity operator in $\mathcal{H}_{[n+1]}$. This enables the construction of an inductive limit C* algebra \mathcal{B}_∞ with a C* embedding $\pi_n : \mathcal{B}_{[n]} \rightarrow \mathcal{B}_\infty$ such that the sequence $\{\pi_n(\mathcal{B}_{[n]})\}$ is increasing in n and $\bigcup_n \pi_n(\mathcal{B}_{[n]})$ is dense in \mathcal{B}_∞ . This yields a normalized positive linear functional ω in \mathcal{B}_∞ such that

$$\omega(\pi_n(X)) = \rho_{[n]}(X), \quad X \in \mathcal{B}_{[n]}, \quad n = 0, 1, 2, \dots$$

In other words $(\mathcal{B}_\infty, \omega)$ is a C* probability space which may be considered as the analogue of Kolmogorov’s measure space constructed from a consistent family of finite dimensional probability distributions. However, there is no limiting Hilbert space in general with a density operator. A similar construction of a C* probability space is possible for a quantum process over any countable index set D .

Definition 2.1. *Suppose D is a countable abelian group with addition operation $+$, $\mathcal{H}_a = \mathcal{H}$ for all $a \in D$, and $\boldsymbol{\rho}$ is a quantum process over D . Then it is said to be strictly stationary or translation invariant if*

$$\rho_{a_1+x, a_2+x, \dots, a_n+x} = \rho_{a_1, a_2, \dots, a_n} \quad \forall x \in D, (a_1, \dots, a_n) \in \mathcal{S}_D.$$

Let $\{\rho_{a_1, \dots, a_n}\}, \{\sigma_{a_1, \dots, a_n}\}, (a_1, \dots, a_n) \in \mathcal{S}_D$ be a pair of consistent families of finite-partite states in $\{\mathcal{H}_{a_1, \dots, a_n}\}$. Then, for any $0 < p < 1$

$$\tau_{a_1, \dots, a_n} = p\rho_{a_1, \dots, a_n} + (1-p)\sigma_{a_1, \dots, a_n}, \quad (a_1, \dots, a_n) \in \mathcal{S}_D$$

yields a consistent family of finite-partite states.

Suppose, $a \mapsto U_a, a \in D$ is any map where U_a is a unitary operator in \mathcal{H}_a for every a . Then

$$\rho'_{a_1, \dots, a_n} = (U_{a_1} \otimes \dots \otimes U_{a_n}) \rho_{a_1, \dots, a_n} (U_{a_1}^\dagger \otimes \dots \otimes U_{a_n}^\dagger), \quad (a_1, \dots, a_n) \in \mathcal{S}_D$$

also yields a consistent family of states. Indeed, this is a consequence of the following proposition.

Proposition 2.1. *Let \mathcal{H}, \mathcal{K} be Hilbert spaces, ρ a state in $\mathcal{H} \otimes \mathcal{K}$, and U, V be unitary operators in \mathcal{H} and \mathcal{K} respectively. Then*

$$\text{Tr}_{\mathcal{K}}(U \otimes V)\rho(U \otimes V)^\dagger = U(\text{Tr}_{\mathcal{K}}\rho)U^\dagger,$$

where $\text{Tr}_{\mathcal{K}}$ is relative trace over \mathcal{K} .

Proof. This is immediate from the fact that relative trace over \mathcal{K} can be computed by using any orthonormal basis $\{e_j\}$ in \mathcal{K} , and if $\{e_j\}$ is one such basis so is $\{V^\dagger e_j\}$. \square

Combining the two elementary remarks above we can construct new quantum processes over D from a given quantum process $\{(\mathcal{H}_{a_1, \dots, a_n}, \rho_{a_1, \dots, a_n}), (a_1, \dots, a_n) \in \mathcal{S}_D\}$ as follows: Start with a probability space $(\Omega, \mathcal{F}, \mathcal{P})$ and a random process $\{U_a(\omega) : a \in D\}$ where $U_a(\omega)$ is a unitary operator in \mathcal{H}_a for every a . Define

$$(2.1) \quad \rho'_{a_1, \dots, a_n} = \int_{\Omega} P(d\omega) (U_{a_1} \otimes \dots \otimes U_{a_n}) \rho_{a_1, \dots, a_n} (U_{a_1} \otimes \dots \otimes U_{a_n})^\dagger.$$

Then $\{\rho'_{a_1, \dots, a_n}, (a_1, \dots, a_n) \in \mathcal{S}_D\}$ is also a consistent family of finite-partite states.

Remark 2.1. *When D is a countable additive abelian group and $\mathcal{H}_a = \mathcal{H}$ for all $a \in D$, ρ is a strictly stationary quantum process and the random process $\{U_a(\omega) : a \in D\}$ is also strictly stationary, then the quantum process ρ' determined by equation (2.1) is also strictly stationary.*

3. MULTI-MODE PROCESSES AND THEIR COVARIANCE KERNELS

We now pass on to the definition of a *k-mode quantum process* over a countable index set D . Let $\mathcal{H}_a = L^2(\mathbb{R}^k)$ for each $a \in D$, where k is a fixed positive integer called the number of modes. We view \mathcal{H}_a as the a -th copy of $L^2(\mathbb{R}^k)$ and introduce the canonical Schrödinger pairs of position and momentum observables q_{aj}, p_{aj} , $1 \leq j \leq k$ given by

$$\begin{aligned} (q_{aj}f)(\mathbf{x}) &= x_j f(\mathbf{x}), \\ (p_{aj}f)(\mathbf{x}) &= \frac{1}{i} \frac{\partial}{\partial x_j} f(\mathbf{x}) \end{aligned}$$

on their respective maximal domains in $L^2(\mathbb{R}^k)$, \mathbf{x} denoting $(x_1, x_2, \dots, x_k) \in \mathbb{R}^k$. We arrange these $2k$ observables as

$$(X_{a1}, X_{a2}, \dots, X_{a2k-1}, X_{a2k}) = (q_{a1}, p_{a1}, \dots, q_{ak}, p_{ak}).$$

Let now ρ be a quantum process over D . Then X_{ar} can be viewed as an amplified observable in $\mathcal{H}_{a_1, a_2, \dots, a_n}$ whenever the element a occurs in the sequence a_1, a_2, \dots, a_n . We assume that all observables which are closures of polynomials of degree not exceeding 2 in $\{X_{ar}, a \in D, r = 1, 2, \dots, 2k\}$ have finite expectations under the process so that X_{ar} and X_{bs} have a well-defined covariance for any $a, b \in D, r, s \in \{1, 2, \dots, 2k\}$. We write for any $a, b \in D$

$$\kappa_{r,s}(a, b) = \text{Cov}(X_{ar}, X_{bs}), \quad r, s \in \{1, 2, \dots, 2k\},$$

where the covariances can be evaluated in any state $\rho_{a_1, a_2, \dots, a_n}$ when both a and b occur in $(a_1, \dots, a_n) \in \mathcal{S}_D$. Indeed, this follows from consistency of the states occurring in the quantum process. We call $\mathcal{K} = [[K(a, b)]]$, $a, b \in D$ the *covariance kernel* of the k -mode quantum process ρ .

If D is an additive abelian group, $\langle X_{aj} \rangle = 0$ for all $a \in D, 1 \leq j \leq 2k$ and $K(a, b) = K(a + c, b + c)$ for all $a, b, c \in D$ we then say that ρ is a mean zero *second order weakly stationary* or simply *weakly stationary k-mode quantum process*. In such a case, there exists a map \tilde{K} from D into the space of $2k \times 2k$ real matrices, such that $K(a, b) = \tilde{K}(b - a)$. This map \tilde{K} is called the *autocovariance map* of the weakly stationary process.

Theorem 3.1. Let $K(a, b) = [[\kappa_{rs}(a, b)]]$, $a, b \in D$ be a family of $2k \times 2k$ real matrices satisfying the following conditions:

$$\kappa_{rs}(a, b) = \kappa_{sr}(b, a), \quad r, s \in \{1, 2, \dots, 2k\}, \quad a, b \in D.$$

Then there exists a k -mode quantum process \mathbf{q} with covariance kernel $K(\cdot, \cdot)$ if and only if for any sequence $(a_1, \dots, a_n) \in \mathcal{S}_D$ the block matrix $[[K(a_i, a_j)]]$ satisfies the matrix inequality

$$(3.1) \quad [[K(a_i, a_j)]] + \frac{\iota}{2} J_{2kn} \geq 0.$$

Proof. Since ρ_{a_1, \dots, a_n} is a kn -mode state and $[[K(a_i, a_j)]]$ is the covariance matrix of the position-momentum observables $(X_{a_1 1}, \dots, X_{a_1 2k}, X_{a_2 1}, \dots, X_{a_2 2k}, \dots, X_{a_n 1}, \dots, X_{a_n 2k})$ in $L^2(\mathbb{R}^{kn})$, necessity is immediate from the uncertainty relation fulfilled by such a covariance matrix [Hol11, ADMS95, Par10]. To prove the converse define ρ_{a_1, \dots, a_n} to be the mean zero kn -mode Gaussian state with covariance matrix $[[K(a_i, a_j)]]$. Then $\{\rho_{a_1, \dots, a_n}\}$ is a consistent family of Gaussian states constituting the required quantum process. \square

Definition 3.1. A kernel

$$\mathcal{K} = [[K(a, b)]], \quad a, b \in D$$

where $K(a, b)$ are real $2k \times 2k$ matrices satisfying the conditions

- (1) $K(a, b)^T = K(b, a)$.
- (2) $[[K(a_i, a_j)]] \geq 0$ for all $(a_1, a_2, \dots, a_n) \in \mathcal{S}_D$

is called a k -mode classical covariance kernel.

If, in addition, the inequality (3.1) is fulfilled, then it is called a (k -mode) quantum covariance kernel.

Corollary 3.1. If \mathcal{K} is a k -mode quantum covariance kernel and \mathcal{C} is a k -mode classical covariance kernel then $\mathcal{K} + \mathcal{C}$ is a k -mode quantum covariance kernel.

Proof. Immediate. \square

Let \mathbf{q} be a k -mode quantum process over D with quantum covariance kernel $\mathcal{K} = [[K(a, b)]]$, $a, b \in D$. Suppose $\mathcal{C} = [[C(a, b)]]$, $a, b \in D$ is the covariance kernel of a real $2k$ -variate classical stochastic process so that the matrix inequalities

$$\sum_{i,j} \alpha_i \alpha_j C(a_i, a_j) \geq 0$$

for all real scalars $\alpha_1, \dots, \alpha_n$, elements $a_1, \dots, a_n \in D$. Then the sum

$$\mathcal{K} + \mathcal{C} = [[K(a, b) + C(a, b)]]$$

is the covariance kernel of a k -mode quantum process σ . We shall now realise such a process σ by an explicit construction which is an interaction between the quantum process \mathbf{q} and a family of unitary conjugations mediated by a classical process with covariance kernel \mathcal{C} .

To this end we start with the 1-mode Hilbert space $L^2(\mathbb{R})$, its Schrödinger position-momentum pair q, p , the associated annihilation-creation pair a, a^\dagger given by $a = 2^{-\frac{1}{2}}(q + ip)$, $a^\dagger = 2^{-\frac{1}{2}}(q - ip)$ and the unitary Weyl (displacement) operators $W(z) = \exp(za^\dagger - \bar{z}a)$, $z \in \mathbb{C}$ satisfying the relations

$$W(z)aW(z)^\dagger = a - z, \quad z \in \mathbb{C}$$

with the convention that z denotes the scalar as well as the operator zI . This leads to the relations

$$(3.2) \quad W(2^{-\frac{1}{2}}z)qW(2^{-\frac{1}{2}}z)^\dagger = q - x,$$

$$(3.3) \quad W(2^{-\frac{1}{2}}z)pW(2^{-\frac{1}{2}}z)^\dagger = p - y,$$

where $x = \operatorname{Re} z$, $y = \operatorname{Im} z$.

Now for $a \in D$, let

$$\begin{aligned} \mathbf{z}_a &= (z_{a1}, z_{a2}, \dots, z_{ak})^T, \\ z_{ar} &= x_{ar} + iy_{ar}, \end{aligned}$$

where $x_{ar} = \operatorname{Re} z_{ar}$, $y_{ar} = \operatorname{Im} z_{ar}$. Viewing $\mathcal{H} = L^2(\mathbb{R}^k)$ as $L^2(\mathbb{R}) \otimes L^2(\mathbb{R}) \otimes \dots \otimes L^2(\mathbb{R})$, k -fold, introduce the k -mode Weyl operators

$$W(\mathbf{z}_a) = W(z_{a1}) \otimes \dots \otimes W(z_{ak}).$$

Then the relations (3.2, 3.3) yield the relations for the operators X_{a1}, \dots, X_{a2k} defined above as

$$(3.4) \quad W(2^{-\frac{1}{2}}\mathbf{z}_a) \begin{bmatrix} X_{a1} \\ X_{a2} \\ \vdots \\ X_{a2k} \end{bmatrix} W(2^{-\frac{1}{2}}\mathbf{z}_a)^\dagger = \begin{bmatrix} X_{a1} - \alpha_{a1} \\ X_{a2} - \alpha_{a2} \\ \vdots \\ X_{a2k} - \alpha_{a2k} \end{bmatrix}$$

where

$$(3.5) \quad (\alpha_{a1}, \alpha_{a2}, \dots, \alpha_{a2k}) = (x_{a1}, y_{a1}, x_{a2}, y_{a2}, \dots, x_{ak}, y_{ak}).$$

Let $(\xi_{a1}, \eta_{a1}, \xi_{a2}, \eta_{a2}, \dots, \xi_{ak}, \eta_{ak})(\omega)$, $\omega \in \Omega$, $a \in D$ be a $2k$ real variate stochastic process defined on a probability space (Ω, \mathcal{F}, P) with zero means and covariance kernel $\mathcal{C} = [[C(a, b)]]$ where

$$(3.6) \quad C(a, b) = \mathbb{E} \begin{bmatrix} \xi_{a1} \\ \eta_{a1} \\ \xi_{a2} \\ \eta_{a2} \\ \vdots \\ \xi_{ak} \\ \eta_{ak} \end{bmatrix} [\xi_{a1}, \eta_{a1}, \xi_{a2}, \eta_{a2}, \dots, \xi_{ak}, \eta_{ak}], \quad a, b \in D.$$

Define the random unitary operators $U_a(\omega)$ in \mathcal{H}_a , $a \in D$ by putting

$$(3.7) \quad \zeta_{ar} = \xi_{ar} + i\eta_{ar}, \quad r = 1, 2, \dots, k$$

$$(3.8) \quad U_a(\omega) = W(2^{-\frac{1}{2}}\zeta_a(\omega))^\dagger$$

where $\zeta_a = (\zeta_{a1}, \dots, \zeta_{ak}) \in \mathbb{C}^k$. By following the remarks in §2 equation (2.1), define the k -mode quantum process σ by

$$(3.9) \quad \sigma_{a_1, \dots, a_n} = \int_{\Omega} P(d\omega) U_{a_1}(\omega) \otimes \dots \otimes U_{a_n}(\omega) \rho_{a_1, \dots, a_n} U_{a_1}(\omega)^\dagger \otimes \dots \otimes U_{a_n}(\omega)^\dagger.$$

Then we have the following theorem:

Theorem 3.2. *The covariance kernel of the σ process determined by the finite-partite states (3.9) is equal to $\mathcal{K} + \mathcal{C}$.*

Proof. Consider the observable X_{aj} . Its expectation under the σ process is given by

$$\begin{aligned}\mathrm{Tr} X_{ar} \sigma_a &= \int P(d\omega) \mathrm{Tr} U_a(\omega)^\dagger X_{ar} U_a(\omega) \rho_a \\ &= \int P(d\omega) \mathrm{Tr} (X_{ar} - \gamma_{ar}) \rho_a\end{aligned}$$

where

$$(\gamma_{a1}(\omega), \dots, \gamma_{a2k}(\omega)) = (\xi_{a1}(\omega), \eta_{a1}(\omega), \dots, \xi_{ak}(\omega), \eta_{ak}(\omega)).$$

Since the classical γ process has mean $\mathbf{0}$ we have

$$(3.10) \quad \langle X_{ar} \rangle_\sigma = \langle X_{ar} \rangle_\rho.$$

Going to second order moments

$$\begin{aligned}\mathrm{Tr} X_{ar} X_{bs} \sigma_{ab} &= \int P(d\omega) \mathrm{Tr} X_{ar} X_{bs} U_a(\omega) \otimes U_b(\omega) \rho_{ab} U_a(\omega)^\dagger \otimes U_b(\omega)^\dagger \\ &= \int P(d\omega) W(2^{-\frac{1}{2}} \zeta_a) \otimes W(2^{-\frac{1}{2}} \zeta_b) X_{ar} X_{bs} W(2^{-\frac{1}{2}} \zeta_a)^\dagger \otimes W(2^{-\frac{1}{2}} \zeta_b)^\dagger \rho_{ab} \\ &= \int P(d\omega) \mathrm{Tr} (X_{ar} - \gamma_{ar}) (X_{bs} - \gamma_{bs}) \rho_{ab} \\ &= \int P(d\omega) \left[\langle X_{ar} X_{bs} \rangle_\rho + \gamma_{ar}(\omega) \gamma_{bs}(\omega) - \gamma_{ar}(\omega) \langle X_{bs} \rangle_\rho - \gamma_{bs}(\omega) \langle X_{ar} \rangle_\rho \right] \\ &= \langle X_{ar} X_{bs} \rangle_\rho + C_{rs}(a, b).\end{aligned}$$

Let $a \neq b$. Then

$$\begin{aligned}\mathrm{Cov}_\sigma(X_{ar}, X_{bs}) &= \langle X_{ar} X_{bs} \rangle_\sigma - \langle X_{ar} \rangle_\sigma \langle X_{bs} \rangle_\sigma \\ &= \langle X_{ar} X_{bs} \rangle_\rho + C_{rs}(a, b) - \langle X_{ar} \rangle_\rho \langle X_{bs} \rangle_\rho \\ &= K_{rs}(a, b) + C_{rs}(a, b).\end{aligned}$$

Let $a = b$. Then $C_{rs}(a, b) = C_{rs}(a, a) = C_{sr}(a, a)$.

$$\begin{aligned}\mathrm{Cov}_\sigma(X_{ar}, X_{as}) &= \left\langle \frac{1}{2} (X_{ar} X_{as} + X_{as} X_{ar}) \right\rangle_\sigma - \langle X_{ar} \rangle_\rho \langle X_{as} \rangle_\rho \\ &= \left\langle \frac{1}{2} (X_{ar} X_{as} + X_{as} X_{ar}) \right\rangle_\rho + C_{rs}(a, a) - \langle X_{ar} \rangle_\rho \langle X_{as} \rangle_\rho \\ &= \mathrm{Cov}_\rho(X_{ar}, X_{as}) + C_{rs}(a, a) \\ &= K_{rs}(a, a) + C_{rs}(a, a).\end{aligned}$$

□

Remark 3.1. If ϱ is a weakly stationary quantum process, and $(\xi_{a_1}, \eta_{a_1}, \dots, \xi_{a_k}, \eta_{a_k})$ is a weakly stationary classical process with mean $\mathbf{0}$ such that

$$\begin{aligned} K(a, b) &= \tilde{K}(b - a), \quad a, b \in D, \\ C(a, b) &= \tilde{C}(b - a), \end{aligned}$$

then σ is also a weakly stationary quantum process.

Remark 3.2. If ϱ is a Gaussian process then so is σ .

4. THE KMW SPECTRUM OF A WEAKLY STATIONARY k -MODE QUANTUM PROCESS

Let ϱ be a weakly stationary k -mode quantum process over a countable discrete additive abelian group D , with autocovariance map \tilde{K} . Let \hat{D} be the compact dual multiplicative group of all characters of D . Denote by \mathcal{F} the Borel σ -algebra on \hat{D} .

Define

$$(4.1) \quad L(a) = \tilde{K}(a) + \frac{i}{2} \mathbb{1}_{\{0\}}(a) J_{2k}, \quad a \in D.$$

Then Theorem 3.1 yields the following proposition.

Proposition 4.1. A real $2k \times 2k$ matrix-valued map \tilde{K} is the autocovariance map of a second order weakly stationary k -mode quantum process if and only if the associated map L defined by (4.1) satisfies the following matrix inequalities

$$[[L(a_s - a_r)]] \geq 0, \quad r, s \in \{1, 2, \dots, n\}$$

for all $(a_1, a_2, \dots, a_n) \in \mathcal{S}_D$, $n = 1, 2, \dots$.

Proof. Immediate. □

Theorem 4.1. A real $2k \times 2k$ matrix-valued map \tilde{K} on D is the autocovariance map of a second order weakly stationary k -mode quantum process on D if and only if there exists a $2k \times 2k$ Hermitian positive matrix-valued measure Φ on (\hat{D}, \mathcal{F}) satisfying the following conditions:

- (1) $\Phi(\hat{D}) = \tilde{K}(0)$,
- (2) $\tilde{K}(a) = \int_{\hat{D}} \chi(a) \Phi(d\chi)$,
- (3) $\Phi(S) + \frac{i}{2} \lambda(S) J_{2k} \geq 0$, $\forall S \in \mathcal{F}$; where λ is the normalized Haar measure of the compact group \hat{D} . In particular, λ is absolutely continuous with respect to $\text{Tr } \Phi$.
- (4) $\Phi(S^{-1}) = \Phi(S) = \Phi(S)^T$, $\forall S \in \mathcal{F}$.

Proof. Let \tilde{K} be the autocovariance map of a weakly stationary k -mode process. Define L by (4.1). By Proposition 4.1 the matrices $[[L(a_s - a_r)]]$, $r, s \in \{1, 2, \dots, n\}$, $(a_1, a_2, \dots, a_n) \in \mathcal{S}_D$ are positive. Hence for any vector $\mathbf{u} \in \mathbb{C}^{2k}$, the function

$$(4.2) \quad \psi_{\mathbf{u}}(a) = \mathbf{u}^\dagger L(a) \mathbf{u}, \quad a \in D$$

is positive definite on the abelian group D in the sense of Bochner. By Bochner's theorem there exists a totally finite measure $\nu_{\mathbf{u}}$ on \mathcal{F} satisfying the relations

$$(4.3) \quad \psi_{\mathbf{u}}(a) = \int_{\widehat{D}} \chi(a) \nu_{\mathbf{u}}(d\chi), \quad a \in D,$$

$$(4.4) \quad \begin{aligned} \psi_{\mathbf{u}}(0) &= \mathbf{u}^\dagger L(0) \mathbf{u} \\ &= \mathbf{u}^\dagger \left(\widetilde{K}(0) + \frac{i}{2} J_{2k} \right) \mathbf{u}, \quad \mathbf{u} \in \mathbb{C}^{2k}. \end{aligned}$$

By (4.2) the left hand side of (4.3) is a quadratic form in \mathbf{u} for each fixed a in D . By the bijective correspondence between totally finite measures on \widehat{D} and their Fourier transforms on D it follows that there exists a $2k \times 2k$ Hermitian positive matrix-valued measure Ψ on \mathcal{F} such that

$$(4.5) \quad L(a) = \int_{\widehat{D}} \chi(a) \Psi(d\chi), \quad a \in D$$

$$(4.6) \quad L(0) = \widetilde{K}(0) + \frac{i}{2} J_{2k} = \Psi(\widehat{D}) \geq 0.$$

Now define

$$\phi_{\mathbf{u}}(a) = \mathbf{u}^\dagger \widetilde{K}(a) \mathbf{u}, \quad a \in D, \quad \mathbf{u} \in \mathbb{C}^{2k}.$$

By (4.1), $\widetilde{K}(a) = \operatorname{Re} L(a)$ and hence $[[\widetilde{K}(a_s - a_r)]] \geq 0$ for any $(a_1, a_2, \dots, a_n) \in \mathcal{S}_D$. In other words, $\phi_{\mathbf{u}}$ is also a positive definite function on D and by the same arguments as employed for L we have the relations

$$(4.7) \quad \widetilde{K}(a) = \int_{\widehat{D}} \chi(a) \Phi(d\chi),$$

$$(4.8) \quad \widetilde{K}(0) = \Phi(\widehat{D}),$$

where Φ is again a $2k \times 2k$ Hermitian positive matrix-valued measure on \mathcal{F} .

By (4.1)

$$(4.9) \quad \begin{aligned} L(a) - \widetilde{K}(a) &= \frac{i}{2} \mathbb{1}_{\{0\}}(a) J_{2k} \\ &= \left[\frac{i}{2} \int_{\widehat{D}} \chi(a) \lambda(d\chi) \right] J_{2k}, \quad a \in D \end{aligned}$$

Subtracting (4.7) from (4.5) and using (4.9) we have

$$\int_{\widehat{D}} \chi(a) (\Psi - \Phi)(d\chi) = \left[\frac{i}{2} \int_{\widehat{D}} \chi(a) \lambda(d\chi) \right] J_{2k}$$

for all $a \in D$. Thus by uniqueness of Fourier transform we have

$$\Psi(S) - \Phi(S) = \frac{i}{2} \lambda(S) J_{2k}, \quad \forall S \in \mathcal{F}.$$

Thus

$$(4.10) \quad \Phi(S) + \frac{i}{2} \lambda(S) J_{2k} \geq 0, \quad \forall S \in \mathcal{F}.$$

Now property (1) follows from (4.8), property (2) from (4.7), and property (3) from (4.10). If $\text{Tr } \Phi(S) = 0$ then $\Phi(S) = 0$ and (4.10) implies

$$\frac{i}{2} \lambda(S) J_{2k} \geq 0$$

and this is positive only if $\lambda(S) = 0$. In other words $\lambda \ll \text{Tr } \Phi$.

To prove property (4) of Φ we introduce the map $\tau : \widehat{D} \rightarrow \widehat{D}$, $\tau(\chi) = \bar{\chi} = \chi^{-1}$ and observe that

$$\begin{aligned} \tilde{K}(a) = \overline{\tilde{K}(a)} &= \int_{\widehat{D}} \overline{\chi(a)} \bar{\Phi}(d\chi) \\ &= \int_{\widehat{D}} \chi(a) \bar{\Phi}\tau^{-1}(d\chi) \\ &= \int_{\widehat{D}} \chi(a) \Phi(d\chi). \end{aligned}$$

Thus

$$\bar{\Phi}\tau^{-1} = \Phi,$$

or

$$\Phi(S^{-1}) = \overline{\Phi(S)} = \Phi^T(S), \quad \forall S \in \mathcal{F}.$$

This completes the proof of necessity. To prove sufficiency consider a $2k \times 2k$ Hermitian positive matrix-valued measure Φ satisfying properties (3) and (4) of the theorem. Define

$$\tilde{K}(a) = \int_{\widehat{D}} \chi(a) \Phi(d\chi).$$

Property (3) implies that the function $L(a)$, $a \in D$ defined by

$$\begin{aligned} L(a) &= \tilde{K}(a) + \frac{i}{2} \mathbb{1}_{\{0\}}(a) J_{2k} \\ &= \int_{\widehat{D}} \chi(a) \left(\Phi + \frac{i}{2} \lambda J_{2k} \right) (d\chi) \end{aligned}$$

satisfies the matrix inequalities $[[L(a_s - a_r)]] \geq 0$ for any sequence $(a_1, \dots, a_n) \in \mathcal{S}_D$. By Proposition 4.1, \tilde{K} is the autocovariance function of a strictly stationary mean zero k -mode quantum Gaussian process. \square

Remark 4.1. *As already described in the introduction, we call equation (2) in Theorem 4.1, the spectral representation of the autocovariance map \tilde{K} and say that Φ is the Kolmogorov-Wiener-Masani (KWM) spectrum of the k -mode weakly stationary quantum process. Theorem 4.1 enables us to construct a whole class of examples of KWM spectra and hence autocovariance maps as follows. Choose and fix any Borel map $\chi \mapsto M(\chi)$, $\chi \in \widehat{D}$ where $M(\chi)$ is a k -mode quantum covariance matrix of order $2k$, so that*

$$M(\chi) + \frac{i}{2} J_{2k} \geq 0, \quad \text{for every } \chi \in D.$$

Assume that $M(\cdot)$ is integrable with respect to the normalised Haar measure λ on \widehat{D} . Let Ψ be any totally finite positive Hermitian $2k \times 2k$ matrix-valued measure on $(\widehat{D}, \mathcal{F})$ satisfying the

conjugate symmetry condition $\Psi(S^{-1}) = \overline{\Psi(S)}$ for any Borel set $S \subset \widehat{D}$. Define

$$\Phi(S) = \int_{\widehat{D}} M(\chi) \lambda(d\chi) + \Psi(S), \quad S \in \mathcal{F}.$$

Then by Theorem 4.1, Φ is the KWM spectrum of a stationary quantum Gaussian process over D with autocovariance map \widetilde{K} given by equation (2) of the theorem.

Remark 4.2. The second part of property (3) of Φ in Theorem 4.1 implies that $\lambda(S) = 0$ whenever $\text{Tr } \Phi(S) = 0$. In other words the KWM spectrum of a weakly stationary k -mode quantum process over D cannot admit a gap of positive Haar measure in \widehat{D} . For example, when $D = \mathbb{Z}$ and \widehat{D} is identified with $[0, 2\pi]$, the KWM spectrum of a stationary k -mode quantum Gaussian process over \mathbb{Z} cannot admit an interval gap.

Remark 4.3. In Theorem 4.1, express the KWM spectrum Φ as

$$\Phi = [[\phi_{rs}]], \quad r, s \in \{1, 2, \dots, 2k\}$$

and write

$$\begin{aligned} \Phi_q &= [[\phi_{2i-1, 2j-1}]], \quad i, j \in \{1, 2, \dots, k\} \\ \Phi_p &= [[\phi_{2i, 2j}]], \quad i, j \in \{1, 2, \dots, k\}. \end{aligned}$$

In the inductive limit C^* probability space $(\mathcal{B}_\infty, \omega)$ associated with the process \mathfrak{q} described in §2 the commuting family of position observables $\{q_{ar} : a \in D, r \in \{1, 2, \dots, k\}\}$ affiliated to \mathcal{B}_∞ execute a classical weakly stationary process with spectrum Φ_q . A similar property holds for the family $\{p_{ar} : a \in D, r \in \{1, 2, \dots, k\}\}$.

Remark 4.4. Following [PS15] one can introduce the observable

$$N_{aj} = \frac{1}{2} (q_{aj}^2 + p_{aj}^2 - 1), \quad 1 \leq j \leq k,$$

which is the number of particles (photons) in the j -th mode at the site a . If the underlying process \mathfrak{q} is Gaussian with mean $\mathbf{0}$ then

$$\langle N_{aj} \rangle = \frac{1}{2} \left\{ \phi_{2j-1, 2j-1}(\widehat{D}) + \phi_{2j, 2j}(\widehat{D}) - \frac{1}{2} \right\}.$$

This is a consequence of property (1) of Φ in the Theorem 4.1 and Corollary 3.1, equation (3.4) [PS15].

This shows that the relationships between photon numbers and KWM spectrum need a deeper exploration.

REFERENCES

- [ADMS95] Arvind, B. Dutta, N. Mukunda, and R. Simon. The real symplectic groups in quantum mechanics and optics. *Pramana*, 45:471–497, 1995.
- [Hol11] Alexander Holevo. *Probabilistic and statistical aspects of quantum theory*, volume 1 of *Quaderni. Monographs*. Edizioni della Normale, Pisa, second edition, 2011. With a foreword from the second Russian edition by K. A. Valiev.
- [Kol41a] A. Kolmogoroff. Interpolation und Extrapolation von stationären zufälligen Folgen. *Bull. Acad. Sci. URSS Sér. Math. [Izvestia Akad. Nauk. SSSR]*, 5:3–14, 1941.

- [Kol41b] A. N. Kolmogoroff. Stationary sequences in Hilbert's space. *Bolletín Moskovskogo Gosudarstvenogo Universiteta. Matematika*, 2:40pp, 1941.
- [Kol74] A. N. Kolmogorov. *Osnovnye ponyatiya teorii veroyatnostei*. Izdat. "Nauka", Moscow, second edition, 1974. Probability Theory and Mathematical Statistics Series, Vol. 16.
- [Par10] K. R. Parthasarathy. What is a Gaussian state? *Commun. Stoch. Anal.*, 4(2):143–160, 2010.
- [PS15] K. R. Parthasarathy and Ritabrata Sengupta. From particle counting to Gaussian tomography. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 18(4):1550023, 21, Dec 2015.
- [Wie49] Norbert Wiener. *Extrapolation, Interpolation, and Smoothing of Stationary Time Series. With Engineering Applications*. The Technology Press of the Massachusetts Institute of Technology, Cambridge, Mass; John Wiley & Sons, Inc., New York, N. Y.; Chapman & Hall, Ltd., London, 1949.
- [WM57] N. Wiener and P. Masani. The prediction theory of multivariate stochastic processes. I. The regularity condition. *Acta Math.*, 98:111–150, 1957.
- [WM58] N. Wiener and P. Masani. The prediction theory of multivariate stochastic processes. II. The linear predictor. *Acta Math.*, 99:93–137, 1958.

THEORETICAL STATISTICS AND MATHEMATICS UNIT, INDIAN STATISTICAL INSTITUTE, DELHI CENTRE, 7
S J S SANSANWAL MARG, NEW DELHI 110 016, INDIA
E-mail address, K R Parthasarathy: krp@isid.ac.in

THEORETICAL STATISTICS AND MATHEMATICS UNIT, INDIAN STATISTICAL INSTITUTE, DELHI CENTRE, 7
S J S SANSANWAL MARG, NEW DELHI 110 016, INDIA
E-mail address, Ritabrata Sengupta: rb@isid.ac.in