

## Homework 7 (Class Test on 06 November)

### 0. Practice Problems from the Textbook:

Exercises **17.1, 17.3, 18.2, 18.3, 18.5** and **19.14** (only for **18.2, 18.3, 18.5**).

1. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function on  $\mathbb{R}$  satisfying the property

$$f(x) < f(0) \text{ for all } x > 1.$$

Now consider the following constrained maximization problem:

$$\left. \begin{array}{l} \text{Maximize } f(x) \\ \text{subject to } x \geq 0. \end{array} \right\}$$

Does this problem have a solution? Explain carefully.

### 2. [The Method of Least Squares]

- Suppose we are given  $n$  points  $(x_i, y_i)$ ,  $i = 1, 2, \dots, n$  in  $\mathbb{R}^2$ . Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be given by  $f(x) = ax + b$ , for all  $x \in \mathbb{R}$ . We wish to find a function  $f$  (that is, we want to choose  $a \in \mathbb{R}$  and  $b \in \mathbb{R}$ ) such that the quantity  $\sum_{i=1}^n [f(x_i) - y_i]^2$  is minimized.
- We can set up the problem as an *unconstrained maximization problem* as follows. Define  $F : \mathbb{R}^2 \rightarrow \mathbb{R}$  by

$$F(a, b) = - \sum_{i=1}^n [ax_i + b - y_i]^2.$$

The maximization problem is:

$$\text{Maximize}_{\{a, b\}} F(a, b).$$

- (a) Write down the first-order conditions of the maximization problem.
- (b) Argue clearly that if  $(a^*, b^*)$  satisfies the first-order conditions, then  $(a^*, b^*)$  is a point of *global maximum* of  $F$ .
- (c) Find out  $(a^*, b^*)$  which is the point of *global maximum* of  $F$ .

3. Recall the following theorem.

Theorem (Taylor's Expansion upto Second Order):

Suppose  $A$  is an open convex subset of  $\mathbb{R}^n$ , and  $f : A \rightarrow \mathbb{R}$  is twice continuously differentiable on  $A$ . Suppose  $x^1$  and  $x^2$  are in  $A$ . Then there is  $0 \leq \theta \leq 1$  such that

$$f(x^2) - f(x^1) = (x^2 - x^1) \cdot \nabla f(x^1) + \frac{1}{2} (x^2 - x^1) \cdot H_f(\theta x^1 + (1 - \theta)x^2) \cdot (x^2 - x^1).$$

Suppose  $D$  is an open convex set in  $\mathbb{R}^n$ , and  $g : D \rightarrow \mathbb{R}$  is twice continuously differentiable and *quasiconcave* on  $D$ . Suppose there exists  $x^* \in D$  satisfying

- (i)  $\nabla g(x^*) = 0$ , and
- (ii)  $H_g(x^*)$  is *negative definite*.

- (a) Use the above theorem to prove that  $g$  has a *strict* local maximum at  $x^*$ .
- (b) Prove further that  $x^*$  is a point of *global maximum* of  $g$  on  $D$ .

4. Consider the following constrained maximization problem:

$$\left. \begin{array}{ll} \text{Maximize} & \prod_{i=1}^n x_i \\ \text{subject to} & \sum_{i=1}^n x_i = n, \\ \text{and} & x_i \geq 0, i = 1, 2, \dots, n. \end{array} \right\} \text{(P)}$$

Using Weierstrass theorem we have proved in class that there exists  $x^* \in C$   
 $= \left\{ x \in \mathbb{R}_+^n : \sum_{i=1}^n x_i = n \right\}$  such that  $x^*$  solves (P).

- (a) Apply Lagrange theorem to find  $x^*$ .
- (b) **The Arithmetic Mean-Geometric Mean Inequality:** Let  $a_1, a_2, \dots, a_n$  be

positive real numbers. Their AM is  $\frac{\sum_{i=1}^n a_i}{n}$ , and GM is  $\left( \prod_{i=1}^n a_i \right)^{\frac{1}{n}}$ .

- Use the conclusion of part (a) to prove that  $\text{AM} \geq \text{GM}$ .

5. Let  $c = (c_1, \dots, c_n)$  be a non-zero vector in  $\mathbb{R}^n$ . Consider the following constrained maximization problem:

$$\left. \begin{array}{ll} \text{Maximize} & \sum_{i=1}^n c_i x_i \\ \text{subject to} & \sum_{i=1}^n x_i^2 = 1, \\ & (x_1, \dots, x_n) \in \mathbb{R}^n. \end{array} \right\} (Q)$$

- (a) Show, by using Weierstrass theorem, that there exists  $(x_1^*, \dots, x_n^*) \in \mathbb{R}^n$  which solves (Q).  
 (b) Use Lagrange theorem to show that

$$\sum_{i=1}^n c_i x_i^* = \|c\|. ^1$$

- (c) Let  $p$  and  $q$  be arbitrary non-zero vectors in  $\mathbb{R}^n$ . Using the result in (b) show that

$$|pq| \leq \|p\| \|q\|. \text{ (Cauchy-Schwarz Inequality)}$$

---

<sup>1</sup>Recall that  $\|y\| = \sqrt{y_1^2 + y_2^2 + \dots + y_n^2}$ .

6. In the lectures we have also discussed (without a proof) the following Theorem. In this question we will prove this Theorem.

- Let  $A \subset \mathbb{R}^n$  be an open convex set, and  $f : A \rightarrow \mathbb{R}$  is twice continuously differentiable on  $A$ . The *bordered Hessian matrix* of  $f$  at  $x \in A$  is denoted by  $G_f(x)$  and is defined by the following  $(n+1) \times (n+1)$  matrix:

$$G_f(x) = \begin{pmatrix} 0 & \nabla f(x) \\ \nabla f(x) & H_f(x) \end{pmatrix}.$$

We denote the  $(k+1)$ th leading principal minor of  $G_f(x)$  by  $|G_f(x; k)|$ , where  $k = 1, 2, \dots, n$ .

- **Theorem:** Suppose  $A \subset \mathbb{R}^n$  is an open convex set, and  $f : A \rightarrow \mathbb{R}$  is twice continuously differentiable on  $A$ . If  $(-1)^k |G_f(x; k)| > 0$  for  $x \in A$ , and  $k = 1, 2, \dots, n$ , then  $f$  is quasi-concave on  $A$ .

- We will prove this Theorem through the following three steps. Fix an arbitrary point  $x^* \in A$ , and define  $g$  by  $g(x) = \nabla f(x^*) \cdot (x^* - x)$ , for  $x \in \mathbb{R}^n$ .
- (a) **Step I:** Given the premise of the Theorem, prove that  $x^*$  is a *local maximum* of  $f$  on the constraint set  $C'[x^*] = \{x \in A \mid g(x) = 0\}$  by showing (i) that the constraint qualification is met at  $x^*$ , (ii) that there exists  $\lambda^* \in \mathbb{R}$  such that  $(x^*, \lambda^*)$  meets the first-order conditions of the Lagrange Theorem, and (iii) that  $(x^*, \lambda^*)$  also meets second-order sufficient conditions for a local maximum.
- **Step II:** It can also be shown that (*you do not have to show this*)  $x^*$  is actually a *global maximum* of  $f$  on the constraint set  $C[x^*] = \{x \in A \mid g(x) \geq 0\}$ .
- (b) **Step III:** Prove the following proposition:

If a  $C^1$  function  $F : A \rightarrow \mathbb{R}$  satisfies the following property for all  $z \in A$ :

$$\nabla F(z) \cdot (z - w) \geq 0 \text{ implies } F(w) \leq F(z), \text{ for all } w \in A,$$

then  $F$  is quasi-concave on  $A$ .

Using this proposition, complete the proof that given the premise of the Theorem,  $f$  is quasi-concave on  $A$ .